

平成20年度 SO₄²⁻濃度 月・年平均值

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均值	最大値 ²⁾	最小値 ²⁾
利尻	24.7	26.4	27.1	11.3	14.0	16.3	25.3	81.6	32.9	56.3	28.2	50.0	30.7	853	0.6
札幌	43.6	18.7	22.1	15.5	21.7	14.9	23.9	36.1	40.5	41.0	39.0	46.5	31.8	170	5.8
落石岬	67.2	27.5	12.2	14.6	9.4	3.2	33.9	40.6	22.4	48.1	*	36.2	21.1	653	1.2
竜飛岬	31.0	44.8	18.4	12.8	12.7	38.1	44.3	33.6	32.8	89.3	51.3	46.2	28.6	914	3.0
八幡平	36.4	24.0	4.4	12.3	5.3	15.3	15.0	28.5	26.4	18.3	19.3	17.5	15.0	299	0.3
篔岳	21.0	11.6	9.3	24.5	7.2	7.3	12.6	10.7	32.1	9.0	26.3	12.0	12.8	253	0.3
尾花沢	15.7	8.7	2.5	21.0	4.6	14.1	11.4	44.1	33.0	13.6	29.4	18.1	16.1	61.4	1.4
筑波	16.3	10.0	18.0	29.7	7.0	7.7	10.9	10.8	7.9	9.4	19.9	13.7	11.8	141	0.6
赤城	7.9	5.8	3.2	10.2	4.3	5.2	9.2	6.1	3.9	3.6	22.1	8.8	5.9	60.2	0.3
小笠原	20.8	11.5	3.8	4.9	4.9	3.4	4.1	8.1	10.5	7.5	11.1	24.9	8.7	207	0.9
佐渡関岬	30.7	13.9	8.3	16.7	11.5	32.6	35.6	47.3	125	51.2	68.9	18.8	30.1	706	3.9
新潟巻	29.7	14.9	5.9	20.5	12.6	18.2	24.2	37.8	43.5	43.1	48.2	29.1	28.3	384	3.0
八方尾根	19.4	6.0	5.5	13.1	5.1	10.5	15.6	7.7	20.8	17.0	19.4	27.3	11.9	305	1.7
越前岬	16.3	6.9	10.4	10.3	16.8	11.3	20.0	74.2	56.3	58.4	26.0	26.4	28.9	377	0.0
伊自良湖	18.7	16.6	12.3	18.2	23.9	27.2	10.6	14.8	31.8	17.5	28.6	13.9	18.1	68.1	1.4
犬山	15.2	11.4	13.1	22.0	16.5	15.1	11.2	17.2	29.9	15.4	21.9	15.1	15.5	166	0.9
潮岬	23.2	11.8	12.8	14.7	13.5	9.1	10.7	12.0	22.8	15.6	35.6	40.7	16.5	180	1.8
京都八幡	17.6	7.6	9.0	14.2	20.1	8.7	11.5	16.3	20.0	11.4	14.7	9.6	12.1	98.3	1.8
尼崎	14.6	8.0	13.3	19.0	20.8	11.3	10.9	13.9	26.0	10.0	16.9	17.6	14.4	99.0	1.1
倉橋島	12.1	15.3	14.6	24.7	13.1	8.5	9.5	18.9	19.8	14.7	18.6	18.1	14.5	129	3.3
隠岐	19.4	18.0	9.5	15.4	13.7	13.4	19.3	43.2	75.2	83.2	69.6	51.3	37.8	675	3.7
蟠竜湖	18.8	9.4	9.4	29.8	14.2	16.7	28.2	37.5	35.1	50.0	23.6	30.5	24.4	115	3.0
禰原	10.9	7.9	7.7	8.3	8.0	4.1	5.5	12.9	24.4	28.8	15.4	11.7	10.3	197	0.8
筑後小郡	17.1	13.1	8.6	18.2	14.5	13.1	24.3	42.8	28.5	47.8	23.0	12.9	16.0	232	1.3
大分久住	16.2	11.3	6.6	19.4	22.9	16.8	7.9	20.8	18.1	18.1	21.0	11.3	13.6	94.4	2.3
対馬	26.8	20.7	17.7	25.1	12.5	17.5	22.8	31.4	54.8	22.9	33.5	26.7	22.4	166	1.4
五島	24.7	14.9	11.3	16.4	17.9	9.6	21.1	36.7	34.8	36.9	30.6	38.5	20.7	216	2.9
えびの	16.7	8.9	8.4	13.3	13.7	13.8	9.8	14.7	21.3	21.4	16.5	14.0	12.5	115	0.5
屋久島	19.5	7.9	7.8	7.4	14.9	10.1	24.3	36.6	35.7	39.9	27.2	29.6	17.3	121	4.1
辺戸岬	10.4	7.4	6.9	7.5	10.0	7.5	9.5	13.4	85.0	42.8	25.9	18.0	13.0	927	2.4
東京	13.1	14.3	18.9	30.1	29.6	17.1	17.3	19.0	14.5	10.8	45.5	21.8	19.5	316	2.9
最大値 ¹⁾	67.2	44.8	27.1	30.1	29.6	38.1	44.3	81.6	125	89.3	69.6	51.3	37.8	927	5.8
最小値 ¹⁾	7.9	5.8	2.5	4.9	4.3	3.2	4.1	6.1	3.9	3.6	11.1	8.8	5.9	60.2	0.0
全国平均	21.8	14.0	10.9	16.8	13.5	13.5	17.4	28.0	34.4	30.7	29.2	24.4	18.7		
標準偏差	11.5	8.1	5.8	6.7	6.3	7.8	9.5	18.5	24.3	22.2	14.6	12.7	7.8		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均值の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 nss-SO₄²⁻濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	16.3	24.8	24.7	10.8	9.5	9.3	8.8	15.1	13.3	31.1	15.9	35.8	16.2	286	0.0
札幌	35.7	17.6	21.5	15.3	18.3	12.2	20.6	21.0	23.0	25.4	23.2	21.0	21.1	163	5.7
落石岬	12.5	12.8	8.6	13.6	7.5	2.4	10.4	12.4	7.7	8.2 *		7.8	8.4	115	0.0
竜飛岬	23.8	30.5	15.1	11.9	8.6	10.6	25.0	19.9	18.5	8.2	20.8	19.5	15.2	697	0.0
八幡平	35.2	22.8	4.2	12.2	5.1	10.9	14.2	24.8	23.0	17.3	16.0	15.2	13.6	272	0.3
篔岳	19.7	10.0	8.2	24.1	6.8	7.1	10.7	10.0	17.0	7.9	15.5	10.4	11.1	249	0.2
尾花沢	14.8	8.1	2.4	20.8	4.5	9.5	10.1	39.6	12.8	9.6	21.5	13.2	12.2	48.2	1.4
筑波	13.8	8.9	17.6	28.6	6.4	7.5	9.4	9.9	6.8	8.7	18.3	11.6	10.7	137	0.6
赤城	7.4	5.6	3.1	10.1	4.3	5.1	8.8	5.8	3.6	3.3	21.1	8.2	5.7	58.1	0.2
小笠原	14.7	7.4	2.1	2.9	2.9	2.0	0.9	1.5	1.1	3.0	7.1	16.3	3.9	201	0.0
佐渡関岬	24.6	12.4	7.6	15.8	9.1	24.3	25.1	16.8	29.7	7.1	32.9	11.6	17.0	146	0.0
新潟巻	27.6	14.0	5.7	20.1	10.3	16.5	17.7	12.2	19.3	17.2	36.8	22.4	17.4	132	2.1
八方尾根	18.8	6.0	5.4	13.1	5.0	10.4	15.3	7.1	19.8	16.4	18.9	26.1	11.6	285	1.7
越前岬	15.2	6.2	8.6	9.7	12.8	9.3	17.3	47.4	31.6	16.8	18.3	18.7	18.0	312	0.0
伊自良湖	17.9	16.1	12.2	17.9	23.4	26.8	10.0	12.3	26.8	16.3	26.3	12.5	17.2	66.2	1.3
犬山	14.6	10.9	13.0	21.5	16.2	14.6	10.4	16.0	26.4	14.9	20.2	13.3	14.8	165	0.9
潮岬	13.3	7.1	7.7	10.9	7.6	5.9	7.1	8.6	7.4	11.2	15.2	12.9	9.1	93.2	0.6
京都八幡	17.2	7.3	8.8	13.8	19.8	8.6	10.8	15.4	18.0	11.2	13.3	9.1	11.7	97.6	1.7
尼崎	14.3	7.7	13.1	18.6	20.5	10.9	10.2	13.1	20.7	9.7	14.2	16.1	13.6	87.7	1.0
倉橋島	12.0	15.0	14.3	24.4	12.5	8.5	9.2	18.0	16.3	14.3	17.3	15.3	13.8	129	3.3
隠岐	10.2	16.1	7.5	8.3	7.7	11.8	14.9	17.8	25.4	15.2	36.0	29.5	16.6	230	0.0
蟠竜湖	17.1	8.7	9.0	29.1	13.3	15.4	26.9	29.1	24.6	29.0	21.1	25.6	19.4	71.7	2.9
禰原	10.6	7.6	7.6	8.1	7.7	4.0	5.2	12.3	21.2	24.8	14.4	10.8	9.7	194	0.8
筑後小郡	16.6	12.7	8.2	17.3	13.9	12.7	23.7	39.9	26.0	37.2	21.6	12.5	15.0	159	1.1
大分久住	15.9	11.0	6.5	19.1	22.2	16.7	7.8	20.5	17.3	17.2	20.5	11.0	13.3	93.7	2.2
対馬	25.3	18.4	16.2	23.7	11.3	16.5	21.3	25.4	51.6	19.8	31.2	22.9	20.4	163	1.0
五島	19.3	13.0	9.6	13.8	15.0	8.8	12.8	27.2	24.2	26.1	24.6	20.2	15.6	155	2.1
えびの	16.2	8.8	7.4	12.6	12.9	13.4	9.5	14.3	19.2	18.7	15.8	13.4	11.7	111	0.3
屋久島	16.1	5.3	6.2	5.0	12.5	5.9	22.2	26.9	24.5	26.5	23.5	23.5	13.1	86.2	3.0
辺戸岬	6.1	5.6	4.2	4.0	7.1	4.5	5.0	5.0	20.2	12.7	10.1	7.5	6.0	417	0.0
東京	11.4	13.0	18.5	28.8	28.5	16.8	15.7	18.3	12.9	9.7	42.8	17.2	18.2	309	2.8
最大値 ¹⁾	35.7	30.5	24.7	29.1	28.5	26.8	26.9	47.4	51.6	37.2	42.8	35.8	21.1	697	5.7
最小値 ¹⁾	6.1	5.3	2.1	2.9	2.9	2.0	0.9	1.5	1.1	3.0	7.1	7.5	3.9	48.2	0.0
全国平均	17.2	12.0	9.8	15.7	11.7	10.9	13.5	18.2	19.7	16.0	21.1	16.5	13.6		
標準偏差	6.9	6.1	5.6	7.1	6.3	5.7	6.6	10.5	9.6	8.3	8.1	6.8	4.3		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 NO₃⁻濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	17.1	23.1	18.9	10.9	13.3	6.8	10.3	18.4	14.7	32.4	12.9	48.0	17.0	344	0.5
札幌	37.6	18.2	22.9	19.5	15.0	11.0	14.7	20.2	25.5	30.6	29.3	24.0	23.0	187	6.5
落石岬	12.4	11.5	8.7	11.4	5.4	2.3	13.9	14.0	9.1	2.6	*	5.4	7.6	144	0.0
竜飛岬	29.5	45.6	22.3	11.8	9.3	9.1	31.9	20.1	28.3	23.1	15.2	27.5	18.1	588	2.2
八幡平	31.2	23.1	5.8	12.8	4.9	10.5	11.3	24.2	23.2	12.5	14.4	17.9	13.1	384	0.4
篔岳	21.7	8.2	10.1	27.9	7.0	7.6	11.0	10.9	23.1	12.9	12.1	10.5	12.1	170	0.6
尾花沢	10.1	9.0	4.8	21.3	6.0	9.9	9.7	38.8	10.5	8.4	21.6	13.2	11.9	52.4	2.0
筑波	14.7	11.1	19.8	59.0	8.7	12.5	13.0	12.3	17.5	12.1	21.7	15.8	14.4	435	1.2
赤城	15.8	13.0	8.8	23.1	6.4	16.8	13.2	14.3	8.9	5.9	30.1	10.9	12.6	191	1.0
小笠原	14.0	3.8	2.8	3.1	2.8	2.9	2.1	2.6	2.2	2.5	5.6	13.3	3.8	100	0.6
佐渡関岬	37.8	14.0	11.9	17.8	10.1	19.1	24.8	20.2	58.2	14.5	39.1	8.8	19.4	461	2.0
新潟巻	37.2	15.4	12.3	26.3	11.8	19.5	18.6	12.0	23.6	14.4	29.8	20.4	18.5	225	0.6
八方尾根	19.9	7.5	7.1	14.6	6.2	7.8	12.5	6.9	19.8	13.2	12.7	22.5	10.7	250	2.4
越前岬	15.4	8.1	13.2	10.8	12.4	12.8	22.3	15.8	15.9	19.2	18.3	20.6	15.2	368	0.0
伊自良湖	20.9	18.8	15.9	27.8	29.9	42.6	12.4	23.1	44.1	19.4	30.0	15.1	22.7	87.8	2.7
犬山	17.2	15.5	17.6	41.0	21.1	20.4	16.2	32.6	33.0	18.8	23.7	16.2	20.6	294	1.3
潮岬	13.5	7.4	8.7	14.3	9.1	6.0	6.6	13.1	16.5	8.9	11.2	11.9	9.3	130	0.4
京都八幡	21.5	10.5	13.8	26.6	27.1	15.8	18.7	26.9	20.6	15.4	15.1	10.5	17.2	141	2.9
尼崎	15.4	10.1	19.6	43.0	24.8	21.6	20.5	20.3	25.5	12.3	15.4	13.3	19.3	190	3.9
倉橋島	14.1	14.7	14.2	30.3	14.3	11.8	12.6	23.0	15.9	19.6	15.8	17.9	15.2	166	2.0
隠岐	11.6	14.3	9.7	16.3	10.1	16.5	25.0	24.1	26.4	24.5	37.1	52.4	21.1	319	0.0
幡竜湖	16.1	8.7	13.2	22.2	14.2	22.6	26.9	40.6	26.4	59.5	21.1	36.7	24.8	158	5.3
禰原	9.4	6.1	5.3	7.8	10.1	4.2	4.0	10.6	25.0	32.8	7.6	9.0	8.8	174	0.8
筑後小郡	11.9	11.9	10.3	27.4	14.5	14.8	24.3	46.2	24.7	67.4	15.7	11.6	16.3	273	1.8
大分久住	9.5	6.3	3.4	7.9	13.1	10.6	3.1	8.5	8.8	18.8	12.3	6.7	7.6	112	1.4
対馬	19.3	14.7	18.1	23.3	13.8	21.2	21.3	34.3	48.7	21.6	28.2	22.5	21.0	180	1.2
五島	11.2	11.0	9.0	14.2	15.2	7.2	8.7	24.9	21.5	34.5	22.4	17.6	14.0	309	0.9
えびの	10.3	4.1	5.3	8.5	9.0	6.3	4.7	6.7	12.6	17.7	9.0	8.4	7.2	161	0.6
屋久島	11.6	6.0	5.4	3.0	10.1	4.3	9.5	21.9	14.7	21.6	19.9	17.6	10.1	71.3	1.0
辺戸岬	8.1	6.0	5.6	5.2	8.0	4.0	4.0	5.7	25.4	15.3	12.9	10.9	6.7	305	0.0
東京	15.6	13.3	20.9	48.8	40.3	25.4	14.6	23.9	22.6	12.9	52.3	19.4	23.7	611	2.5
最大値 ¹⁾	37.8	45.6	22.9	59.0	40.3	42.6	31.9	46.2	58.2	67.4	52.3	52.4	24.8	611	6.5
最小値 ¹⁾	8.1	3.8	2.8	3.0	2.8	2.3	2.1	2.6	2.2	2.5	5.6	5.4	3.8	52.4	0.0
全国平均	17.8	12.6	11.8	20.6	13.0	13.0	14.3	19.9	22.4	20.2	20.4	17.9	14.9		
標準偏差	8.5	7.9	6.0	13.4	8.1	8.5	7.6	10.6	11.7	14.1	10.4	10.9	5.7		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 C1⁻濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	162	26.7	42.6	8.8	74.4	123	296	1220	350	453	215	232	259	15700	0.6
札幌	151	20.6	13.4	5.8	73.1	45.5	57.8	293	335	301	290	459	201	911	4.1
落石岬	1030	281	68.5	19.5	34.5	14.9	432	518	271	756	*	564	243	12000	5.0
竜飛岬	123	260	61.8	15.7	78.5	535	356	251	265	1500	555	484	246	12400	1.5
八幡平	23.7	22.2	3.1	2.5	3.0	85.8	13.9	70.6	65.5	18.0	59.2	43.5	26.6	718	0.0
籠岳	23.6	30.2	21.0	7.3	6.7	3.9	35.9	15.9	285	21.5	208	33.3	32.0	3820	1.0
尾花沢	16.5	11.6	2.3	5.2	2.5	88.5	23.9	83.9	396	78.8	152	98.3	77.8	466	0.5
筑波	47.1	21.9	9.8	20.2	10.3	5.1	27.2	14.8	19.5	13.1	32.0	44.7	22.2	522	0.7
赤城	8.8	4.8	2.7	2.7	1.7	3.9	7.2	7.5	7.0	4.8	19.1	11.0	4.7	104	0.4
小笠原	114	78.4	32.1	36.6	38.3	28.5	61.1	121	177	86.0	71.6	166	90.2	2240	6.1
佐渡関岬	109	33.7	14.3	17.2	49.4	157	204	535	1850	768	612	133	236	9240	2.3
新潟巻	43.6	18.4	5.1	11.8	45.5	33.1	128	500	474	504	220	120	212	5440	2.9
八方尾根	12.7	2.3	1.8	2.4	2.3	3.5	7.8	12.0	17.2	15.2	10.3	14.8	6.2	191	0.1
越前岬	20.5	11.3	30.5	9.9	73.1	32.9	51.9	460	458	817	139	141	201	2900	1.4
伊自良湖	12.8	8.6	4.0	8.5	10.4	10.3	12.7	49.3	98.9	25.5	44.4	27.9	18.5	154	2.1
犬山	11.0	7.6	4.8	11.0	6.5	8.9	15.0	21.5	64.3	9.5	32.2	34.2	12.7	173	1.0
潮岬	180	88.4	93.2	64.6	101	59.0	69.5	58.8	276	83.3	386	530	138	2940	4.3
京都八幡	10.2	8.2	5.6	9.4	7.6	5.6	13.4	19.3	38.3	6.7	27.0	10.2	10.8	101	2.6
尼崎	6.9	7.0	5.2	10.5	7.0	7.0	11.8	14.4	100	6.1	50.5	29.9	16.4	259	1.6
倉橋島	5.8	6.7	6.5	8.3	14.2	5.6	12.3	19.3	67.5	10.9	25.8	55.4	15.7	417	0.8
隠岐	178	38.6	38.4	135	112	31.6	85.1	530	957	1300	633	415	409	12300	6.0
幡竜湖	34.0	15.2	10.5	15.9	17.9	26.4	25.2	159	199	396	47.3	91.4	94.3	971	2.9
禰原	6.7	5.2	2.0	3.0	5.5	2.9	4.9	12.8	62.5	78.4	19.7	17.6	12.4	645	0.3
筑後小郡	6.0	5.7	4.4	14.1	9.3	6.2	10.0	56.4	50.5	211	23.5	5.7	17.0	1300	1.0
大分久住	4.1	3.6	2.5	8.4	15.2	6.9	4.3	9.1	20.5	19.0	14.4	5.1	7.2	107	0.3
対馬	28.4	43.7	25.1	25.6	22.6	19.2	26.8	110	61.4	56.8	43.0	70.5	36.8	446	1.2
五島	98.2	38.2	33.6	47.1	54.6	15.3	143	187	201	192	115	369	95.9	2360	1.3
えびの	9.9	3.1	5.1	14.8	16.3	7.0	7.3	9.8	44.0	50.2	15.8	11.9	10.8	694	0.5
屋久島	68.0	60.4	32.0	48.2	56.6	98.0	46.5	224	242	299	81.2	135	95.2	762	17.2
辺戸岬	76.8	32.9	49.9	66.4	53.4	54.0	83.1	157	1170	545	288	196	129	16600	5.2
東京	28.8	25.6	8.0	23.5	23.2	8.1	30.6	16.2	29.7	20.9	65.8	84.8	24.8	601	0.6
最大値 ¹⁾	1030	281	93.2	135	112	535	432	1220	1850	1500	633	564	409	16600	17.2
最小値 ¹⁾	4.1	2.3	1.8	2.4	1.7	2.9	4.3	7.5	7.0	4.8	10.3	5.1	4.7	101	0.0
全国平均	85.5	39.4	20.6	21.9	33.1	49.5	74.3	186	279	279	150	150	96.8		
標準偏差	184	65.3	23.0	27.1	31.6	98.3	107	262	393	390	181	172	104		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 NH₄⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	27.4	31.8	33.5	22.9	26.1	15.4	21.4	36.9	14.6	33.8	15.9	44.4	25.4	430	0.6
札幌	51.8	12.5	27.7	18.9	49.9	18.3	24.0	30.4	32.7	31.6	31.3	34.5	27.3	280	0.0
落石岬	11.8	14.7	7.2	11.9	4.3	0.7	8.8	5.9	3.8	23.6	*	31.0	9.1	358	0.0
竜飛岬	34.2	31.6	13.4	15.3	7.3	8.0	24.5	17.7	18.6	13.0	20.8	19.6	16.3	1070	0.0
八幡平	42.4	29.7	7.1	20.6	5.8	13.6	17.0	26.0	23.3	15.3	14.5	16.8	16.1	417	0.7
篔岳	29.3	11.1	10.8	34.5	8.1	10.5	11.1	10.1	19.2	12.1	7.3	9.2	13.7	241	1.0
尾花沢	22.4	9.6	5.1	28.6	5.5	10.6	10.9	45.2	15.3	9.8	21.6	14.2	14.4	56.7	1.9
筑波	20.8	15.0	24.2	53.5	11.2	15.3	13.7	11.4	13.0	10.8	17.7	20.7	16.3	306	1.3
赤城	14.4	10.4	7.8	31.2	6.1	13.6	13.2	12.5	8.9	3.1	21.9	9.3	11.7	180	0.0
小笠原	16.6	2.4	0.6	2.2	1.3	1.1	0.5	1.2	1.3	2.1	3.8	21.2	3.2	75.6	0.0
佐渡関岬	36.4	10.9	8.6	20.3	8.0	22.9	26.0	18.6	48.1	10.2	43.9	7.0	18.8	484	0.0
新潟巻	34.9	12.7	8.8	29.8	10.0	12.9	16.5	10.2	16.5	12.0	34.0	20.9	16.9	196	1.7
八方尾根	21.3	5.7	7.2	17.9	4.4	5.3	13.7	6.3	21.9	13.5	19.0	23.9	11.3	148	0.0
越前岬	18.6	8.2	10.0	14.2	15.6	10.9	21.2	17.6	17.8	23.7	18.7	21.3	16.3	235	0.0
伊自良湖	19.2	20.9	9.0	24.8	21.4	30.8	8.1	10.2	29.1	14.7	27.3	14.4	18.0	93.7	1.0
犬山	17.9	14.1	12.0	33.2	16.6	18.1	11.9	17.6	32.2	15.7	25.0	16.0	17.4	214	1.7
潮岬	11.5	4.2	7.5	7.7	3.2	2.9	3.0	8.6	7.8	6.0	17.7	12.4	7.1	189	0.1
京都八幡	16.7	8.8	10.7	18.9	25.0	9.9	7.7	16.7	18.9	12.8	16.7	12.1	13.5	139	1.5
尼崎	15.0	6.5	9.8	27.1	29.3	18.1	14.0	15.5	19.4	11.4	16.1	18.7	15.9	115	3.3
倉橋島	8.0	10.9	8.8	24.1	8.0	4.7	5.9	14.5	13.3	13.6	12.0	11.6	9.7	140	0.0
隠岐	11.3	12.4	4.7	8.4	5.7	10.9	14.2	17.9	24.7	20.4	36.8	47.3	17.2	223	0.8
蟠竜湖	13.7	6.7	7.0	28.8	9.2	13.3	17.9	28.6	23.1	37.8	15.0	29.5	18.3	96.8	2.7
禰原	9.4	4.8	3.8	7.9	5.3	2.1	2.8	9.5	16.1	24.9	10.1	12.5	7.6	157	0.0
筑後小郡	22.5	20.1	16.6	38.5	23.7	19.6	22.7	50.1	34.6	54.0	30.7	20.8	23.9	200	1.9
大分久住	18.2	8.6	5.6	14.9	16.3	16.2	3.2	9.5	12.9	17.6	20.7	14.2	11.2	87.3	2.1
対馬	25.2	16.6	13.9	17.1	10.0	10.5	14.8	21.2	46.4	14.5	30.4	21.8	17.7	157	1.7
五島	18.0	11.7	13.5	12.5	15.6	7.0	5.7	20.9	17.4	25.4	20.4	17.5	14.2	158	1.4
えびの	16.7	8.0	8.1	15.8	16.0	8.7	8.5	11.5	13.4	19.1	17.4	14.7	11.6	179	0.0
屋久島	13.4	3.4	4.4	4.0	7.6	2.1	15.4	16.3	22.7	18.7	22.9	22.3	9.8	111	0.6
辺戸岬	8.5	5.8	4.6	2.7	6.4	3.7	4.1	3.6	15.2	14.4	13.0	11.0	6.1	278	0.2
東京	22.2	24.9	26.7	49.7	41.6	28.1	17.7	27.7	23.7	22.4	74.8	28.3	29.1	469	0.0
最大値 ¹⁾	51.8	31.8	33.5	53.5	49.9	30.8	26.0	50.1	48.1	54.0	74.8	47.3	29.1	1070	3.3
最小値 ¹⁾	8.0	2.4	0.6	2.2	1.3	0.7	0.5	1.2	1.3	2.1	3.8	7.0	3.2	56.7	0.0
全国平均	21.0	12.7	10.9	21.2	13.7	11.8	12.9	17.7	20.2	18.0	22.6	20.0	15.0		
標準偏差	10.3	8.0	7.6	12.5	11.3	7.6	7.0	11.4	10.6	10.6	13.2	9.6	6.0		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 N a ⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	140	27.0	39.4	8.9	75.2	117	273	1110	325	418	204	236	241	13800	0.1
札幌	131	18.3	10.1	2.0	57.5	44.8	54.4	250	289	258	261	422	177	760	0.5
落石岬	906	243	59.7	16.6	30.6	13.3	391	468	245	662	*	471	212	10900	3.6
竜飛岬	121	238	55.5	13.9	68.1	456	319	226	238	1370	564	443	228	11500	1.1
八幡平	20.1	19.9	2.7	1.6	2.5	72.8	12.5	60.9	56.1	15.3	53.7	38.4	23.1	639	0.0
篁岳	21.4	26.7	19.2	6.4	6.6	2.9	30.9	12.2	251	17.6	179	27.2	27.9	3530	0.7
尾花沢	15.1	10.2	1.7	3.4	2.0	75.8	21.0	75.4	335	66.5	131	81.2	65.9	394	0.3
筑波	41.6	18.7	5.2	17.7	9.6	4.1	24.7	14.5	17.1	10.4	25.3	34.7	18.4	439	0.0
赤城	7.2	3.2	2.0	2.0	0.9	1.7	6.3	4.1	6.0	4.9	17.6	9.1	3.4	103	0.0
小笠原	102	68.0	28.3	32.1	33.4	24.6	52.8	109	161	75.6	65.7	145	80.1	1980	4.9
佐渡関岬	101	24.8	11.4	14.9	40.8	137	174	531	1640	732	597	121	221	9290	1.8
新潟巻	35.9	14.1	3.2	7.0	38.2	27.1	107	425	401	429	189	111	180	4580	1.5
八方尾根	10.2	1.0	0.6	1.3	1.0	2.2	5.2	9.4	15.8	11.4	7.7	19.6	4.9	343	0.4
越前岬	18.3	10.5	30.2	9.5	66.8	33.3	45.4	446	409	690	128	127	180	2577	0.9
伊自良湖	12.3	6.9	1.8	5.5	7.7	7.2	10.3	41.0	82.4	19.9	37.5	23.5	15.0	125	0.6
犬山	10.3	8.1	2.1	7.4	5.0	6.9	13.9	20.2	57.0	8.4	27.9	30.5	10.9	155	0.9
潮岬	165	78.0	84.5	62.5	97.3	53.1	59.6	55.6	256	73.4	337	462	122	2550	3.5
京都八幡	7.1	5.7	4.3	6.2	5.0	2.8	10.6	15.8	33.4	3.6	23.4	7.9	8.2	88.8	0.7
尼崎	5.0	4.8	3.3	6.3	5.1	5.9	11.0	12.6	89.3	4.9	44.2	25.2	13.5	233	0.7
倉橋島	2.5	5.1	4.2	5.2	9.3	1.1	4.9	15.4	58.0	6.9	21.2	46.1	11.6	363	0.0
隠岐	153	32.4	33.7	117	99.8	26.8	72.3	465	832	1130	557	363	356	10900	3.7
蟠竜湖	29.0	12.1	7.6	12.5	15.5	21.5	20.7	140	174	349	41.6	81.2	82.4	876	1.0
禰原	5.7	4.3	1.8	3.4	4.8	2.4	4.6	10.7	53.6	66.5	17.0	15.6	10.7	553	0.0
筑後小郡	7.7	6.2	6.1	15.0	10.0	6.9	9.8	47.9	41.8	176	23.1	7.4	16.0	1200	1.0
大分久住	4.8	3.9	2.0	5.0	10.6	2.9	1.4	4.5	13.0	14.8	8.7	4.0	4.8	89.3	0.5
対馬	25.1	38.5	23.8	23.1	19.7	17.3	24.3	99.4	53.1	51.5	38.8	61.6	33.0	383	0.9
五島	88.8	31.7	27.5	42.7	48.4	13.2	137.3	158	177	178	99.7	305	83.5	2100	3.1
えびの	7.9	1.9	16.8	11.4	13.9	5.3	4.9	6.0	34.3	44.0	12.0	9.7	12.2	631	0.2
屋久島	56.0	43.8	26.7	39.6	40.1	70.7	34.0	161	186	222	61.3	101	71.1	574	13.9
辺戸岬	70.8	29.7	44.7	58.2	48.3	48.9	74.4	139.5	1080	511	262	174	117	16300	4.9
東京	28.1	21.6	5.3	20.8	17.7	5.2	25.9	11.7	25.3	18.0	45.4	75.4	20.8	510	0.3
最大値 ¹⁾	906	243	84.5	117	100	456	391	1110	1640	1370	597	471	356	16300	13.9
最小値 ¹⁾	2.5	1.0	0.6	1.3	0.9	1.1	1.4	4.1	6.0	3.6	7.7	4.0	3.4	88.8	0.0
全国平均	75.8	34.1	18.2	18.7	28.7	42.3	65.7	166	246	246	136	132	85.6		
標準偏差	162	58.1	20.8	24.2	28.9	84.3	97.2	239	351	350	172	152	93.6		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 K⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	3.5	4.6	3.6	1.2	1.1	2.6	6.0	29.5	7.8	11.7	5.1	7.4	6.8	324	0.0
札幌	4.4	3.3	1.0	0.5	2.5	0.6	2.7	6.9	8.2	7.7	7.0	10.2	5.3	24.2	0.1
落石岬	18.2	5.5	1.5	0.7	0.7	0.4	8.6	10.6	5.6	16.0	*	10.1	4.7	221	0.2
竜飛岬	3.9	9.5	2.1	0.7	1.5	9.9	12.3	8.0	6.0	28.6	11.2	9.0	5.7	315	0.0
八幡平	1.5	1.4	0.2	0.2	0.1	1.6	0.6	2.4	2.3	0.8	1.6	1.3	0.9	33.7	0.0
籠岳	1.4	1.0	0.8	1.0	1.1	0.5	1.0	0.7	6.2	1.2	4.4	1.1	1.3	76.4	0.2
尾花沢	1.1	3.5	0.2	0.9	0.5	2.1	1.3	4.3	7.7	2.1	4.1	2.4	2.2	21.9	0.0
筑波	2.2	1.8	1.2	2.7	1.6	0.8	1.6	1.6	2.0	1.1	1.5	3.9	1.8	42.9	0.2
赤城	1.1	0.8	0.6	1.1	1.5	0.8	0.5	0.8	0.9	0.2	2.3	1.3	0.9	10.8	0.0
小笠原	3.5	4.8	0.8	1.0	0.9	0.7	1.2	2.5	3.7	1.9	2.0	5.3	2.5	977	0.1
佐渡関岬	3.9	9.5	1.1	1.1	1.0	6.6	8.5	11.6	35.4	14.4	13.3	3.5	6.8	590	0.0
新潟巻	2.0	0.8	0.3	0.6	0.8	1.0	2.8	9.0	9.3	9.6	6.2	3.1	4.4	94.9	0.1
八方尾根	1.2	0.3	0.3	0.4	0.3	0.3	0.7	0.6	3.0	1.1	2.0	2.0	0.8	14.7	0.0
越前岬	1.3	0.8	1.4	0.6	3.0	2.5	2.0	9.6	9.8	17.2	4.4	5.7	5.1	65.4	0.0
伊自良湖	0.9	0.3	0.1	0.2	0.3	0.6	0.5	1.1	2.7	1.0	2.1	1.0	0.7	5.4	0.0
犬山	1.1	0.8	0.5	0.8	0.7	0.7	0.7	1.3	2.4	1.0	1.8	1.5	0.9	15.2	0.1
潮岬	3.6	1.9	2.2	1.4	1.5	1.0	1.3	1.5	5.8	1.8	8.0	10.4	2.8	61.6	0.1
京都八幡	2.0	1.3	1.5	1.6	1.4	1.2	1.2	1.8	2.0	1.0	1.4	1.2	1.5	11.2	0.5
尼崎	1.3	0.7	2.0	1.3	1.7	1.0	0.9	1.5	2.9	1.1	2.5	1.7	1.5	14.4	0.0
倉橋島	1.0	0.9	0.6	0.8	0.9	0.5	0.6	1.0	1.5	1.1	1.4	1.9	0.9	14.3	0.2
隠岐	4.4	1.5	1.5	3.6	3.3	1.8	2.7	29.2	20.1	26.3	15.6	11.2	11.2	241	0.4
蟠竜湖	1.4	0.8	0.6	0.9	0.7	0.9	1.0	4.8	5.2	9.9	2.0	3.5	2.8	23.2	0.3
禰原	0.4	0.2	0.1	0.2	0.2	0.1	0.2	0.5	1.5	2.6	0.9	1.1	0.5	15.1	0.0
筑後小郡	1.8	1.8	1.8	3.8	2.6	1.8	1.4	3.8	4.8	9.3	3.0	2.1	2.5	70.6	0.7
大分久住	0.5	1.1	0.1	0.3	0.3	0.3	0.1	0.3	0.6	1.4	1.2	0.7	0.4	6.9	0.0
対馬	1.7	3.0	1.3	2.1	0.3	1.0	5.4	4.3	4.5	2.4	2.8	2.2	2.0	36.0	0.0
五島	10.0	5.0	3.7	1.0	1.1	0.4	4.4	7.0	6.8	8.7	3.7	7.3	4.3	107	0.0
えびの	0.8	0.2	0.3	0.4	0.8	0.7	0.6	0.8	2.8	1.8	0.9	1.0	0.6	37.6	0.0
屋久島	1.2	1.2	1.1	1.1	1.4	1.4	1.0	3.4	4.2	5.4	2.3	2.6	1.9	15.1	0.3
辺戸岬	2.0	1.0	1.3	1.7	1.4	1.8	2.0	3.4	24.6	12.2	6.6	4.6	3.1	351	0.2
東京	1.4	1.0	0.8	2.0	1.2	0.5	0.8	0.7	1.4	0.9	2.5	1.7	1.1	17.6	0.2
最大値 ¹⁾	18.2	9.5	3.7	3.8	3.3	9.9	12.3	29.5	<u>35.4</u>	<u>28.6</u>	15.6	11.2	11.2	977	0.7
最小値 ¹⁾	0.4	0.2	0.1	0.2	0.1	0.1	0.1	0.3	0.6	0.2	0.9	0.7	0.4	5.4	0.0
全国平均	2.7	2.3	1.1	1.2	1.2	1.5	2.4	5.3	6.5	6.5	4.1	3.9	2.8		
標準偏差	3.4	2.4	0.9	0.9	0.8	2.0	2.9	7.2	7.5	7.6	3.7	3.3	2.5		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 Ca²⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	8.9	8.9	3.3	1.0	3.1	4.2	8.9	27.9	11.4	15.6	10.9	68.8	12.1	641	0.0
札幌	32.4	11.8	4.9	2.4	8.1	5.2	6.0	11.8	15.6	12.2	18.8	24.3	12.0	59.5	1.9
落石岬	25.5	8.9	2.1	0.6	1.0	0.5	8.4	9.8	8.4	18.7	*	15.9	6.3	260	0.2
竜飛岬	10.6	22.0	5.9	2.6	9.7	12.3	12.6	9.2	9.2	31.3	14.8	23.2	10.1	380	1.0
八幡平	6.0	8.9	1.3	1.1	0.7	3.2	1.9	6.3	10.6	3.6	9.0	8.0	3.9	270	0.0
籠岳	4.8	2.0	2.1	2.6	1.6	1.0	2.0	1.3	7.6	2.3	16.5	3.8	2.9	79.7	0.2
尾花沢	2.4	1.5	0.4	1.4	0.5	2.9	1.6	6.8	8.5	2.7	12.4	6.1	3.6	18.1	0.2
筑波	5.9	3.4	1.7	8.1	1.2	1.3	2.7	1.5	2.6	4.1	7.9	6.4	3.6	95.9	0.1
赤城	2.6	2.3	0.9	2.4	0.9	1.5	1.8	2.2	1.9	1.6	16.2	6.0	1.9	53.1	0.0
小笠原	5.4	2.5	0.9	1.5	1.0	0.6	1.0	2.0	2.9	1.8	1.7	8.1	2.2	252	0.0
佐渡関岬	15.5	7.4	2.7	3.2	4.3	5.3	9.9	14.5	54.6	16.0	26.9	5.4	9.7	248	0.9
新潟巻	12.0	3.3	1.0	1.6	1.4	1.7	4.4	11.5	12.0	10.8	20.3	21.0	8.5	110	0.3
八方尾根	8.1	2.0	1.7	0.8	0.8	0.5	2.0	2.3	8.8	7.0	7.6	23.1	4.4	332	0.0
越前岬	3.7	1.5	1.4	0.8	2.2	1.3	4.0	10.6	9.3	16.0	9.8	11.8	6.2	342	0.0
伊自良湖	2.8	1.9	0.8	0.6	2.5	1.5	1.0	2.1	5.2	2.0	8.7	3.6	2.4	23.2	0.0
犬山	3.2	2.4	1.6	2.5	3.6	2.7	1.9	4.5	7.4	4.7	12.9	5.4	3.6	91.5	0.1
潮岬	7.3	4.5	3.0	1.8	10.2	5.6	6.3	1.9	7.8	2.4	13.4	15.1	6.0	81.1	0.2
京都八幡	4.4	3.2	2.0	3.0	2.8	1.5	2.2	5.9	4.5	2.2	5.6	4.7	3.1	31.8	0.5
尼崎	4.2	3.1	5.0	3.2	3.4	3.0	3.5	3.8	8.1	3.7	7.7	6.8	4.5	99.9	0.8
倉橋島	2.0	2.9	1.9	1.6	1.5	1.3	1.5	1.6	2.7	1.8	3.3	3.8	2.2	50.3	0.2
隠岐	6.2	3.0	2.4	5.1	3.1	1.6	5.6	12.8	26.8	31.6	37.9	32.2	13.9	287	0.4
蟠竜湖	4.6	1.9	1.6	1.4	1.1	1.3	3.2	6.9	11.0	18.8	7.9	9.3	6.0	57.3	0.6
禱原	1.8	1.3	0.5	0.5	0.7	0.4	0.6	1.2	4.4	5.5	2.8	5.9	1.7	86.8	0.0
筑後小郡	5.2	4.6	2.2	6.9	3.1	3.3	2.4	6.4	5.6	21.0	5.6	3.9	4.2	134	0.7
大分久住	2.5	2.3	0.7	1.1	1.0	0.7	0.9	0.9	2.0	3.3	4.9	5.7	1.7	18.3	0.0
対馬	4.8	5.4	2.2	2.3	1.3	1.7	2.1	5.6	10.8	4.8	8.4	5.4	3.9	148	0.0
五島	3.5	5.2	8.4	3.5	2.6	0.7	4.2	6.7	8.2	9.5	8.4	10.3	5.3	62.2	0.0
えびの	3.7	0.8	0.6	1.0	1.3	0.9	1.0	1.2	3.0	3.1	2.2	4.6	1.4	68.4	0.0
屋久島	2.6	1.5	1.4	1.5	1.4	2.0	1.4	4.7	5.7	7.4	3.8	5.3	2.7	22.0	0.5
辺戸岬	3.9	1.7	1.8	2.1	2.2	1.7	2.4	4.0	33.0	16.1	8.9	5.7	4.0	364	0.5
東京	3.9	3.2	3.0	6.5	4.9	3.0	2.4	3.4	3.9	2.0	12.7	9.0	3.9	153	0.5
最大値 ¹⁾	32.4	22.0	8.4	8.1	10.2	12.3	12.6	27.9	54.6	31.6	37.9	68.8	13.9	641	1.9
最小値 ¹⁾	1.8	0.8	0.4	0.5	0.5	0.4	0.6	0.9	1.9	1.6	1.7	3.6	1.4	18.1	0.0
全国平均	6.8	4.4	2.2	2.4	2.7	2.4	3.6	6.2	10.1	9.1	10.9	11.9	5.1		
標準偏差	6.7	4.3	1.7	1.9	2.5	2.3	3.0	5.6	10.6	8.5	7.7	12.9	3.3		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 nss-C a²⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	5.9	8.3	2.4	0.8	1.5	1.7	3.0	3.9	4.4	6.6	6.4	63.7	6.9	630	0.0
札幌	29.6	11.4	4.7	2.3	6.9	4.2	4.8	6.4	9.4	6.6	13.1	15.2	8.2	49.6	1.8
落石岬	5.9	3.6	0.8	0.3	0.4	0.2	0.4	0.4	3.2	4.4 *		5.7	1.8	92.4	0.0
竜飛岬	8.0	16.9	4.7	2.3	8.2	2.5	5.7	4.3	4.1	3.3	3.0	13.7	5.3	302	0.0
八幡平	5.6	8.4	1.2	1.0	0.6	1.6	1.7	5.0	9.4	3.3	7.8	7.1	3.4	260	0.0
籠岳	4.4	1.4	1.7	2.4	1.5	0.9	1.3	1.0	2.2	1.9	12.7	3.2	2.3	40.5	0.1
尾花沢	2.0	1.3	0.4	1.4	0.4	1.3	1.1	5.2	1.3	1.3	9.6	4.3	2.2	17.8	0.2
筑波	5.0	3.0	1.6	7.7	1.0	1.2	2.2	1.2	2.3	3.9	7.3	5.6	3.2	91.2	0.1
赤城	2.4	2.2	0.9	2.4	0.9	1.5	1.7	2.1	1.8	1.5	15.8	5.8	1.8	52.5	0.0
小笠原	3.2	1.1	0.3	0.9	0.4	0.1	0.1	0.1	0.0	0.2	0.4	5.1	0.7	250	0.0
佐渡関岬	13.3	6.8	2.4	2.9	3.4	2.4	6.1	3.2	19.2	0.9	14.0	2.7	5.0	99.8	0.0
新潟巻	11.3	3.0	0.9	1.4	0.5	1.1	2.1	2.4	3.4	1.8	16.2	18.6	4.6	87.6	0.0
八方尾根	7.9	2.0	1.7	0.8	0.8	0.5	1.9	2.1	8.5	6.8	7.4	22.7	4.3	324	0.0
越前岬	3.3	1.3	0.7	0.6	0.8	0.6	3.1	1.0	0.7	1.9	7.1	9.1	2.4	318	0.0
伊自良湖	2.5	1.7	0.7	0.6	2.4	1.3	0.8	1.2	3.4	1.6	7.9	3.1	2.1	22.3	0.0
犬山	3.0	2.2	1.6	2.3	3.5	2.5	1.6	4.0	6.1	4.5	12.3	4.8	3.4	88.1	0.1
潮岬	3.7	2.8	1.2	0.6	8.1	4.5	5.1	0.7	2.3	0.9	6.1	5.1	3.3	56.5	0.0
京都八幡	4.3	3.1	1.9	2.8	2.7	1.4	1.9	5.5	3.8	2.1	5.1	4.5	3.0	31.2	0.4
尼崎	4.1	3.0	5.0	3.1	3.3	2.9	3.2	3.5	6.2	3.6	6.7	6.2	4.2	98.5	0.7
倉橋島	1.9	2.8	1.8	1.4	1.3	1.3	1.4	1.2	1.5	1.6	2.8	2.8	2.0	49.6	0.1
隠岐	2.9	2.3	1.7	2.6	1.0	1.0	4.0	2.7	8.9	7.2	25.9	24.4	6.2	213	0.2
蟠竜湖	3.9	1.6	1.4	1.2	0.8	0.9	2.8	3.9	7.2	11.3	7.0	7.6	4.2	41.9	0.4
禱原	1.7	1.2	0.5	0.4	0.6	0.4	0.5	1.0	3.3	4.0	2.4	5.5	1.5	76.5	0.0
筑後小郡	5.1	4.5	2.1	6.5	2.9	3.2	2.2	5.4	4.7	17.2	5.1	3.8	3.8	108	0.6
大分久住	2.4	2.2	0.6	1.0	0.8	0.7	0.9	0.8	1.7	3.0	4.7	5.6	1.6	17.4	0.0
対馬	4.2	4.5	1.7	1.8	0.9	1.4	1.6	3.5	9.7	3.6	7.5	4.1	3.1	147	0.0
五島	1.5	4.6	7.8	2.5	1.5	0.5	1.3	3.3	4.3	5.6	6.2	3.7	3.5	45.7	0.0
えびの	3.6	0.8	0.4	0.7	1.0	0.8	0.9	1.1	2.3	2.2	1.9	4.4	1.2	66.3	0.0
屋久島	1.4	0.5	0.9	0.6	0.6	0.5	0.7	1.3	1.7	2.6	2.5	3.1	1.2	17.7	0.2
辺戸岬	2.4	1.0	0.8	0.8	1.2	0.6	0.8	1.0	9.8	5.0	3.3	2.2	1.5	120	0.0
東京	3.3	2.8	2.9	6.1	4.5	2.9	1.8	3.1	3.3	1.6	11.7	7.4	3.5	149	0.4
最大値 ¹⁾	29.6	16.9	7.8	7.7	8.2	4.5	6.1	6.4	19.2	17.2	25.9	63.7	8.2	630	1.8
最小値 ¹⁾	1.4	0.5	0.3	0.3	0.4	0.1	0.1	0.1	0.0	0.2	0.4	2.2	0.7	17.4	0.0
全国平均	5.1	3.6	1.8	2.0	2.1	1.5	2.2	2.6	4.8	3.9	8.0	9.1	3.3		
標準偏差	5.3	3.5	1.6	1.8	2.2	1.1	1.6	1.8	3.9	3.4	5.4	11.6	1.8		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 Mg²⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	21.4	5.1	5.7	1.3	9.8	15.2	33.9	129	38.3	49.7	25.3	31.0	29.4	1550	0.0
札幌	18.0	3.5	2.1	0.8	7.8	5.9	7.9	30.8	34.7	31.6	33.0	50.0	22.0	89.3	0.5
落石岬	105	28.4	7.5	1.8	3.7	1.8	42.5	51.9	28.5	75.5	*	52.1	24.2	1150	0.3
竜飛岬	14.4	28.8	9.2	2.1	9.2	50.0	32.9	26.6	25.6	147	51.2	45.2	24.5	1120	0.6
八幡平	3.4	3.8	0.5	0.4	0.4	8.3	1.7	7.3	7.6	2.0	6.5	5.4	3.0	87.3	0.0
篔岳	3.1	3.1	2.3	0.9	0.9	0.7	3.6	1.4	28.8	2.1	20.2	3.2	3.3	394	0.0
尾花沢	1.8	0.6	0.2	0.5	0.2	8.8	2.4	9.3	38.3	7.5	15.9	9.7	7.7	45.2	0.0
筑波	6.2	3.0	0.9	4.1	0.9	0.6	3.2	2.3	4.7	1.7	3.6	4.6	2.8	52.8	0.0
赤城	1.3	0.7	0.5	1.0	0.4	1.2	2.2	2.1	2.1	2.1	5.9	3.7	1.1	16.1	0.0
小笠原	12.9	8.8	3.7	4.4	5.0	3.3	7.2	13.5	20.3	10.3	8.6	18.2	10.4	234	0.6
佐渡関岬	13.7	3.8	2.1	2.4	5.1	15.7	20.6	51.7	164	76.7	51.5	13.6	23.1	769	0.7
新潟巻	6.4	2.2	0.6	1.2	4.3	3.2	12.7	49.7	45.4	47.6	23.4	13.8	21.0	507	0.2
八方尾根	2.6	0.2	0.1	0.1	0.1	0.2	0.7	1.6	2.1	1.7	1.7	4.4	0.9	48.7	0.0
越前岬	2.7	1.4	3.3	0.9	7.2	3.2	5.3	49.4	45.5	76.4	15.3	15.2	20.2	287	0.0
伊自良湖	1.7	0.9	0.1	0.5	0.7	0.4	1.1	4.6	9.4	2.3	5.0	3.0	1.7	14.4	0.0
犬山	1.7	1.2	0.5	1.2	0.8	1.0	1.6	2.7	6.8	1.3	4.7	4.1	1.6	35.6	0.0
潮岬	18.4	9.5	9.9	7.5	2.2	1.1	1.4	6.2	30.2	7.8	38.3	52.2	12.3	295	0.1
京都八幡	1.8	1.6	1.4	1.5	1.4	1.1	1.9	2.5	4.2	1.2	3.2	1.8	1.8	8.5	0.4
尼崎	1.3	1.1	2.5	1.2	1.2	0.8	2.0	1.8	11.1	1.2	5.9	4.1	2.4	29.4	0.0
倉橋島	1.0	1.2	1.1	1.1	1.5	1.3	1.5	1.5	5.1	0.7	2.3	4.6	1.7	40.1	0.2
隠岐	18.2	4.3	4.1	14.3	11.4	3.4	9.4	47.2	95.6	128	64.5	44.9	40.6	1220	0.8
蟠竜湖	4.1	1.8	1.2	1.7	1.8	2.6	2.9	16.2	20.3	40.3	5.2	10.0	9.8	99.6	0.4
禰原	1.0	0.5	0.2	0.4	0.6	0.3	0.5	1.2	6.5	8.3	2.3	2.1	1.4	55.5	0.0
筑後小郡	1.8	1.6	0.8	2.6	1.3	1.0	0.9	6.7	4.1	28.9	2.2	0.8	2.3	281	0.4
大分久住	0.8	0.8	0.2	0.7	1.1	0.5	0.2	0.5	2.3	2.5	2.2	1.6	0.8	11.9	0.0
対馬	4.5	5.1	2.9	3.2	2.3	2.5	2.8	9.2	8.1	5.0	5.7	6.7	4.1	33.8	0.0
五島	7.0	3.3	3.2	4.9	5.5	1.5	11.0	15.9	17.3	16.2	10.8	29.3	8.1	151	0.8
えびの	1.8	0.6	0.4	1.1	1.6	0.6	1.1	0.8	3.8	5.0	1.4	1.4	1.1	75.6	0.0
屋久島	6.9	5.7	3.4	4.9	4.7	8.7	4.2	19.7	23.1	28.6	7.6	12.4	8.9	75.2	1.6
辺戸岬	8.5	3.5	5.2	6.7	5.6	6.0	8.8	16.0	119	54.6	29.9	19.7	13.3	1580	0.6
東京	3.6	3.1	2.0	2.9	2.6	0.9	3.0	1.7	3.1	2.2	6.0	9.1	2.9	55.7	0.0
最大値 ¹⁾	105	28.8	9.9	14.3	11.4	50.0	42.5	129	<u>164</u>	<u>147</u>	64.5	52.2	40.6	1580	1.6
最小値 ¹⁾	0.8	0.2	0.1	0.1	0.1	0.2	0.2	0.5	<u>2.1</u>	0.7	1.4	<u>0.8</u>	0.8	8.5	0.0
全国平均	9.6	4.5	2.5	2.5	3.3	4.9	7.5	18.7	27.6	27.9	15.3	15.4	9.9		
標準偏差	18.7	6.8	2.6	2.9	3.1	9.3	10.7	26.9	36.6	38.3	17.1	16.8	10.5		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 H⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	4.5	21.3	19.4	5.9	5.1	6.8	8.8	15.8	8.6	26.0	10.3	11.5	11.6	97.7	0.1
札幌	19.9	13.6	23.5	24.5	8.3	9.7	18.2	29.7	27.9	37.5	21.5	15.4	23.9	174	0.5
落石岬	14.7	13.7	15.1	23.1	13.3	6.4	17.2	23.4	10.9	12.0	*	5.6	12.9	59.7	0.6
竜飛岬	41.3	32.9	25.3	20.0	13.3	16.7	43.5	23.8	34.2	8.7	13.2	10.1	21.2	245	0.3
八幡平	40.8	20.7	5.8	16.3	9.3	17.4	20.5	34.1	24.4	24.7	13.0	14.3	17.0	115	1.0
篔岳	22.2	15.9	13.0	39.8	13.1	10.5	19.0	19.4	31.1	14.6	6.5	13.8	17.5	282	1.0
尾花沢	12.7	12.9	7.1	34.0	11.3	20.8	16.8	49.0	22.7	16.0	17.7	18.0	18.5	53.7	3.8
筑波	10.7	8.1	29.3	48.2	11.6	13.8	16.0	19.2	14.8	11.5	23.0	7.3	14.0	216	0.3
赤城	15.7	15.8	9.9	16.9	14.6	18.4	21.9	24.1	11.3	10.6	17.4	7.8	15.2	106	2.4
小笠原	21.1	14.0	7.4	8.9	7.7	7.0	5.2	6.8	6.2	7.6	12.6	16.1	8.8	61.2	1.7
佐渡関岬	33.7	15.5	15.9	29.5	18.5	37.8	36.4	23.0	38.9	18.9	19.8	20.6	25.5	107	0.1
新潟巻	28.6	22.9	13.0	38.6	22.7	37.2	31.8	20.7	31.8	30.7	37.0	12.9	26.8	148	0.1
八方尾根	14.5	7.3	6.6	22.8	7.0	18.8	24.2	7.9	17.6	21.9	17.9	12.6	13.3	70.8	0.1
越前岬	23.8	10.9	21.3	20.0	24.6	23.7	32.5	29.4	29.4	29.5	24.5	18.2	23.8	166	0.1
伊自良湖	29.7	26.3	29.6	37.6	49.4	60.5	21.0	32.9	56.0	32.7	33.2	16.3	33.0	107	4.7
犬山	25.3	21.4	34.5	37.6	28.5	27.4	20.9	36.5	40.2	23.7	12.7	14.9	26.4	229	1.9
潮岬	21.9	12.4	13.4	23.5	14.1	14.6	15.7	23.1	20.6	23.2	20.4	18.4	17.4	145	0.9
京都八幡	27.7	13.8	19.9	30.7	38.5	24.7	29.5	27.1	29.3	20.6	16.2	9.9	22.7	126	3.8
尼崎	22.2	12.6	24.5	42.7	31.8	24.0	23.7	24.9	41.7	14.2	15.2	17.7	23.2	135	1.3
倉橋島	25.8	25.3	30.1	56.9	27.9	20.0	23.4	41.2	32.1	33.0	35.3	28.1	28.7	275	6.0
隠岐	11.1	23.6	12.9	15.1	14.4	22.9	29.2	37.7	36.6	25.7	18.2	16.6	23.2	394	0.6
蟠竜湖	19.1	11.1	16.4	41.1	23.1	31.7	51.0	54.2	34.6	51.2	25.2	34.6	30.1	129	5.5
禰原	21.0	16.7	17.3	19.7	25.3	12.5	15.2	30.4	43.3	52.9	22.9	12.5	21.1	240	0.9
筑後小郡	15.1	12.4	10.7	15.0	17.1	17.5	38.4	62.5	38.6	38.2	12.4	5.5	17.4	204	0.6
大分久住	20.0	13.0	12.0	28.9	34.8	26.2	14.7	37.7	34.0	24.9	23.6	11.2	20.4	120	0.5
対馬	26.3	22.5	28.3	39.4	23.1	37.8	39.5	47.3	66.9	31.0	38.6	34.1	32.2	182	1.4
五島	21.3	11.5	4.0	21.7	23.3	17.9	18.5	42.9	36.8	40.0	32.3	31.1	21.2	251	0.1
えびの	19.1	12.0	7.9	14.4	11.5	19.8	12.1	24.7	33.8	27.1	23.2	12.0	14.9	245	0.4
屋久島	24.3	11.4	11.9	8.4	28.4	10.3	34.5	60.7	34.1	42.8	36.4	31.4	22.4	112	5.2
辺戸岬	7.0	9.2	6.5	7.7	12.4	6.2	8.2	9.0	20.8	10.6	10.9	10.8	8.6	87.5	1.1
東京	9.2	10.3	27.8	47.4	42.0	24.8	24.6	26.0	19.5	13.3	39.0	15.9	24.3	483	1.2
最大値 ¹⁾	41.3	32.9	34.5	56.9	49.4	60.5	51.0	62.5	66.9	52.9	39.0	34.6	33.0	483	6.0
最小値 ¹⁾	4.5	7.3	4.0	5.9	5.1	6.2	5.2	6.8	6.2	7.6	6.5	5.5	8.6	53.7	0.1
全国平均	21.0	15.8	16.8	27.0	20.2	20.8	23.6	30.5	30.0	25.0	21.7	16.3	20.6		
標準偏差	8.7	6.1	8.6	13.3	11.0	11.6	10.8	14.1	13.3	12.1	9.3	7.9	6.4		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 pH 月・年平均値

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	5.34	4.67	4.71	5.23	5.29	5.17	5.05	4.80	5.06	4.59	4.99	4.94	4.94	6.92	4.01
札幌	4.70	4.87	4.63	4.61	5.08	5.01	4.74	4.53	4.55	4.43	4.67	4.81	4.62	6.29	3.76
落石岬	4.83	4.86	4.82	4.64	4.87	5.19	4.76	4.63	4.96	4.92	*	5.25	4.89	6.21	4.22
竜飛岬	4.38	4.48	4.60	4.70	4.88	4.78	4.36	4.62	4.47	5.06	4.88	5.00	4.67	6.55	3.61
八幡平	4.39	4.68	5.24	4.79	5.03	4.76	4.69	4.47	4.61	4.61	4.89	4.84	4.77	6.01	3.94
篔岳	4.65	4.80	4.89	4.40	4.88	4.98	4.72	4.71	4.51	4.84	5.19	4.86	4.76	6.01	3.55
尾花沢	4.90	4.89	5.15	4.47	4.95	4.68	4.78	4.31	4.64	4.80	4.75	4.75	4.73	5.42	4.27
筑波	4.97	5.09	4.53	4.32	4.94	4.86	4.80	4.72	4.83	4.94	4.64	5.14	4.85	6.49	3.67
赤城	4.80	4.80	5.00	4.77	4.84	4.74	4.66	4.62	4.95	4.98	4.76	5.11	4.82	5.61	3.98
小笠原	4.68	4.85	5.13	5.05	5.12	5.16	5.29	5.17	5.21	5.12	4.90	4.79	5.06	5.78	4.21
佐渡関岬	4.47	4.81	4.80	4.53	4.73	4.42	4.44	4.64	4.41	4.72	4.70	4.69	4.59	7.08	3.97
新潟巻	4.54	4.64	4.89	4.41	4.64	4.43	4.50	4.68	4.50	4.51	4.43	4.89	4.57	6.91	3.83
八方尾根	4.84	5.14	5.18	4.64	5.16	4.72	4.62	5.10	4.75	4.66	4.75	4.90	4.88	6.87	4.15
越前岬	4.62	4.96	4.67	4.70	4.61	4.63	4.49	4.53	4.53	4.53	4.61	4.74	4.62	7.30	3.78
伊自良湖	4.53	4.58	4.53	4.43	4.31	4.22	4.68	4.48	4.25	4.49	4.48	4.79	4.48	5.33	3.97
犬山	4.60	4.67	4.46	4.43	4.54	4.56	4.68	4.44	4.40	4.62	4.90	4.83	4.58	5.73	3.64
潮岬	4.66	4.91	4.87	4.63	4.85	4.84	4.80	4.64	4.69	4.64	4.69	4.74	4.76	6.05	3.84
京都八幡	4.56	4.86	4.70	4.51	4.41	4.61	4.53	4.57	4.53	4.69	4.79	5.01	4.64	5.42	3.90
尼崎	4.65	4.90	4.61	4.37	4.50	4.62	4.63	4.60	4.38	4.85	4.82	4.75	4.63	5.88	3.87
倉橋島	4.59	4.60	4.52	4.25	4.55	4.70	4.63	4.39	4.49	4.48	4.45	4.55	4.54	5.22	3.56
隠岐	4.95	4.63	4.89	4.82	4.84	4.64	4.53	4.42	4.44	4.59	4.74	4.78	4.63	6.21	3.41
蟠竜湖	4.72	4.95	4.79	4.39	4.64	4.50	4.29	4.27	4.46	4.29	4.60	4.46	4.52	5.26	3.89
禰原	4.68	4.78	4.76	4.71	4.60	4.90	4.82	4.52	4.36	4.28	4.64	4.90	4.68	6.06	3.62
筑後小郡	4.82	4.91	4.97	4.82	4.77	4.76	4.42	4.20	4.41	4.42	4.91	5.26	4.76	6.22	3.69
大分久住	4.70	4.89	4.92	4.54	4.46	4.58	4.83	4.42	4.47	4.60	4.63	4.95	4.69	6.29	3.92
対馬	4.58	4.65	4.55	4.40	4.64	4.42	4.40	4.32	4.17	4.51	4.41	4.47	4.49	5.86	3.74
五島	4.67	4.94	5.40	4.66	4.63	4.75	4.73	4.37	4.43	4.40	4.49	4.51	4.67	6.88	3.60
えびの	4.72	4.92	5.10	4.84	4.94	4.70	4.92	4.61	4.47	4.57	4.63	4.92	4.83	6.41	3.61
屋久島	4.61	4.94	4.93	5.08	4.55	4.99	4.46	4.22	4.47	4.37	4.44	4.50	4.65	5.28	3.95
辺戸岬	5.16	5.04	5.19	5.11	4.91	5.21	5.09	5.04	4.68	4.98	4.96	4.97	5.07	5.94	4.06
東京	5.04	4.99	4.56	4.32	4.38	4.60	4.61	4.59	4.71	4.88	4.41	4.80	4.62	5.93	3.32
最大値 ¹⁾	5.34	5.14	5.40	5.23	5.29	5.21	5.29	5.17	5.21	5.12	5.19	5.26	5.07	7.30	4.27
最小値 ¹⁾	4.38	4.48	4.46	4.25	4.31	4.22	4.29	4.20	4.17	4.28	4.41	4.46	4.48	5.22	3.32
全国平均	4.72	4.83	4.84	4.63	4.76	4.75	4.68	4.57	4.57	4.66	4.70	4.83	4.71		
標準偏差	0.21	0.16	0.25	0.25	0.25	0.24	0.22	0.23	0.23	0.23	0.20	0.21	0.15		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

平成20年度 電気伝導率 月・年平均値

単位：mS m⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	3.26	2.20	2.35	0.96	1.84	2.41	4.92	17.8	6.17	8.83	4.29	6.01	4.86	213	0.17
札幌	4.47	1.64	2.12	1.69	2.50	1.75	2.39	5.94	6.52	6.66	5.92	7.97	4.61	15.2	0.80
落石岬	15.1	4.68	1.77	1.53	1.18	0.53	6.72	7.22	4.43	7.92	*	7.71	3.85	34.8	0.06
竜飛岬	4.05	5.14	2.52	1.49	2.19	8.46	6.76	4.63	5.43	16.8	8.58	7.45	4.46	91.6	0.45
八幡平	3.13	1.85	0.47	1.14	0.62	2.22	1.44	3.11	2.69	1.64	1.89	1.72	1.50	21.7	0.16
籠岳	1.92	1.36	1.11	2.55	0.88	0.74	1.60	1.23	6.04	1.23	3.83	1.37	1.55	58.2	0.19
尾花沢	1.45	0.99	0.46	2.16	0.64	2.26	1.44	4.65	6.64	2.07	3.72	2.45	2.25	7.48	0.31
筑波	1.65	1.00	1.76	3.29	0.96	1.01	1.53	1.48	1.33	0.98	2.08	1.49	1.33	13.0	0.20
赤城	1.23	1.11	0.63	1.48	0.83	1.17	1.42	1.38	0.86	0.62	1.71	0.79	1.03	6.68	0.14
小笠原	2.96	1.76	0.85	0.98	0.94	0.78	1.10	1.99	2.75	1.63	1.78	3.39	1.73	20.0	0.37
佐渡関岬	3.84	1.41	1.13	1.99	1.75	4.24	5.13	8.52	4.57	10.4	6.12	3.03	4.39	24.1	0.60
新潟巻	2.80	1.60	0.80	2.37	1.82	2.30	3.51	7.85	8.04	8.08	5.22	3.06	4.38	44.2	0.43
八方尾根	1.48	0.55	0.54	1.35	0.56	1.12	1.43	0.62	1.60	1.63	1.50	1.69	1.03	13.9	0.25
越前岬	1.73	0.87	1.58	1.25	2.38	1.60	2.43	7.81	7.81	12.3	3.51	3.30	4.05	44.3	0.32
伊自良湖	1.89	1.65	1.52	2.20	2.71	3.39	1.27	2.34	4.46	2.12	2.83	1.49	2.08	7.11	0.29
犬山	1.54	1.25	1.62	2.71	1.78	1.73	1.44	2.33	3.21	1.66	1.78	1.55	1.72	13.8	0.22
潮岬	3.93	1.96	2.15	2.29	2.35	1.52	1.78	2.04	4.98	2.35	6.26	8.05	2.85	41.0	0.32
京都八幡	1.66	0.93	1.13	1.94	2.42	1.38	1.70	1.98	2.32	1.31	1.55	0.89	1.45	8.28	0.21
尼崎	1.38	0.93	1.52	2.81	2.21	1.51	1.49	1.75	3.51	1.02	1.83	1.67	1.67	7.91	0.29
倉橋島	1.51	1.57	1.75	3.06	1.75	1.26	1.44	2.47	2.83	2.09	2.29	2.27	1.82	15.5	0.45
隠岐	2.93	2.08	1.43	3.03	2.50	1.73	2.86	8.75	14.4	18.0	10.4	7.69	6.79	161	0.55
蟠竜湖	1.91	1.06	1.24	2.83	1.70	2.13	3.09	5.25	4.95	8.49	2.43	3.66	3.21	18.6	0.48
禰原	1.11	0.86	0.83	0.93	1.05	0.58	0.67	1.53	3.20	3.84	1.56	1.24	1.20	13.6	0.23
筑後小郡	1.47	1.16	0.86	1.81	1.42	1.30	2.10	4.52	3.05	6.23	1.72	0.84	1.55	34.0	0.21
大分久住	1.33	0.83	0.63	1.64	2.02	1.52	0.77	2.03	2.01	1.75	1.68	0.92	1.21	6.95	0.32
対馬	2.36	2.12	2.01	2.62	1.64	2.25	2.63	4.28	5.20	2.67	3.03	2.98	2.45	11.9	0.36
五島	3.02	1.56	1.21	2.03	2.27	1.08	3.22	4.92	4.72	4.40	3.51	6.63	2.65	27.7	0.37
えびの	1.38	0.71	0.74	1.13	0.97	0.84	0.85	1.29	2.37	2.29	1.57	1.03	1.02	12.3	0.11
屋久島	2.31	1.38	1.11	1.15	2.22	1.83	2.57	6.08	5.59	6.67	3.18	3.66	2.54	17.8	0.63
辺戸岬	1.74	0.98	1.23	1.44	1.53	1.24	1.63	2.81	18.2	7.25	4.99	3.63	2.35	227	0.35
東京	1.33	1.31	1.78	3.26	2.83	1.71	1.91	1.91	1.73	1.18	4.08	2.26	1.93	28.8	0.35
最大値 ¹⁾	15.1	5.14	2.52	3.29	2.83	8.46	6.76	17.8	18.2	18.0	10.4	8.05	6.79	227	0.80
最小値 ¹⁾	1.11	0.55	0.46	0.93	0.56	0.53	0.67	0.62	0.86	0.62	1.50	0.79	1.02	6.68	0.06
全国平均	2.64	1.56	1.32	1.97	1.69	1.86	2.36	4.21	4.89	4.97	3.49	3.29	2.57		
標準偏差	2.50	0.99	0.57	0.73	0.68	1.46	1.58	3.49	3.61	4.64	2.21	2.43	1.44		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。