

**OResults of Radioactive Material Monitoring of Aquatic Organisms (Location K off the mouth of the Abukuma River)**

<Location K off the mouth of the Abukuma River: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
K-3	○	○	○	○	○	○

<Location K off the mouth of the Abukuma River: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment			Other		
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
K-3(Surface layer)	38.0458°	140.9518°	2017/6/27	08:25	08:34	19.8	12.2	Silt with sand	5Y4/2	Juvenile shellfish a little	21.0	2.3
K-3(Bottom layer)				07:53		12.7						

<Location K off the mouth of the Abukuma River: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
K-3(Surface layer)	38.0458°	140.9518°	2017/6/27	08:25	8.2	1.1	3.4	8.9	4600	30.13	1.6	2	2.3	N.D.(0.00095)	0.0060	-
K-3(Bottom layer)				07:53	8.0	<0.5	2.2	8.5	4970	33.38	1.2	4	2.5	0.0013	0.0091	0.0011

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location K off the mouth of the Abukuma River: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>NHE</sub> (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
K-3	38.0458°	140.9518°	2017/6/27	08:34	7.4	-23	37.5	4.6	7.4	2.689	0.0	0.1	0.4	29.3	55.7	14.5	0.050	2.0	28	190	N.D.(0.13)

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location K off the mouth of the Abukuma River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
Surrounding water area off the mouth of the Abukuma River	Sea area in front of the Abukuma River Estuary	-	-	2017/6/17	Vertebrata	Osteichthyes	Scorpaeniformes	Hexagrammidae	<i>Hexagrammos otakii</i>	Fat greenling	2	0.29	Immature fish,Mature fish	Shrimp	Viscera removed	1.63	0.43	1.2	-
					Vertebrata	Osteichthyes	Scorpaeniformes	Scorpaenidae	<i>Sebastes cheni</i>	Rockfish	1	0.064	Immature fish	Shrimp	Viscera removed	1.1	N.D.(1.2)	1.1	-
					Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Kareius bicoloratus</i>	Stone flounder	1	0.53	Immature fish	Shellfish	Viscera removed	0.47	N.D.(0.37)	0.47	-
					Vertebrata	Osteichthyes	Pleuronectiformes	Paralichthyidae	<i>Paralichthys olivaceus</i>	Bastard halibut	2	0.69	Immature fish	Anchovy	Viscera removed	0.39	N.D.(0.27)	0.39	-
					Vertebrata	Osteichthyes	Perciformes	Lateolabracidae	<i>Lateolabrax japonicus</i>	Japanese sea bass	2	1.9	Immature fish,Mature fish	Fish	Viscera removed	0.89	N.D.(0.33)	0.89	N.D.(0.015)
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Acanthogobius flavimanus</i>	Yellowfin Goby	27	0.42	Immature fish,Mature fish	-	-	2.36	0.26	2.1	-

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.