

Table 33 General appearance in F1 and F2 male rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal day 0				
		Control	HBCD (ppm)	150	1500	15000
F1	Number of litters examined	24		21	20	18
	Number of pups examined	164		132	113	139
	Number of pups with abnormal findings	0 (0.00)		3 (2.38)	2 (2.92)	1 (1.11)
	Findings ^a					
	Rudimentary tail	0 (0.00)		1 (0.60)	0 (0.00)	0 (0.00)
	Found dead	0 (0.00)		2 (1.79)	2 (2.92)	1 (1.11)
F2	Number of litters examined	23		22	20	21
	Number of pups examined	159		157	139 ^b	134
	Number of pups with abnormal findings	2 (1.48)		1 (0.57)	6 (4.11)	4 (2.75)
	Findings ^a					
	Found dead	2 (1.48)		1 (0.57)	6 (4.11)	4 (2.75)

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of pups that showed abnormal findings.

b: Including one animal that was not distinguished its sex because of maternal cannibalism.

Table 33 (continued-1) General appearance in F1 and F2 male rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 1-4			
		HBCD (ppm)			
		Control	150	1500	15000
F1	Number of litters examined	24	21	20	18
	Number of pups examined	164	130	111	138
	Number of pups with abnormal findings	11 (5.34)	3 (2.15)	1 (0.56)	6 (3.33)
	Findings ^a				
	Rudimentary tail	0 (0.00)	1 (0.68)	0 (0.00)	0 (0.00)
	Found dead/lost	11 (5.34)	2 (1.47)	1 (0.56)	6 (3.33)
F2	Number of litters examined	23	22	20	21
	Number of pups examined	157	156	133	130
	Number of pups with abnormal findings	19 (11.83)	13 (8.54)	14 (11.42)	36 (29.16)
	Findings ^a				
	Found dead/lost	19 (11.83)	13 (8.54)	14 (11.42)	36 (29.16)

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated.

a: Values represent the number of pups that showed abnormal findings.

Table 33 (continued-2) General appearance in F1 and F2 male rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 5-21			
		HBCD (ppm)			
		Control	150	1500	15000
F1	Number of litters examined	24	21	20	18
	Number of pups examined	99	81	76	80
	Number of pups with abnormal findings	9 (9.38)	0 (0.00)*	2 (2.50)	5 (5.56)
	Findings ^a				
	Moribund condition	4 (4.17)	0 (0.00)	2 (2.50)	0 (0.00)
	Found dead/lost	9 (9.38)	0 (0.00)*	2 (2.50)	5 (5.56)
F2	Number of litters examined	22	22	20	19
	Number of pups examined	82	88	77	70
	Number of pups with abnormal findings	11 (12.27)	9 (10.00)	25 (35.50)*	33 (49.91)**
	Findings ^a				
	Enlargement of eyeball	0 (0.00)	0 (0.00)	0 (0.00)	1 (1.32)
	Found dead/lost	11 (12.27)	9 (10.00)	25 (35.50)*	32 (48.60)**

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated.

a: Values represent the number of pups that showed abnormal findings.

*: Significantly different from the control at $p \leq 0.05$ by Wilcoxon rank-sum test.

**: Significantly different from the control at $p \leq 0.01$ by Wilcoxon rank-sum test.

Table 33 (continued-3) General appearance in F1 and F2 male rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 22-26			
		Control	HBCD (ppm)		
F1	Number of litters examined	23	21	20	17
	Number of weanlings examined	90	80	70	72
	Number of weanlings with abnormal findings	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
F2	Number of litters examined	22	22	18	13
	Number of weanlings examined	71	79	52	38
	Number of weanlings with abnormal findings	0 (0.00)	0 (0.00)	1 (1.39)	2 (4.49)
	Findings ^a				
	Enlargement of eyeball	0 (0.00)	0 (0.00)	1 (1.39)	1 (2.56)
	Opacity of eyeball	0 (0.00)	0 (0.00)	0 (0.00)	1 (1.92)

Values in parentheses represent the means of incidences of weanlings with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of weanlings that showed abnormal findings.

Table 34 General appearance in F1 and F2 female rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal day 0			
		Control	150	1500	15000
F1	Number of litters examined	23	21	20	18
	Number of pups examined	149	148	152	104
	Number of pups with abnormal findings	1 (0.87)	5 (3.07)	1 (0.71)	2 (1.73)
	Findings ^a				
	Kinked tail	0 (0.00)	0 (0.00)	0 (0.00)	1 (0.62)
	Found dead	1 (0.87)	5 (3.07)	1 (0.71)	1 (1.11)
F2	Number of litters examined	23	23	20	21
	Number of pups examined	145	162	129	142
	Number of pups with abnormal findings	1 (1.09)	6 (3.97)	6 (3.33)	3 (1.87)
	Findings ^a				
	Black discoloration of tail	0 (0.00)	0 (0.00)	0 (0.00)	1 (0.60)
	Found dead	1 (1.09)	6 (3.97)	6 (3.33)	2 (1.28)

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of pups that showed abnormal findings.

Table 34 (continued-1) General appearance in F1 and F2 female rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 1-4			
		HBCD (ppm)			
		Control	150	1500	15000
F1	Number of litters examined	23	21	20	18
	Number of pups examined	148	143	151	103
	Number of pups with abnormal findings	5 (3.80)	2 (1.16)	3 (1.93)	5 (6.17)
	Findings ^a				
	Kinked tail	0 (0.00)	0 (0.00)	0 (0.00)	1 (0.62)
	Found dead/lost	5 (3.80)	2 (1.16)	3 (1.93)	4 (5.56)
F2	Number of litters examined	23	23	20	21
	Number of pups examined	144	156	123	140
	Number of pups with abnormal findings	26 (15.20)	16 (13.45)	9 (6.87)	47 (32.12)
	Findings ^a				
	Black discoloration of tail	0 (0.00)	0 (0.00)	0 (0.00)	1 (0.60)
	Found dead/lost	26 (15.20)	16 (13.45)	9 (6.87)	47 (32.12)

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated.

a: Values represent the number of pups that showed abnormal findings.

7
107

Table 34 (continued-2) General appearance in F1 and F2 female rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 5-21			
		Control	HBCD (ppm)	150	1500
F1	Number of litters examined	23	21	20	18
	Number of pups examined	92	87	81	62
	Number of pups with abnormal findings	4 (4.35)	1 (0.79)	1 (1.25)	4 (6.94)
	Findings ^a				
	Moribund condition	1 (1.09)	1 (0.79)	0 (0.00)	0 (0.00)
	Found dead/lost	4 (4.35)	1 (0.79)	1 (1.25)	4 (6.94)
F2	Number of litters examined	22	22	20	20
	Number of pups examined	84	87	83	66
	Number of pups with abnormal findings	15 (17.05)	9 (11.14)	21 (25.35)	30 (48.33)*
	Findings ^a				
	Found dead/lost	15 (17.05)	9 (11.14)	21 (25.35)	30 (48.33)*

(to be continued)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of pups that showed abnormal findings.

*: Significantly different from the control at $p \leq 0.05$ by Wilcoxon rank-sum test.

Table 34 (continued-3) General appearance in F1 and F2 female rat pups and weanlings during postnatal days 0-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Postnatal days 22-26			
		Control	HBCD (ppm)	150	1500
F1	Number of litters examined	23	21	20	17
	Number of weanlings examined	88	85	73	55
	Number of weanlings with abnormal findings	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
F2	Number of litters examined	21	22	20	13
	Number of weanlings examined	69	78	62	36
	Number of weanlings with abnormal findings	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)

Values in parentheses represent the means of incidences of weanlings with abnormal findings (%), the litter is the unit evaluated).

79
60

Table 35 Body weights of F1 and F2 rat pups treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of litters	Male					Female				
			Body weight (g) on postnatal day					Body weight (g) on postnatal day				
			0	4	7	14	21	0	4	7	14	21
F1	Control	24	Mean	6.8	10.2	16.4	(23)	(23)	(23)	(23)	(23)	(23)
			S.D.	0.5	1.7	3.1	4.8	7.1	0.5	1.4	2.8	5.3
	HBCD 150 ppm	21	Mean	6.9	10.7	17.5	36.3	62.3	6.6	10.3	17.0	35.5
			S.D.	0.6	1.8	2.4	3.6	6.5	0.7	1.8	2.5	6.4
	HBCD 1500 ppm	20	Mean	7.2	10.8	16.9	36.1	61.9	6.8 *	10.4	16.9	35.7
			S.D.	0.7	1.6	2.2	3.9	6.5	0.6	1.5	2.3	5.9
	HBCD 15000 ppm	18	Mean	6.8	9.5	15.6	33.5	55.4 *	6.5	9.2	15.1	32.6
			S.D.	0.6	1.8	2.0	2.6	4.0	0.7	1.6	1.6	4.7
F2	Control	23	Mean	6.8	(22)	(22)	(22)	(22)	(22)	(21)	(21)	(21)
			S.D.	0.8	2.3	3.9	8.0	12.6	0.8	2.3	3.5	6.5
	HBCD 150 ppm	23	Mean	6.7	(22)	(22)	(22)	(22)	(22)	(22)	(22)	(22)
			S.D.	0.7	1.3	2.8	5.0	6.7	0.6	1.3	2.8	5.1
	HBCD 1500 ppm	20	Mean	7.1	9.0	14.3	31.0	54.1	6.7	8.8	13.5	29.3
			S.D.	0.6	1.8	3.6	7.2	10.1	0.6	1.8	3.9	10.8
	HBCD 15000 ppm	21	Mean	6.6	(19)	(17)	(14)	(13)	(20)	(17)	(13)	(13)
			S.D.	0.6	1.3	2.9	6.6	8.3	0.6	1.3	2.6	8.4

Values in parentheses are the number of litters examined.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

ss: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

Table 36 Anogenital distance of F1 and F2 rat pups on postnatal day 4 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of litters	Male		Female	
			AGD (mm)	AGD/(BW) ^{1/3}	AGD (mm)	AGD/(BW) ^{1/3}
F1	Control	24	Mean	5.37	2.49	(23)
			S.D.	0.41	0.11	1.22
	HBCD 150 ppm	21	Mean	5.44	2.48	2.67
			S.D.	0.36	0.10	1.23
F1	HBCD 1500 ppm	20	Mean	5.38	2.44	2.62
			S.D.	0.32	0.12	1.20
	HBCD 15000 ppm	18	Mean	5.20	2.46	2.57
			S.D.	0.51	0.14	1.23
F2	Control	22	Mean	5.12	2.46	2.69
			S.D.	0.54	0.12	1.30
	HBCD 150 ppm	22	Mean	5.12	2.44	2.71
			S.D.	0.41	0.13	1.33
F2	HBCD 1500 ppm	20	Mean	5.04	2.43	2.71
			S.D.	0.42	0.08	1.32
	HBCD 15000 ppm	20	Mean	(19)	(19)	2.54
			S.D.	4.84	2.42	1.32

Values in parentheses are the number of litters examined.

AGD: Anogenital distance.

BW: Body weight.

Table 37 Physical development in F1 and F2 rat pups treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of litters	Male			Female		
			Pinna detachment (%)	Incisor eruption (%)	Eye opening (%)	Pinna detachment (%)	Incisor eruption (%)	Eye opening (%)
F1	Control	24	Mean	86.0	(23)	(23)	(23)	(23)
			S.D.	26.5	91.6 17.6	48.2 41.5	85.8 29.5	94.9 11.4
	HBGD 150 ppm	21	Mean	92.5	96.4	56.7	94.7	66.7
			S.D.	16.5	12.0	37.9	14.7	10.1
	HBCD 1500 ppm	20	Mean	93.6	92.1	77.1 *	97.3	92.5
			S.D.	15.7	17.0	36.3	7.5	20.0
	HBCD 15000 ppm	18	Mean	81.3	(17)	(17)	(17)	(17)
			S.D.	27.9	89.7 19.9	45.8 34.6	86.4 23.8	92.2 15.4
F2	Control	23	Mean	(22)	(22)	(22)	(21)	(21)
			S.D.	79.9 36.4	86.4 25.3	72.7 40.0	73.6 39.6	85.7 26.9
	HBGD 150 ppm	22	Mean	90.5	92.8	62.5	90.6	90.9
			S.D.	22.8	19.6	40.6	22.8	26.2
	HBCD 1500 ppm	20	Mean	82.1	(18)	(18)	97.5	53.8 *
			S.D.	29.8	97.2 11.8	47.2 44.8	81.5 31.1	11.2
	HBCD 15000 ppm	21	Mean	(20)	(14)	(14)	(15)	(13)
			S.D.	70.1 39.2	86.3 27.7	33.9 ** 34.7	66.8 40.9	90.0 28.0

Pinna detachment, incisor eruption and eye opening were observed on postnatal day 3, 11 and 14, respectively.

Values in parentheses are the number of litters examined.

*: Significantly different from the control at $p \leq 0.05$ by Wilcoxon rank-sum test.

**: Significantly different from the control at $p \leq 0.01$ by Wilcoxon rank-sum test.

Table 38 Reflex response tests in F1 and F2 rat pups treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of animals	Male						Female					
			Surface righting reflex		Negative geotaxis reflex		Mid-air righting reflex		Surface righting reflex		Negative geotaxis reflex		Mid-air righting reflex	
			Success rate (%)	Response time (sec)	Completion ratio (%)	Response time (sec)	Success rate (%)	Response time (sec)	Success rate (%)	Response time (sec)	Completion ratio (%)	Response time (sec)	Success rate (%)	Response time (sec)
F1	Control	24	Mean	100.0	2.3	24/24	17.7	100.0	(23)	(23)	(23)	(23)	(23)	(23)
			S.D.	0.0	1.1	(100.0)	7.1	0.0		0.0	23/23	13.9	100.0	0.0
	HBCD 150 ppm	21	Mean	100.0	2.0	21/21	16.8	100.0	100.0	2.4	21/21	11.5	100.0	0.0
			S.D.	0.0	0.6	(100.0)	8.0	0.0	0.0	1.5	(100.0)	6.2	100.0	
	HBCD 1500 ppm	20	Mean	100.0	1.8	20/20	15.2	100.0	100.0	2.9	20/20	12.7	100.0	0.0
			S.D.	0.0	0.5	(100.0)	7.8	0.0	0.0	2.6	(100.0)	6.3	100.0	
	HBCD 15000 ppm	17	Mean	100.0	1.6 ^{ss}	17/17	19.4	100.0	100.0	2.6	17/17	17.0	100.0	0.0
			S.D.	0.0	0.3	(100.0)	5.9	0.0	0.0	2.6	(100.0)	6.9	100.0	
F2	Control	22	Mean	100.0	2.1	22/22	17.3	100.0	100.0	2.3	21/21	(21)	(21)	100.0
			S.D.	0.0	1.7	(100.0)	8.6	0.0	0.0	0.9	(100.0)	5.3	100.0	0.0
	HBCD 150 ppm	22	Mean	100.0	2.0	21/22	14.7	100.0	100.0	2.4	22/22	12.0	100.0	0.0
			S.D.	0.0	1.5	(95.5)	6.8	0.0	0.0	1.7	(100.0)	5.2	100.0	
	HBCD 1500 ppm	20	Mean	100.0	2.8	19/19	15.2	98.2	100.0	2.1	20/20	16.7	95.0	0.0
			S.D.	0.0	2.5	(100.0)	6.4	7.8	0.0	0.9	(100.0)	6.4	16.3	
	HBCD 15000 ppm	19	Mean	100.0	2.2	13/16	14.1	100.0	88.9	(18)	(16)	(15)	(13)	79.5 *
			S.D.	0.0	2.3	(81.3)	6.7	0.0	32.3	3.7	(88.2)	6.6	39.8	

Surface righting reflex, negative geotaxis reflex and mid-air righting reflex were examined on postnatal day 5, 8 and 18, respectively.

Completion ratio (%) = (number of animals showing positive response/number of animals examined) x 100.

Values in parentheses are the number of animals examined.

*: Significantly different from the control at $p \leq 0.05$ by Wilcoxon rank-sum test.

^{ss}: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

Table 39 Autopsy findings in F1 and F2 male rat pups euthanized on postnatal day 4 or found dead on postnatal days 0-4 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of litters examined	19	19	16	16
	Number of pups examined	60	50	37	63 ^b
	Number of pups with abnormal findings	0 (0.00)	1 (1.32)	1 (1.25)	2 (4.17)
	Findings ^a				
	Lung: White mass	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)
	Liver: Fine yellowish white patch, multifocal	0 (0.00)	0 (0.00)	0 (0.00)	1 (2.08)
	Kidney: Dilatation, renal pelvis	0 (0.00)	0 (0.00)	1 (1.25)	0 (0.00)
	External: Rudimentary tail	0 (0.00)	1 (1.32)	0 (0.00)	0 (0.00)
F2	Number of litters examined	19	21	17	17
	Number of pups examined	66	67	58 ^c	40
	Number of pups with abnormal findings	1 (2.63)	1 (1.59)	3 (3.43)	1 (1.96)
	Findings ^a				
	Liver: Yellowish white discoloration	0 (0.00)	0 (0.00)	0 (0.00)	1 (1.96)
	Digestive tract: Retention, gas	1 (2.63)	1 (1.59)	3 (3.43)	0 (0.00)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of pups that showed abnormal findings.

b: Including pups that were euthanized because of maternal death during delivery.

c: Including one animal that was not distinguished its sex because of maternal cannibalism.

Table 40 Autopsy findings in F1 and F2 female rat pups euthanized on postnatal day 4 or found dead on postnatal days 0-4 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of litters examined	21	19	18	12
	Number of pups examined	57	61	70	43 ^b
	Number of pups with abnormal findings	1 (1.19)	3 (3.38)	2 (3.70)	3 (7.92)
	Findings ^a				
	Liver: Yellowish brown discoloration	1 (1.19)	0 (0.00)	0 (0.00)	0 (0.00)
	Dark red mass, multifocal	0 (0.00)	1 (0.88)	0 (0.00)	0 (0.00)
	Jejunum: Obliteration	0 (0.00)	0 (0.00)	0 (0.00)	1 (4.17)
	Kidney: Dilatation, renal pelvis	0 (0.00)	2 (2.51)	2 (3.70)	1 (2.08)
	Abdominal cavity: Ascites	0 (0.00)	0 (0.00)	0 (0.00)	1 (4.17)
	External: Kinked tail	0 (0.00)	0 (0.00)	0 (0.00)	1 (1.67)
F2	Number of litters examined	18	22	13	18
	Number of pups examined	48	68	43	55
	Number of pups with abnormal findings	3 (4.17)	4 (6.44)	2 (5.77)	0 (0.00)
	Findings ^a				
	Digestive tract: Retention, gas	2 (2.78)	4 (6.44)	2 (5.77)	0 (0.00)
	Kidney: Dilatation, renal pelvis	1 (1.39)	0 (0.00)	0 (0.00)	0 (0.00)

Values in parentheses represent the means of incidences of pups with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of pups that showed abnormal findings.

b: Including pups that were euthanized because of maternal death during delivery.

Table 41 Autopsy findings in F1 and F2 male rat weanlings euthanized on postnatal day 26 or animals found dead on postnatal days 5-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of litters examined	24	21	20	17
	Number of animals examined	72	57	51	51
	Number of animals with abnormal findings	3 (4.17)	2 (3.17)	7 (12.50)	11 (22.16)*
	Findings ^a				
	Ileum: Diverticulum	0 (0.00)	0 (0.00)	1 (1.67)	0 (0.00)
	Kidney: Dilatation, renal pelvis	3 (4.17)	2 (3.17)	6 (10.83)	11 (22.16)*
F2	Number of litters examined	22	22	19	16
	Number of animals examined	75	81	64	52
	Number of animals with abnormal findings	1 (1.14)	2 (2.27)	4 (5.26)	3 (4.69)
	Findings ^a				
	Eyeball: Swelling	0 (0.00)	0 (0.00)	1 (1.32)	1 (1.56)
	Opacity	0 (0.00)	0 (0.00)	0 (0.00)	1 (1.56)
	Ileum: Diverticulum	1 (1.14)	0 (0.00)	3 (3.95)	1 (1.56)
	Kidney: Dilatation, renal pelvis	0 (0.00)	2 (2.27)	0 (0.00)	0 (0.00)

Values in parentheses represent the means of incidences of animals with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of animals that showed abnormal findings.

*: Significantly different from the control at $p \leq 0.05$ by Wilcoxon rank-sum test.

Table 42 Autopsy findings in F1 and F2 female rat weanlings euthanized on postnatal day 26 or animals found dead on postnatal days 5-26 treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of litters examined	23	21	20	14
	Number of animals examined	67	63	57	35
	Number of animals with abnormal findings	0 (0.00)	2 (3.17)	0 (0.00)	3 (7.14)
	Findings ^a				
	Spleen: Hypertrophy	0 (0.00)	1 (1.59)	0 (0.00)	0 (0.00)
	Kidney: Dilatation, renal pelvis	0 (0.00)	1 (1.59)	0 (0.00)	3 (7.14)
F2	Number of litters examined	22	22	20	19
	Number of animals examined	78	81	71	53
	Number of animals with abnormal findings	2 (2.27)	1 (1.14)	2 (2.50)	0 (0.00)
	Findings ^a				
	Ileum: Diverticulum	1 (1.14)	1 (1.14)	0 (0.00)	0 (0.00)
	Kidney: Dilatation, renal pelvis	1 (1.14)	0 (0.00)	2 (2.50)	0 (0.00)

Values in parentheses represent the means of incidences of animals with abnormal findings (%), the litter is the unit evaluated).

a: Values represent the number of animals that showed abnormal findings.

8117

Table 43 Absolute and relative organ weights in F1 and F2 male rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of animals	Body weight		Brain		Thymus		Liver		Kidney ^a		Spleen	
			g	g	%		mg	10 ⁻³ %	g	%	mg	10 ⁻³ %	mg	10 ⁻³ %
F1	Control	23	Mean	85.7	1.64	1.94	342	398	3.94	4.60	996	1165	336	394
			S.D.	10.9	0.09	0.19	68	55	0.63	0.37	125	74	62	64
	HBCD 150 ppm	21	Mean	89.6	1.66	1.87	339	379	4.12	4.60	1035	1155	327	366
			S.D.	8.1	0.05	0.17	50	45	0.48	0.32	131	92	41	42
	HBCD 1500 ppm	20	Mean	87.7	1.62	1.86	369	421	4.43 *	5.05 **	1004	1146	334	383
			S.D.	9.2	0.07	0.18	59	55	0.59	0.32	109	70	43	46
	HBCD 15000 ppm	17	Mean	78.3 *	1.55 **	1.99	317	405	4.71 **	6.00 **	894 *	1140	309	395
			S.D.	5.8	0.06	0.13	57	70	0.58	0.44	99	78	69	81
F2	Control	22	Mean	82.2	1.62	2.08	343	414	3.87	4.72	965	1201	360	443
			S.D.	17.1	0.13	0.58	92	97	0.90	0.59	167	173	83	77
	HBCD 150 ppm	22	Mean	84.6	1.65	1.96	336	397	4.02	4.74	958	1134 **	361	429
			S.D.	8.7	0.08	0.16	57	54	0.55	0.35	99	56	54	64
	HBCD 1500 ppm	18	Mean	81.3	1.60	2.01	360	441	4.12	5.04 §	933	1155	346	426
			S.D.	13.4	0.10	0.29	88	69	0.83	0.40	135	85	78	69
	HBCD 15000 ppm	13	Mean	64.7 **	1.46 **	2.31 **	282	434	3.88	6.00 **	749 **	1170	263 **	411
			S.D.	11.2	0.09	0.33	71	81	0.68	0.25	100	96	50	66

(to be continued)

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

§: Significantly different from the control at $p \leq 0.05$ by Mann-Whitney U-test.

**: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

Table 43 (continued) Absolute and relative organ weights in F1 and F2 male rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of animals	Adrenal ^a		Testis ^a		Epididymis ^a		Prostate		
			mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	
F1	Control	23	Mean	23.9	28.0	488	565	73.2	85.9	40.0	46.4
			S.D.	3.0	2.6	100	65	9.5	9.8	12.0	10.3
	HBCD 150 ppm	21	Mean	25.0	28.0	550 *	614 *	77.4	86.7	42.0	47.1
			S.D.	3.3	3.9	70	56	9.8	10.3	7.7	8.8
F1	HBCD 1500 ppm	20	Mean	26.1	29.9	541	615 *	78.3	89.3	42.1	48.2
			S.D.	3.7	4.3	92	61	9.9	7.5	7.1	7.3
	HBCD 15000 ppm	17	Mean	22.8	29.2	494	631 **	70.1	89.9	34.8	44.5
			S.D.	3.6	4.8	70	73	11.6	15.3	9.4	11.1
F2	Control	22	Mean	23.4	28.7	476	574	73.7	90.7	40.6	50.2
			S.D.	5.1	4.4	138	123	16.8	14.1	9.7	9.3
	HBCD 150 ppm	22	Mean	25.1	29.7	510	600	73.6	87.2	42.3	50.2
			S.D.	3.6	3.2	81	55	10.7	10.6	9.5	10.7
F2	HBCD 1500 ppm	18	Mean	24.3	29.9	475	572	71.8	87.3	41.7	50.8
			S.D.	5.2	4.0	136	93	17.5	9.6	12.1	9.6
	HBCD 15000 ppm	13	Mean	19.6 *	30.4	385	589	61.7 ^s	96.2	29.5 **	47.3
			S.D.	3.2	2.0	92	54	9.5	10.5	6.8	15.8

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

^s: Significantly different from the control at $p \leq 0.05$ by Mann-Whitney U-test.

Table 44 Absolute and relative organ weights in F1 and F2 female rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of animals	Body weight		Brain		Thymus		Liver		Kidney ^a		Spleen	
			g	g	%	mg	10 ⁻³ %	g	%	mg	10 ⁻³ %	mg	10 ⁻³ %	
F1	Control	23	Mean	78.9	1.58	2.04	335	423	3.61	4.57	932	1189	311	399
			S.D.	10.6	0.09	0.23	64	58	0.55	0.35	102	85	53	75
	HBCD 150 ppm	21	Mean	83.2	1.61	1.96	330	397	3.83	4.59	945	1136	306	370
			S.D.	9.7	0.07	0.19	58	63	0.55	0.28	112	63	44	51
	HBCD 1500 ppm	20	Mean	83.9	1.59	1.91	370	441	4.22 **	5.02 **	958	1143	304	363
			S.D.	8.3	0.08	0.14	58	53	0.56	0.32	115	81	59	67
	HBCD 15000 ppm	14	Mean	72.1	1.51 *	2.10	305	422	4.37 **	6.07 **	815 **	1129	280	388
			S.D.	5.3	0.06	0.16	31	33	0.41	0.36	85	72	40	48
F2	Control	21	Mean	75.3	1.57	2.14	338	447	3.55	4.70	916	1226	325	436
			S.D.	12.5	0.11	0.37	85	81	0.64	0.27	131	93	59	61
	HBCD 150 ppm	22	Mean	75.8	1.58	2.11	324	429	3.57	4.70	885	1169	302	399
			S.D.	8.5	0.07	0.20	50	57	0.48	0.28	98	65	42	43
	HBCD 1500 ppm	20	Mean	73.1	1.55	2.17	331	451	3.63	4.94	868	1194	299	412
			S.D.	12.8	0.12	0.35	69	51	0.74	0.32	144	84	62	61
	HBCD 15000 ppm	13	Mean	57.9 **	1.41 **	2.48 **	260 **	445	3.42	5.89 **	679 **	1177	225 **	392
			S.D.	11.6	0.15	0.34	80	83	0.77	0.44	138	103	45	53

(to be continued)

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Mann-Whitney U-test.

Table 44 (continued) Absolute and relative organ weights in F1 and F2 female rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Group	Number of animals	Adrenal ^a		Ovary ^a		Uterus		
			mg	10 ⁻³ %	mg	10 ⁻³ %	mg	10 ⁻³ %	
F1	Control	23	Mean	21.9	27.8	20.8	26.5	57.0	73.6
			S.D.	3.5	3.8	3.7	4.5	10.9	17.5
	HBCD 150 ppm	21	Mean	23.7	28.7	22.8	27.5	62.0	74.9
			S.D.	2.8	4.0	3.6	4.1	14.1	17.7
	HBCD 1500 ppm	20	Mean	24.2	28.9	21.0	25.0	64.1	76.0
			S.D.	3.8	4.0	4.0	3.8	18.6	18.4
	HBCD 15000 ppm	14	Mean	20.9	28.9	20.9	28.9	51.9	71.9
			S.D.	3.4	4.1	3.4	3.7	12.4	16.2
F2	Control	21	Mean	22.1	29.5	20.0	26.9	60.8	80.9
			S.D.	4.2	4.1	3.9	5.1	16.1	16.3
	HBCD 150 ppm	22	Mean	21.5	28.4	22.9 *	30.5 ^s	63.6	84.4
			S.D.	2.6	3.4	2.6	3.9	15.1	21.0
	HBCD 1500 ppm	20	Mean	21.5	29.4	20.9	28.8	57.0	78.7
			S.D.	4.3	3.1	3.9	4.2	15.7	21.7
	HBCD 15000 ppm	13	Mean	17.6 **	30.7	18.2	32.1 ^s	47.6 *	83.7
			S.D.	3.1	2.6	4.0	7.5	11.4	20.3

a: Values represent the total weights of the organs of both sides.

*: Significantly different from the control at $p \leq 0.05$ by Dunnett's test.

**: Significantly different from the control at $p \leq 0.01$ by Dunnett's test.

^s: Significantly different from the control at $p \leq 0.05$ by Mann-Whitney U-test.

Table 45 Histopathological findings in F1 and F2 male rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of weanlings examined ^a	23	21	20	17
	Number of weanlings with abnormal findings	0	0	0	1
Findings^b					
	Liver: Necrosis, focal	0	0	0	1
F2	Number of weanlings examined ^a	22	22	18	13
	Number of weanlings with abnormal findings	0	0	0	0

a: Weanlings were examined on the thyroid in the all groups and on the liver in the control and 15000 ppm groups.

b: Values represent the number of animals that showed abnormal findings.

922

Table 46 Histopathological findings in F1 and F2 female rat weanlings treated with 1,2,5,6,9,10-hexabromocyclododecane (HBCD) in the two-generation reproductive toxicity study (SR04222)

Gener- ation	Item	Control	HBCD (ppm)		
			150	1500	15000
F1	Number of weanlings examined ^a	23	21	20	14
	Number of weanlings with abnormal findings	0	0	0	0
F2	Number of weanlings examined ^a	21	22	20	13
	Number of weanlings with abnormal findings	0	0	0	0

a: Weanlings were examined on the thyroid in the all groups and on the liver in the control and 15000 ppm groups.