

Purpose

- To identify areas requiring emergency measures in response to the accident at Tokyo Electric Power Company (TEPCO)'s Fukushima Daiichi NPS
- To estimate exposure doses for the first one year after the accident for that purpose
- To assess health risks of people in Japan and the whole world based on the estimated doses

Assessment method

- Set conservative conditions for dose estimation and assess exposure doses
- Estimate doses both from internal and external exposure
- Estimate exposure doses by age (one year old (infants), 10 years old (children), and 20 years old (adults)) and by area

The WHO is an organization responsible for assessing health risks posed by radiation in an emergency. Therefore, after the accident at TEPCO's Fukushima Daiichi NPS, it conducted assessment of exposure doses for the first one year regarding people in Japan and the whole world for the purpose of identifying areas and groups of people for which emergency measures should be taken.

The WHO assessed doses due to exposure to radiation via four pathways: (i) external exposure from the ground surface, (ii) external exposure from radioactive plumes (p.29 of Vol. 1, "Effects of Reactor Accidents"), (iii) internal exposure through inhalation, and (iv) internal exposure through ingestion. Doses due to external exposure via (i) and (ii) and internal exposure via (iii) were estimated through simulation based on information on contamination density on the soil surface as of September 2011, while doses due to internal exposure via (iv) were estimated based on the measurement values for foods and drinking water.

People's exposure doses are to be calculated by summing up estimated values for (i) to (iv), but in order to avoid underestimation, the WHO set conservative assumptions and calculated the largest exposure doses imaginable. Concretely, the WHO adopted the preconditions that protective measures such as deliberate evacuation, sheltering indoors, or shipping restrictions on foods were not at all taken.

As exposure doses vary by area and age, the WHO estimated doses by dividing areas into Fukushima Prefecture, neighboring prefectures (Chiba, Gunma, Ibaraki, Miyagi and Tochigi Prefectures), the rest of Japan, neighboring countries and the rest of the world, and by dividing people by age into those aged one year old (infants), 10 years old (children), and 20 years old (adults) at the time of the accident.

Included in this reference material on March 31, 2015

Updated on March 31, 2023