

WHO Reports (1/2): Outline of the WHO's Dose Assessment [Reference]

Purpose

- To identify areas requiring emergency measures in response to the accident at Tokyo Electric Power Company (TEPCO)'s Fukushima Daiichi NPS
- To estimate exposure doses for the first one year after the accident for that purpose
- To assess health risks of people in Japan and the whole world based on the estimated doses

Assessment method

- Set conservative conditions for dose estimation and assess exposure doses
- Estimate doses both from internal and external exposure
- Estimate exposure doses by age (one year old (infants), 10 years old (children), and 20 years old (adults)) and by area

Assumptions for risk assessment

- Assuming that there is no threshold dose for radiation carcinogenesis, the linear model and the linear quadratic model were adopted for dose-response relationships for solid cancer and leukemia, respectively.
- Dose and dose-rate effectiveness factors (DDREF) were not applied.

Results

- People's exposure doses were below all thresholds of deterministic effects (tissue reactions).
- Even in the area where the highest exposure dose was estimated, no significant increase would be observed in risks of childhood thyroid cancer and other types of cancer or leukemia and increased incidence of these diseases exceeding natural variation is hardly expected.
- Risks of heritable effects due to radiation exposure are further smaller than the risks of generating cancer.
- The results suggest that increases in the incidence of diseases attributable to the additional radiation exposure are likely to remain below detectable levels.

Conclusion

- Values in this Report are for roughly ascertaining current risk levels and are not intended to predict future health effects.