

平成 27～31 年度に対象自治体等から報告された結果の取りまとめ データ編

(1) 参加者¹数

参加者の属性は表 1 に、各検査等の項目別参加者数は表 2 に示すとおりである。

<表 1>

参加者は 60～79 歳の者が多く、男性の方が女性よりも多かった。

<表 2>

実施方法①と②を合わせると、延べ人数で石綿ばく露の聴取を受けた者は 11,012 人、CT 検査を受けた者は 8,579 人（表 2 - 1、表 2 - 2 を合計、以下同様）、肺がん検診等の既存検診を受診しエックス線検査画像の読影を行った人数は 5,556 人、保健指導を受けた者は 6,996 人であった。

表 1 参加者の属性(年齢階層別、性別、平成 27～31 年度延べ人数)

(単位:人)

	男性		女性		合計	
40 歳未満	36	0.5%	41	0.9%	77	0.7%
40～49 歳	421	6.4%	280	6.3%	701	6.4%
50～59 歳	762	11.6%	608	13.7%	1,370	12.4%
60～69 歳	1,893	28.7%	1,394	31.5%	3,287	29.8%
70～79 歳	2,697	40.9%	1,642	37.1%	4,339	39.4%
80～89 歳	750	11.4%	444	10.0%	1,194	10.8%
90 歳以上	31	0.5%	13	0.3%	44	0.4%
合 計	6,590	100.0%	4,422	100.0%	11,012	100.0%

※ 小数点以下第2位を四捨五入したため、合計が 100%にならない場合がある

¹ 調査対象者のうち「石綿ばく露の聴取」を受けた者を「参加者」とする。

表2-1 項目別参加者数((A)実施方法①、平成 27~31 年度延べ人数)

(単位:人)

		石綿ばく露 の聴取	CT検査				保健指導	
			うちエックス線検査					
		人	人	%	人	%	人	%
埼玉県	さいたま市	233	220	94.4%	191	82.0%	220	94.4%
神奈川県	横浜市鶴見区	316	308	97.5%	225	71.2%	316	100.0%
岐阜県	羽島市	947	674	71.2%	728	76.9%	91	9.6%
大阪府	大阪市	281	273	97.2%	0	0.0%	273	97.2%
	堺市	211	192	91.0%	161	76.3%	34	16.1%
	岸和田市	103	84	81.6%	36	35.0%	96	93.2%
	貝塚市	49	49	100.0%	0	0.0%	48	98.0%
	八尾市	51	51	100.0%	0	0.0%	51	100.0%
	泉佐野市	96	84	87.5%	0	0.0%	61	63.5%
	河内長野市	392	385	98.2%	294	75.0%	392	100.0%
	和泉市	18	18	100.0%	0	0.0%	18	100.0%
	東大阪市	64	29	45.3%	29	45.3%	56	87.5%
	泉南市	252	249	98.8%	7	2.8%	195	77.4%
	阪南市	189	189	100.0%	10	5.3%	189	100.0%
	熊取町	9	9	100.0%	0	0.0%	9	100.0%
	田尻町	1	1	100.0%	0	0.0%	1	100.0%
	岬町	37	37	100.0%	0	0.0%	36	97.3%
兵庫県	尼崎市	2,943	1,814	61.6%	2,031	69.0%	2,943	100.0%
	西宮市	254	161	63.4%	96	37.8%	161	63.4%
	芦屋市	46	46	100.0%	46	100.0%	46	100.0%
	加古川市	121	112	92.6%	112	92.6%	112	92.6%
	宝塚市	4	4	100.0%	4	100.0%	4	100.0%
奈良県		2,229	2,157	96.8%	74	3.3%	175	7.9%
福岡県	北九州市門司区	517	317	61.3%	321	62.1%	516	99.8%
佐賀県	鳥栖市	175	118	67.4%	173	98.9%	175	100.0%
転居者調査		464	345	74.4%	8	1.7%	78	16.8
合計		10,002	7,926	79.2%	4,546	45.5%	6,296	62.9%

※ 割合は、石綿ばく露の聴取を実施した者に対する割合

※ 小数点以下第2位を四捨五入

※ さいたま市は平成 29 年度のみ中央区・大宮区だけを対象地域とした

表2－2 項目別参加者数((B)実施方法②、平成 29～31 年度)

(単位:人)

			石綿ばく露 の聴取	エックス検査				保健指導	
				うちCT線検査					
			人	人	%	人	%	人	%
東京都	大田区	H31	18	18	100.0%	3	16.7%	18	100.0%
兵庫県	神戸市	H29	198	198	100.0%	116	58.6%	124	62.6%
		H30	651	651	100.0%	441	67.7%	459	70.5%
		H31	143	143	100.0%	93	65.0%	99	69.2%
		計	992	992	100.0%	650	65.5%	682	68.8%
合計			1,010	1,010	100.0%	653	64.7%	700	69.3%

※ 割合は、石綿ばく露の聴取を実施した者に対する割合

※ 小数点以下第2位を四捨五入

※ (B)実施方法②は平成 29 年度より実施

（２）有所見者数

各検査方法によって確認された石綿関連所見（疑いを含む。）の有所見者数は表 3、平成 31 年度にＣＴ検査を受けた継続者のうち、過去もＣＴ検査を受けた人数は表 4、各検査方法によって確認された石綿関連所見（疑いを含む。）と年齢階層・性別の関係及びばく露・性別の関係は表 5 に示すとおりである。

表３－１ 検査によって確認された石綿関連所見の有所見者数

((A)実施方法①、平成 27～31 年度)

	合計			新規(リスク調査含)			継続(リスク調査含)		
	CT検査 受診者	有所見者		CT検査 受診者	有所見者		CT検査 受診者	有所見者	
平成 27 年度	1,566	604	38.6%	－	－	－	－	－	－
平成 28 年度	1,481	563	38.0%	375	106	28.3%	1,106	457	41.3%
平成 29 年度	1,659	618	37.3%	394	126	32.0%	1,265	492	38.9%
平成 30 年度	1,610	652	40.5%	340	119	35.0%	1,270	533	42.0%
平成 31 年度	1,610	645	40.1%	299	95	31.8%	1,311	550	42.0%

※過去に試行調査又はリスク調査において CT 検査を受診していない者を新規、
過去に試行調査又はリスク調査において１度でも CT 検査を実施した者を継続とした

表３－２ 検査によって確認された石綿関連所見の自治体別有所見者数

((B)実施方法②、平成 29～31 年度)

(単位:人)

		合計			新規(リスク調査含)			継続(リスク調査含)		
		検査 受診者	有所見者		検査 受診者	有所見者		検査 受診者	有所見者	
平成 29 年度	胸部 X 線 のみ	82	2	2.4%	82	2	2.4%	0	0	－
	胸部 X 線 及び胸部 CT	116	11	9.5%	116	11	9.5%	0	0	－
	計	198	13	6.6%	198	13	6.6%	0	0	－
平成 30 年度	胸部 X 線 のみ	210	15	7.1%	203	15	7.4%	7	0	0.0%
	胸部 X 線 及び胸部 CT	441	54	12.2%	441	54	12.2%	0	0	－
	計	651	69	10.6%	644	69	10.7%	7	0	0.0%
平成 31 年度	胸部 X 線 のみ	65	6	9.2%	60	6	10.0%	5	0	0.0%
	胸部 X 線 及び胸部 CT	96	15	15.6%	87	13	14.9%	9	2	22.2%
	計	161	21	13.0%	147	19	12.9%	14	2	14.3%

(表3-1 参考) 検査によって確認された石綿関連所見の自治体別有所見者数
((A)実施方法①、平成 28 年度)

(単位:人)

		合計			新規(リスク調査含)			継続(リスク調査含)		
		CT検査 受診者	有所見者		CT検査 受診者	有所見者		CT検査 受診者	有所見者	
神奈川県	横浜市鶴見区	35	14	40.0%	27	7	25.9%	8	7	87.5%
岐阜県	羽島市	98	62	63.3%	1	1	100.0%	97	61	62.9%
大阪府	大阪市	53	24	45.3%	8	1	12.5%	45	23	51.1%
	堺市	26	11	42.3%	21	8	38.1%	5	3	60.0%
	岸和田市	15	6	40.0%	1	0	0.0%	14	6	42.9%
	貝塚市	11	3	27.3%	3	0	0.0%	8	3	37.5%
	八尾市	13	3	23.1%	13	3	23.1%	0	0	－
	泉佐野市	15	3	20.0%	1	0	0.0%	14	3	21.4%
	河内長野市	85	15	17.6%	6	1	16.7%	79	14	17.7%
	和泉市	2	0	0.0%	2	0	0.0%	0	0	－
	東大阪市	11	0	0.0%	11	0	0.0%	0	0	－
	泉南市	42	18	42.9%	9	6	66.7%	33	12	36.4%
	阪南市	35	7	20.0%	7	0	0.0%	28	7	25.0%
	熊取町	1	1	100.0%	1	1	100.0%	0	0	－
	田尻町	1	0	0.0%	0	0	－	1	0	0.0%
	岬町	7	3	42.9%	2	0	0.0%	5	3	60.0%
兵庫県	尼崎市	326	124	38.0%	117	31	26.5%	209	93	44.5%
	西宮市	43	29	67.4%	35	23	65.7%	8	6	75.0%
	芦屋市	12	0	0.0%	6	0	0.0%	6	0	0.0%
	加古川市	58	14	24.1%	58	14	24.1%	0	0	－
奈良県		451	169	37.5%	8	3	37.5%	443	166	37.5%
福岡県	北九州市門司区	56	18	32.1%	25	3	12.0%	31	15	48.4%
佐賀県	鳥栖市	29	23	79.3%	1	1	100.0%	28	22	78.6%
転居者調査		56	16	28.6%	12	3	25.0%	44	13	29.5%
合計		1,481	563	38.0%	375	106	28.3%	1,106	457	41.3%

※過去に試行調査又はリスク調査において CT 検査を受診していない者を新規、
過去に試行調査又はリスク調査において1度でも CT 検査を実施した者を継続とした

(表3-1 参考) 検査によって確認された石綿関連所見の自治体別有所見者数
((A)実施方法①、平成 29 年度)

(単位:人)

		合計			新規(リスク調査含)			継続(リスク調査含)		
		CT検査 受診者	有所見者		CT検査 受診者	有所見者		CT検査 受診者	有所見者	
埼玉県	さいたま市 中央区・大宮区	84	27	32.1%	84	27	32.1%	0	0	-
神奈川県	横浜市鶴見区	62	34	54.8%	34	10	29.4%	28	24	85.7%
岐阜県	羽島市	168	91	54.2%	2	1	50.0%	166	90	54.2%
大阪府	大阪市	55	19	34.5%	6	2	33.3%	49	17	34.7%
	堺市	45	14	31.1%	33	7	21.2%	12	7	58.3%
	岸和田市	20	7	35.0%	0	0	-	20	7	36.8%
	貝塚市	9	2	22.2%	0	0	-	9	2	22.2%
	八尾市	14	2	14.3%	13	2	15.4%	1	0	0.0%
	泉佐野市	18	6	33.3%	0	0	-	18	6	33.3%
	河内長野市	71	8	11.3%	0	0	-	71	8	11.3%
	和泉市	9	1	11.1%	9	1	11.1%	0	0	-
	東大阪市	3	0	0.0%	2	0	0.0%	1	0	0.0%
	泉南市	46	18	39.1%	5	1	20.0%	41	17	41.5%
	阪南市	40	6	15.0%	3	0	0.0%	37	6	16.2%
	熊取町	0	0	-	0	0	-	0	0	-
	田尻町	0	0	-	0	0	-	0	0	-
	岬町	6	3	50.0%	2	0	0.0%	4	3	75.0%
兵庫県	尼崎市	347	149	42.9%	133	52	39.1%	214	97	45.3%
	西宮市	31	15	48.4%	21	12	57.1%	10	3	30.0%
	芦屋市	9	0	0.0%	1	0	0.0%	8	0	0.0%
	加古川市	18	5	27.8%	14	3	21.4%	4	2	50.0%
奈良県		441	153	34.7%	6	0	0.0%	435	153	35.2%
福岡県	北九州市門司区	83	20	24.1%	16	3	18.8%	67	17	25.4%
佐賀県	鳥栖市	23	20	87.0%	1	1	100.0%	22	19	86.4%
転居者調査		57	18	31.6%	9	4	44.4%	48	14	29.2%
合計		1,659	618	37.3%	394	126	32.0%	1,265	492	38.9%

(表3-1 参考) 検査によって確認された石綿関連所見の自治体別有所見者数
((A)実施方法①、平成30年度)

(単位:人)

		合計			新規(リスク調査含)			継続(リスク調査含)		
		CT検査 受診者	有所見者		CT検査 受診者	有所見者		CT検査 受診者	有所見者	
埼玉県	さいたま市	75	22	29.3%	72	20	27.8%	3	2	66.7%
神奈川県	横浜市鶴見区	66	48	72.7%	30	17	56.7%	36	31	86.1%
岐阜県	羽島市	132	79	59.8%	4	2	50.0%	128	77	60.2%
大阪府	大阪市	56	18	32.1%	10	3	30.0%	46	15	32.6%
	堺市	47	19	40.4%	16	6	37.5%	31	13	41.9%
	岸和田市	18	6	33.3%	0	0	—	18	6	33.3%
	貝塚市	8	4	50.0%	0	0	—	8	4	50.0%
	八尾市	18	3	16.7%	13	1	7.7%	5	2	40.0%
	泉佐野市	18	7	38.9%	1	1	100.0%	17	6	35.3%
	河内長野市	75	15	20.0%	1	0	0.0%	74	15	20.3%
	和泉市	4	0	0.0%	4	0	0.0%	0	0	—
	東大阪市	7	2	28.6%	2	1	50.0%	5	1	20.0%
	泉南市	47	14	29.8%	9	2	22.2%	38	12	31.6%
	阪南市	39	7	17.9%	3	2	66.7%	36	5	13.9%
	熊取町	2	1	50.0%	2	1	50.0%	0	0	—
	田尻町	0	0	—	0	0	—	0	0	—
	岬町	9	3	33.3%	3	1	33.3%	6	2	33.3%
兵庫県	尼崎市	388	184	47.4%	106	41	38.7%	282	143	50.7%
	西宮市	27	11	40.7%	12	5	41.7%	15	6	40.0%
	芦屋市	5	0	0.0%	1	0	0.0%	4	0	0.0%
	加古川市	19	6	31.6%	18	5	27.8%	1	1	100.0%
奈良県		414	152	36.7%	8	3	37.5%	406	149	36.7%
福岡県	北九州市門司区	50	14	28.0%	2	0	0.0%	48	14	29.2%
佐賀県	鳥栖市	20	16	80.0%	3	1	33.3%	17	15	88.2%
転居者調査		66	21	31.8%	20	7	35.0%	46	14	30.4%
合計		1,610	652	40.5%	340	119	35.0%	1,270	533	42.0%

(表3-1 参考) 検査によって確認された石綿関連所見の自治体別有所見者数
((A)実施方法①、平成31年度)

(単位:人)

		合計			新規(リスク調査含)			継続(リスク調査含)		
		CT検査 受診者	有所見者		CT検査 受診者	有所見者		CT検査 受診者	有所見者	
埼玉県	さいたま市	61	23	37.7%	59	21	35.6%	2	2	100.0%
神奈川県	横浜市鶴見区	63	49	77.8%	17	9	52.9%	46	40	87.0%
岐阜県	羽島市	165	76	46.1%	4	1	25.0%	161	75	46.6%
大阪府	大阪市	58	21	36.2%	9	4	44.4%	49	17	34.7%
	堺市	28	10	35.7%	5	2	40.0%	23	8	34.8%
	岸和田市	14	6	42.9%	2	2	100.0%	12	4	33.3%
	貝塚市	10	3	30.0%	0	0	－	10	3	30.0%
	八尾市	6	3	50.0%	5	2	40.0%	1	1	100.0%
	泉佐野市	13	5	38.5%	0	0	－	13	5	38.5%
	河内長野市	57	11	19.3%	3	0	0.0%	54	11	20.4%
	和泉市	3	0	0.0%	3	0	0.0%	0	0	－
	東大阪市	8	2	25.0%	4	0	0.0%	4	2	50.0%
	泉南市	53	20	37.7%	12	5	41.7%	41	15	36.6%
	阪南市	38	10	26.3%	0	0	－	38	10	26.3%
	熊取町	4	1	25.0%	4	1	25.0%	0	0	－
	田尻町	0	0	－	0	0	－	0	0	－
	岬町	8	1	12.5%	2	0	0.0%	6	1	16.7%
兵庫県	尼崎市	409	199	48.7%	103	36	35.0%	306	163	53.3%
	西宮市	28	7	25.0%	15	2	13.3%	13	5	38.5%
	芦屋市	8	0	0.0%	1	0	0.0%	7	0	0.0%
	加古川市	17	2	11.8%	17	2	11.8%	0	0	－
	宝塚市	4	1	25.0%	4	1	25.0%	0	0	－
奈良県		385	138	35.8%	7	2	28.6%	378	136	36.0%
福岡県	北九州市門司区	84	17	20.2%	16	3	18.8%	68	14	20.6%
佐賀県	鳥栖市	21	21	100.0%	0	0	－	21	21	100.0%
転居者調査		65	19	29.2%	7	2	28.6%	58	17	29.3%
合計		1,610	645	40.1%	299	95	31.8%	1,311	550	42.0%

(表3-2 参考) 検査によって確認された石綿関連所見の自治体別有所見者数
((B)実施方法②、平成 29～31 年度)

(単位:人)

				合計			新規(リスク調査含)			継続(リスク調査含)		
				検査 受診者	有所見者		検査 受診者	有所見者		検査 受診者	有所見者	
東京都	大田区	H31	胸部 X 線 のみ	15	3	20.0%	15	3	20.0%	0	0	－
			胸部 X 線 及び胸部 CT	3	2	66.7%	3	2	66.7%	0	0	－
			計	18	5	27.8%	18	5	27.8%	0	0	－
兵庫県	神戸市	H29	胸部 X 線 のみ	82	2	2.4%	82	2	2.4%	0	0	－
			胸部 X 線 及び胸部 CT	116	11	9.5%	116	11	9.5%	0	0	－
			計	198	13	6.6%	198	13	6.6%	0	0	－
		H30	胸部 X 線 のみ	210	15	7.1%	203	15	7.4%	7	0	0.0%
			胸部 X 線 及び胸部 CT	441	54	12.2%	441	54	12.2%	0	0	－
			計	651	69	10.6%	644	69	10.7%	7	0	0.0%
		H31	胸部 X 線 のみ	50	3	6.0%	45	3	6.7%	5	0	0.0%
			胸部 X 線 及び胸部 CT	93	13	14.0%	84	11	13.1%	9	2	22.2%
			計	143	16	11.2%	129	14	10.9%	14	2	14.3%
		計	胸部 X 線 のみ	342	20	5.8%	330	20	6.1%	12	0	0.0%
			胸部 X 線 及び胸部 CT	650	78	12.0%	641	76	11.9%	9	2	22.2%
			計	992	98	9.9%	971	96	9.9%	21	2	9.5%
合計			胸部 X 線 のみ	357	23	6.4%	345	23	6.7%	12	0	0.0%
			胸部 X 線 及び胸部 CT	653	80	12.3%	644	78	12.1%	9	2	22.2%
			計	1,010	103	10.2%	989	101	10.2%	21	2	9.5%

表4-1 平成28年度にCT検査を受けた継続者のうち、過去もCT検査を受けた人数

(単位:人)

過去の 直近の検査結果	過去にCT検査を受けた直近の年 ^{※1}										計
	H18	H19	H20	H21	H22	H23	H24	H25	H26	H27	
所見 ^{※2} あり	0	1	0	0	2	2	3	4	73	486	571
所見 ^{※2} なし	1	2	9	3	2	8	12	6	95	397	535
計	1	3	9	3	4	10	15	10	168	883	1,106

表4-2 平成29年度にCT検査を受けた継続者のうち、過去もCT検査を受けた人数

(単位:人)

過去の 直近の検査結果	過去にCT検査を受けた直近の年 ^{※1}											計
	H18	H19	H20	H21	H22	H23	H24	H25	H26	H27	H28	
所見 ^{※2} あり	2	1	1	0	0	2	2	5	44	53	393	503
所見 ^{※2} なし	2	4	2	2	7	5	4	5	128	95	508	762
計	4	5	3	2	7	7	6	10	172	148	901	1,265

表4-3 平成30年度にCT検査を受けた継続者のうち、過去もCT検査を受けた人数

(単位:人)

過去の 直近の検査結果	過去にCT検査を受けた直近の年 ^{※1}												計
	H18	H19	H20	H21	H22	H23	H24	H25	H26	H27	H28	H29	
所見 ^{※2} あり	0	1	1	2	1	0	4	3	21	20	52	448	553
所見 ^{※2} なし	0	0	0	5	5	3	5	4	35	35	91	534	717
計	0	1	1	7	6	3	9	7	56	55	143	982	1,270

※1 平成26年度リスク調査参加者は原則全員CT検査受診としている。

※2 所見は胸水貯留、胸膜プラーク、びまん性胸膜肥厚、胸膜腫瘍、肺野の間質影、円形無気肺、肺野の腫瘤状陰影、リンパ節の腫大(①～⑧)

表4-4 平成31年度にCT検査を受けた継続者のうち、過去もCT検査を受けた人数

(単位:人)

過去の直近の 検査結果	過去にCT検査を受けた直近の年※1													
	H18	H19	H20	H21	H22	H23	H24	H25	H26	H27	H28	H29	H30	計
所見※2あり	2	0	2	1	2	5	4	5	84	33	41	152	896	1,227
所見※2なし	1	0	0	0	1	2	0	2	1	6	7	15	55	90
計	3	0	2	1	3	7	4	7	85	39	48	167	951	1,317

※1 平成26年度リスク調査参加者は原則全員CT検査受診としている。

※2 所見は胸水貯留、胸膜プラーク、びまん性胸膜肥厚、胸膜腫瘍、肺野の間質影、円形無気肺、肺野の腫瘤状陰影、リンパ節の腫大(①～⑧)

表5-1 年齢階層・性別検査所見((A)実施方法①、平成28～31年度延べ人数)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
CT検査 受診者	合計	6,360		35		367		787		1,816		2,553		779		23	
	男性	3,804		11		206		427		1,055		1,593		494		18	
	女性	2,556		24		161		360		761		960		285		5	
石綿関連 所見(疑い を含む)実 人数	合計	2478(151)	39.0%	1(0)	2.9%	30(3)	8.2%	173(27)	22.0%	657(38)	36.2%	1156(65)	45.3%	444(18)	57.0%	19(0)	82.6%
	男性	1715(106)	45.1%	1(0)	9.1%	12(1)	5.8%	110(21)	25.8%	432(22)	40.9%	840(49)	52.7%	305(13)	61.7%	16(0)	88.9%
	女性	763(45)	29.9%	0(0)	0.0%	18(2)	11.2%	63(6)	17.5%	225(16)	29.6%	316(16)	32.9%	139(5)	48.8%	3(0)	60.0%
①胸水貯 留	小計	22(1)	0.3%	0(0)	0.0%	0(0)	0.0%	2(1)	0.3%	3(0)	0.2%	6(0)	0.2%	8(0)	1.0%	3(0)	13.0%
	男性	19(0)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.3%	6(0)	0.4%	7(0)	1.4%	3(0)	16.7%
	女性	3(1)	0.1%	0(0)	0.0%	0(0)	0.0%	2(1)	0.6%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%
②胸膜ブ ラーク(胸 膜肥厚 斑)	小計	2157(124)	33.9%	0(0)	0.0%	23(2)	6.3%	156(21)	19.8%	587(32)	32.3%	980(53)	38.4%	395(15)	50.7%	16(0)	69.6%
	男性	1493(92)	39.2%	0(0)	0.0%	9(1)	4.4%	101(19)	23.7%	382(17)	36.2%	715(43)	44.9%	272(12)	55.1%	14(0)	77.8%
	女性	664(32)	26.0%	0(0)	0.0%	14(1)	8.7%	55(3)	15.3%	205(15)	26.9%	265(10)	27.6%	123(3)	43.2%	2(0)	40.0%
③びまん 性胸膜肥 厚	小計	63(8)	1.0%	0(0)	0.0%	1(0)	0.3%	9(0)	1.1%	10(1)	0.6%	24(6)	0.9%	17(1)	2.2%	2(0)	8.7%
	男性	49(7)	1.3%	0(0)	0.0%	1(0)	0.5%	2(0)	0.5%	10(1)	0.9%	19(5)	1.2%	16(1)	3.2%	1(0)	5.6%
	女性	14(1)	0.5%	0(0)	0.0%	0(0)	0.0%	7(0)	1.9%	0(0)	0.0%	5(1)	0.5%	1(0)	0.4%	1(0)	20.0%
④胸膜腫 瘍(中皮 腫)疑い	小計	8(2)	0.1%	0(0)	0.0%	0(0)	0.0%	2(1)	0.3%	0(0)	0.0%	4(1)	0.2%	2(0)	0.3%	0(0)	0.0%
	男性	6(2)	0.2%	0(0)	0.0%	0(0)	0.0%	2(1)	0.5%	0(0)	0.0%	3(1)	0.2%	1(0)	0.2%	0(0)	0.0%
	女性	2(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.1%	1(0)	0.4%	0(0)	0.0%
⑤肺野の 間質影	小計	420(25)	6.6%	0(0)	0.0%	2(0)	0.5%	7(2)	0.9%	73(8)	4.0%	234(11)	9.2%	100(4)	12.8%	4(0)	17.4%
	男性	331(19)	8.7%	0(0)	0.0%	1(0)	0.5%	5(1)	1.2%	55(6)	5.2%	185(8)	11.6%	82(4)	16.6%	3(0)	16.7%
	女性	89(6)	3.5%	0(0)	0.0%	1(0)	0.6%	2(1)	0.6%	18(2)	2.4%	49(3)	5.1%	18(0)	6.3%	1(0)	20.0%
⑥円形無 気肺	小計	27(3)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	15(1)	0.6%	12(2)	1.5%	0(0)	0.0%
	男性	23(3)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	12(1)	0.8%	11(2)	2.2%	0(0)	0.0%
	女性	4(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.3%	1(0)	0.4%	0(0)	0.0%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	99(25)	1.6%	1(0)	2.9%	5(1)	1.4%	11(3)	1.4%	26(6)	1.4%	46(12)	1.8%	10(3)	1.3%	0(0)	0.0%
	男性	66(17)	1.7%	1(0)	9.1%	2(0)	1.0%	5(1)	1.2%	18(6)	1.7%	34(8)	2.1%	6(2)	1.2%	0(0)	0.0%
	女性	33(8)	1.3%	0(0)	0.0%	3(1)	1.9%	6(2)	1.7%	8(0)	1.1%	12(4)	1.3%	4(1)	1.4%	0(0)	0.0%
⑧リンパ 節の腫大	小計	38(3)	0.6%	0(0)	0.0%	0(0)	0.0%	4(0)	0.5%	8(1)	0.4%	14(1)	0.5%	12(1)	1.5%	0(0)	0.0%
	男性	22(2)	0.6%	0(0)	0.0%	0(0)	0.0%	2(0)	0.5%	6(1)	0.6%	10(1)	0.6%	4(0)	0.8%	0(0)	0.0%
	女性	16(1)	0.6%	0(0)	0.0%	0(0)	0.0%	2(0)	0.6%	2(0)	0.3%	4(0)	0.4%	8(1)	2.8%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

表5-2 ばく露分類・性別検査所見((A)実施方法①、平成27~31年度延べ人数)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	7,926		1,994		762		1,006		746		3,418	
	男性	4,747		1,713		600		300		535		1,599	
	女性	3,179		281		162		706		211		1,819	
石綿関連 所見(疑い を含む)実 人数	合計	3,082(219)	38.9%	952(66)	47.7%	320(17)	42.0%	422(23)	41.9%	267(20)	35.8%	1,121(93)	32.8%
	男性	2,129(147)	44.8%	867(55)	50.6%	266(16)	44.3%	157(10)	52.3%	210(18)	39.3%	629(48)	39.3%
	女性	953(72)	30.0%	85(11)	30.2%	54(1)	33.3%	265(13)	37.5%	57(2)	27.0%	492(45)	27.0%
①胸水貯 留	小計	31(1)	0.4%	22(0)	1.1%	0(0)	0.0%	3(0)	0.3%	2(0)	0.3%	4(1)	0.1%
	男性	27(0)	0.6%	22(0)	1.3%	0(0)	0.0%	1(0)	0.3%	2(0)	0.4%	2(0)	0.1%
	女性	4(1)	0.1%	0(0)	0.0%	0(0)	0.0%	2(0)	0.3%	0(0)	0.0%	2(1)	0.1%
②胸膜プ ラーク(胸膜 肥厚斑)	小計	2,672(175)	33.7%	823(53)	41.3%	282(15)	37.0%	387(17)	38.5%	215(16)	28.8%	964(74)	28.2%
	男性	1,846(121)	38.9%	753(45)	44.0%	234(15)	39.0%	144(10)	48.0%	167(14)	31.2%	548(37)	34.3%
	女性	826(54)	26.0%	70(8)	24.9%	48(0)	29.6%	243(7)	34.4%	48(2)	22.7%	417(37)	22.9%
③びまん性 胸膜肥厚	小計	82(13)	1.0%	30(2)	1.5%	7(3)	0.9%	15(0)	1.5%	10(2)	1.3%	20(6)	0.6%
	男性	64(11)	1.3%	27(2)	1.6%	7(3)	1.2%	4(0)	1.3%	10(2)	1.9%	16(4)	1.0%
	女性	18(2)	0.6%	3(0)	1.1%	0(0)	0.0%	11(0)	1.6%	0(0)	0.0%	4(2)	0.2%
④胸膜腫 瘍(中皮腫) 疑い	小計	15(6)	0.2%	2(2)	0.1%	2(1)	0.3%	3(0)	0.3%	3(1)	0.4%	5(2)	0.1%
	男性	9(4)	0.2%	2(2)	0.1%	1(0)	0.2%	1(0)	0.3%	3(1)	0.6%	2(1)	0.1%
	女性	6(2)	0.2%	0(0)	0.0%	1(1)	0.6%	2(0)	0.3%	0(0)	0.0%	3(1)	0.2%
⑤肺野の 間質影	小計	523(33)	6.6%	198(15)	9.9%	55(1)	7.2%	50(3)	5.0%	55(4)	7.4%	165(10)	4.8%
	男性	409(27)	8.6%	184(14)	10.7%	51(1)	8.5%	19(1)	6.3%	48(3)	9.0%	107(8)	6.7%
	女性	114(6)	3.6%	14(1)	5.0%	4(0)	2.5%	31(2)	4.4%	7(1)	3.3%	58(2)	3.2%
⑥円形無 気肺	小計	34(4)	0.4%	15(2)	0.8%	2(1)	0.3%	0(0)	0.0%	1(0)	0.1%	16(1)	0.5%
	男性	30(4)	0.6%	15(2)	0.9%	2(1)	0.3%	0(0)	0.0%	1(0)	0.2%	12(1)	0.8%
	女性	4(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(0)	0.2%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	118(27)	1.5%	33(8)	1.7%	10(1)	1.3%	15(4)	1.5%	13(4)	1.7%	47(10)	1.4%
	男性	76(17)	1.6%	29(6)	1.7%	7(1)	1.2%	5(0)	1.7%	11(4)	2.1%	24(6)	1.5%
	女性	42(10)	1.3%	4(2)	1.4%	3(0)	1.9%	10(4)	1.4%	2(0)	0.9%	23(4)	1.3%
⑧リンパ節 の腫大	小計	73(3)	0.9%	25(1)	1.3%	8(1)	1.0%	8(1)	0.8%	11(0)	1.5%	21(0)	0.6%
	男性	43(2)	0.9%	19(1)	1.1%	8(1)	1.3%	2(0)	0.7%	4(0)	0.7%	10(0)	0.6%
	女性	30(1)	0.9%	6(0)	2.1%	0(0)	0.0%	6(1)	0.8%	7(0)	3.3%	11(0)	0.6%

※ ()内は疑い(内数)

※ ①~⑧で複数の所見が出た者は全て計上している。①~⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

表5-3 年齢階層・性別検査所見((B)実施方法②、平成29～31年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
検査 受診者	合計	1,010		16		104		145		262		427		53		3	
	男性	727		10		74		94		183		321		42		3	
	女性	283		6		30		51		79		106		11		0	
石綿関連 所見(疑い を含む)実 人数	合計	103(0)	10.2%	0(0)	0.0%	2(0)	1.9%	4(0)	2.8%	22(0)	8.4%	63(0)	14.8%	10(0)	18.9%	2(0)	66.7%
	男性	86(0)	11.8%	0(0)	0.0%	0(0)	0.0%	4(0)	4.3%	18(0)	9.8%	56(0)	17.4%	6(0)	14.3%	2(0)	66.7%
	女性	17(0)	6.0%	0(0)	0.0%	2(0)	6.7%	0(0)	0.0%	4(0)	5.1%	7(0)	6.6%	4(0)	36.4%	0(0)	－
①胸水貯 留	小計	2(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	46(0)	4.6%	0(0)	0.0%	1(0)	1.0%	1(0)	0.7%	6(0)	2.3%	32(0)	7.5%	4(0)	7.5%	2(0)	66.7%
	男性	36(0)	5.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.1%	4(0)	2.2%	27(0)	8.4%	2(0)	4.8%	2(0)	66.7%
	女性	10(0)	3.5%	0(0)	0.0%	1(0)	3.3%	0(0)	0.0%	2(0)	2.5%	5(0)	4.7%	2(0)	18.2%	0(0)	－
③びまん 性胸膜肥 厚	小計	29(0)	2.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	11(0)	4.2%	17(0)	4.0%	1(0)	1.9%	0(0)	0.0%
	男性	29(0)	4.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	11(0)	6.0%	17(0)	5.3%	1(0)	2.4%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
④胸膜腫 瘍(中皮 腫)疑い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑤肺野の 間質影	小計	28(0)	2.8%	0(0)	0.0%	0(0)	0.0%	2(0)	1.4%	4(0)	1.5%	19(0)	4.4%	3(0)	5.7%	0(0)	0.0%
	男性	25(0)	3.4%	0(0)	0.0%	0(0)	0.0%	2(0)	2.1%	3(0)	1.6%	18(0)	5.6%	2(0)	4.8%	0(0)	0.0%
	女性	3(0)	1.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.3%	1(0)	0.9%	1(0)	9.1%	0(0)	－
⑥円形無 気肺	小計	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.9%	0(0)	0.0%
	男性	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.4%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	10(0)	1.0%	0(0)	0.0%	1(0)	1.0%	1(0)	0.7%	3(0)	1.1%	3(0)	0.7%	2(0)	3.8%	0(0)	0.0%
	男性	6(0)	0.8%	0(0)	0.0%	0(0)	0.0%	1(0)	1.1%	2(0)	1.1%	2(0)	0.6%	1(0)	2.4%	0(0)	0.0%
	女性	4(0)	1.4%	0(0)	0.0%	1(0)	3.3%	0(0)	0.0%	1(0)	1.3%	1(0)	0.9%	1(0)	9.1%	0(0)	－
⑧リンパ 節の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

表5-4 ばく露分類・性別検査所見((B)実施方法②、平成 29～31 年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
検査 受診者	合計	1,010		371		241		58		173		167	
	男性	727		357		196		13		83		78	
	女性	283		14		45		45		90		89	
石綿関連 所見(疑い を含む)実 人数	合計	103(0)	10.2%	53(0)	14.3%	24(0)	10.0%	2(0)	3.4%	10(0)	5.8%	14(0)	8.4%
	男性	86(0)	11.8%	53(0)	14.8%	20(0)	10.2%	0(0)	0.0%	7(0)	8.4%	6(0)	7.7%
	女性	17(0)	6.0%	0(0)	0.0%	4(0)	8.9%	2(0)	4.4%	3(0)	3.3%	8(0)	9.0%
①胸水貯 留	小計	2(0)	0.2%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	0(0)	0.0%
	男性	2(0)	0.3%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	1(0)	1.2%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜プ ラーク(胸膜 肥厚斑)	小計	46(0)	4.6%	27(0)	7.3%	7(0)	2.9%	1(0)	1.7%	2(0)	1.2%	9(0)	5.4%
	男性	36(0)	5.0%	27(0)	7.6%	5(0)	2.6%	0(0)	0.0%	1(0)	1.2%	3(0)	3.8%
	女性	10(0)	3.5%	0(0)	0.0%	2(0)	4.4%	1(0)	2.2%	1(0)	1.1%	6(0)	6.7%
③びまん性 胸膜肥厚	小計	29(0)	2.9%	21(0)	5.7%	6(0)	2.5%	0(0)	0.0%	2(0)	1.2%	0(0)	0.0%
	男性	29(0)	4.0%	21(0)	5.9%	6(0)	3.1%	0(0)	0.0%	2(0)	2.4%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
④胸膜腫 瘍(中皮腫) 疑い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	28(0)	2.8%	11(0)	3.0%	8(0)	3.3%	1(0)	1.7%	4(0)	2.3%	4(0)	2.4%
	男性	25(0)	3.4%	11(0)	3.1%	7(0)	3.6%	0(0)	0.0%	3(0)	3.6%	4(0)	5.1%
	女性	3(0)	1.1%	0(0)	0.0%	1(0)	2.2%	1(0)	2.2%	1(0)	1.1%	0(0)	0.0%
⑥円形無 気肺	小計	1(0)	0.1%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	1(0)	0.1%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	10(0)	1.0%	3(0)	0.8%	4(0)	1.7%	0(0)	0.0%	1(0)	0.6%	2(0)	1.2%
	男性	6(0)	0.8%	3(0)	0.8%	3(0)	1.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	4(0)	1.4%	0(0)	0.0%	1(0)	2.2%	0(0)	0.0%	1(0)	1.1%	2(0)	2.2%
⑧リンパ節 の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-1 参考) 年齢階層・性別検査所見((A)実施方法①、平成27年度)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
CT検査 受診者	合計	1,566							
	男性	943							
	女性	623							
石綿関連 所見(疑い を含む)実 人数	合計	604(68) 38.6%							
	男性	414(41) 43.9%							
	女性	190(27) 30.5%							
①胸水貯 留	小計	9(0) 0.6%	0(0) -	0(0) -	1(0) -	0(0) -	3(0) -	3(0) -	2(0) -
	男性	8(0) 0.8%	0(0) -	0(0) -	1(0) -	0(0) -	3(0) -	3(0) -	1(0) -
	女性	1(0) 0.2%	0(0) -	0(0) -	0(0) -	0(0) -	0(0) -	0(0) -	1(0) -
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	515(51) 32.9%	0(0) -	8(0) -	53(5) -	165(17) -	220(20) -	67(9) -	2(0) -
	男性	353(29) 37.4%	0(0) -	4(0) -	30(2) -	110(9) -	159(13) -	48(5) -	2(0) -
	女性	162(22) 26.0%	0(0) -	4(0) -	23(3) -	55(8) -	61(7) -	19(4) -	0(0) -
③びまん 性胸膜肥 厚	小計	19(5) 1.2%	0(0) -	0(0) -	1(0) -	8(2) -	5(2) -	5(1) -	0(0) -
	男性	15(4) 1.6%	0(0) -	0(0) -	0(0) -	6(2) -	4(1) -	5(1) -	0(0) -
	女性	4(1) 0.6%	0(0) -	0(0) -	1(0) -	2(0) -	1(1) -	0(0) -	0(0) -
④胸膜腫 瘍(中皮 腫)疑い	小計	7(4) 0.4%	0(0) -	0(0) -	2(0) -	1(1) -	1(0) -	3(3) -	0(0) -
	男性	3(2) 0.3%	0(0) -	0(0) -	1(0) -	0(0) -	0(0) -	2(2) -	0(0) -
	女性	4(2) 0.6%	0(0) -	0(0) -	1(0) -	1(1) -	1(0) -	1(1) -	0(0) -
⑤肺野の 間質影	小計	103(8) 6.6%	0(0) -	1(0) -	7(1) -	25(1) -	50(6) -	20(0) -	0(0) -
	男性	78(8) 8.3%	0(0) -	0(0) -	6(1) -	17(1) -	36(6) -	19(0) -	0(0) -
	女性	25(0) 4.0%	0(0) -	1(0) -	1(0) -	8(0) -	14(0) -	1(0) -	0(0) -
⑥円形無 気肺	小計	7(1) 0.4%	0(0) -	0(0) -	0(0) -	0(0) -	3(1) -	4(0) -	0(0) -
	男性	7(1) 0.7%	0(0) -	0(0) -	0(0) -	0(0) -	3(1) -	4(0) -	0(0) -
	女性	0(0) 0.0%	0(0) -	0(0) -	0(0) -	0(0) -	0(0) -	0(0) -	0(0) -
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	19(2) 1.2%	0(0) -	0(0) -	1(0) -	4(0) -	12(1) -	1(1) -	1(0) -
	男性	10(0) 1.1%	0(0) -	0(0) -	0(0) -	3(0) -	7(0) -	0(0) -	0(0) -
	女性	9(2) 1.4%	0(0) -	0(0) -	1(0) -	1(0) -	5(1) -	1(1) -	1(0) -
⑧リンパ 節の腫大	小計	35(0) 2.2%	2(0) -	1(0) -	1(0) -	14(0) -	12(0) -	5(0) -	0(0) -
	男性	21(0) 2.2%	0(0) -	1(0) -	0(0) -	6(0) -	11(0) -	3(0) -	0(0) -
	女性	14(0) 2.2%	2(0) -	0(0) -	1(0) -	8(0) -	1(0) -	2(0) -	0(0) -

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 「CT検査受診者」及び「石綿関連所見(疑いを含む)実人数」の年齢階層別人数はデータがないため記載していない

※ 小数点以下第2位を四捨五入

(表5-1 参考) 年齢階層・性別検査所見((A)実施方法①、平成28年度)

(単位:人)

		合計	40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
CT検査 受診者	合計	1,481	11		92		177		471		570		159		1	
	男性	910	4		50		91		278		375		111		1	
	女性	571	7		42		86		193		195		48		0	
石綿関連 所見(疑い を含む)実 人数	合計	563(47) 38.0%	0(0)	0.0%	6(2)	6.5%	50(10)	28.2%	162(9)	34.4%	249(22)	43.7%	95(4)	59.7%	1(0)	100.0%
	男性	392(35) 43.1%	0(0)	0.0%	1(1)	2.0%	31(9)	34.1%	102(5)	36.7%	187(18)	49.9%	70(2)	63.1%	1(0)	100.0%
	女性	171(12) 29.9%	0(0)	0.0%	5(1)	11.9%	19(1)	22.1%	60(4)	31.1%	62(4)	31.8%	25(2)	52.1%	0(0)	-
①胸水貯 留	小計	4(0) 0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	1.9%	1(0)	100.0%
	男性	4(0) 0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	2.7%	1(0)	100.0%
	女性	0(0) 0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	493(42) 33.3%	0(0)	0.0%	4(2)	4.3%	46(8)	26.0%	151(10)	32.1%	207(18)	36.3%	84(4)	52.8%	1(0)	100.0%
	男性	334(29) 36.7%	0(0)	0.0%	1(1)	2.0%	27(7)	29.7%	93(6)	33.5%	152(13)	40.5%	60(2)	54.1%	1(0)	100.0%
	女性	159(13) 27.8%	0(0)	0.0%	3(1)	7.1%	19(1)	22.1%	58(4)	30.1%	55(5)	28.2%	24(2)	50.0%	0(0)	-
③びまん 性胸膜肥 厚	小計	19(4) 1.3%	0(0)	0.0%	0(0)	0.0%	3(0)	1.7%	4(0)	0.8%	7(3)	1.2%	5(1)	3.1%	0(0)	0.0%
	男性	16(3) 1.8%	0(0)	0.0%	0(0)	0.0%	1(0)	1.1%	4(0)	1.4%	6(2)	1.6%	5(1)	4.5%	0(0)	0.0%
	女性	3(1) 0.5%	0(0)	0.0%	0(0)	0.0%	2(0)	2.3%	0(0)	0.0%	1(1)	0.5%	0(0)	0.0%	0(0)	-
④胸膜腫 瘍(中皮 腫)疑い	小計	2(0) 0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%
	男性	1(0) 0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%
	女性	1(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	0(0)	0.0%	0(0)	-
⑤肺野の 間質影	小計	107(6) 7.2%	0(0)	0.0%	1(0)	1.1%	2(1)	1.1%	23(3)	4.9%	55(2)	9.6%	26(0)	16.4%	0(0)	0.0%
	男性	89(5) 9.8%	0(0)	0.0%	0(0)	0.0%	2(1)	2.2%	19(2)	6.8%	44(2)	11.7%	24(0)	21.6%	0(0)	0.0%
	女性	18(1) 3.2%	0(0)	0.0%	1(0)	2.4%	0(0)	0.0%	4(1)	2.1%	11(0)	5.6%	2(0)	4.2%	0(0)	-
⑥円形無 気肺	小計	7(1) 0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.4%	5(1)	3.1%	0(0)	0.0%
	男性	6(1) 0.7%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%	5(1)	4.5%	0(0)	0.0%
	女性	1(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	0(0)	0.0%	0(0)	-
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	22(7) 1.5%	0(0)	0.0%	1(0)	1.1%	2(1)	1.1%	2(1)	0.4%	14(5)	2.5%	3(0)	1.9%	0(0)	0.0%
	男性	18(7) 2.0%	0(0)	0.0%	0(0)	0.0%	2(1)	2.2%	2(1)	0.7%	13(5)	3.5%	1(0)	0.9%	0(0)	0.0%
	女性	4(0) 0.7%	0(0)	0.0%	1(0)	2.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	2(0)	4.2%	0(0)	-
⑧リンパ 節の腫大	小計	11(1) 0.7%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	2(1)	0.4%	6(0)	1.1%	2(0)	1.3%	0(0)	0.0%
	男性	8(1) 0.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(1)	0.7%	5(0)	1.3%	1(0)	0.9%	0(0)	0.0%
	女性	3(0) 0.5%	0(0)	0.0%	0(0)	0.0%	1(0)	1.2%	0(0)	0.0%	1(0)	0.5%	1(0)	2.1%	0(0)	-

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-1 参考) 年齢階層・性別検査所見((A)実施方法①、平成29年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
CT検査 受診者	合計	1,659		11		110		208		496		632		195		7	
	男性	993		2		65		109		285		406		123		3	
	女性	666		9		45		99		211		226		72		4	
石綿関連 所見(疑い を含む)実 人数	合計	618(46)	37.3%	0(0)	0.0%	7(0)	6.4%	52(8)	25.0%	176(15)	35.1%	272(18)	43.0%	110(5)	56.4%	3(0)	42.9%
	男性	433(35)	43.6%	0(0)	0.0%	4(0)	6.2%	33(6)	30.3%	123(9)	42.8%	201(15)	49.5%	72(5)	58.5%	1(0)	33.3%
	女性	185(11)	27.8%	0(0)	0.0%	3(0)	6.7%	19(2)	19.2%	53(6)	24.6%	71(3)	31.4%	38(0)	52.8%	2(0)	50.0%
①胸水貯 留	小計	6(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	2(0)	0.4%	3(0)	0.5%	0(0)	0.0%	0(0)	0.0%
	男性	5(0)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.7%	3(0)	0.7%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	535(38)	32.2%	0(0)	0.0%	7(0)	6.4%	46(6)	22.1%	148(11)	29.8%	231(17)	36.6%	100(3)	51.3%	3(0)	42.9%
	男性	374(29)	37.7%	0(0)	0.0%	4(0)	6.2%	30(5)	27.5%	104(6)	36.5%	170(15)	41.9%	65(3)	52.8%	1(0)	33.3%
	女性	161(9)	24.2%	0(0)	0.0%	3(0)	6.7%	16(2)	16.2%	44(5)	20.9%	61(2)	27.0%	35(0)	48.6%	2(0)	50.0%
③びまん 性胸膜肥 厚	小計	13(1)	0.8%	0(0)	0.0%	0(0)	0.0%	2(0)	1.0%	4(0)	0.8%	3(1)	0.5%	3(0)	1.5%	1(0)	14.3%
	男性	10(1)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(0)	1.4%	3(1)	0.7%	3(0)	2.4%	0(0)	0.0%
	女性	3(0)	0.5%	0(0)	0.0%	0(0)	0.0%	2(0)	2.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	25.0%
④胸膜腫 瘍(中皮 腫)疑い	小計	2(2)	0.1%	0(0)	0.0%	0(0)	0.0%	1(1)	0.5%	0(0)	0.0%	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%
	男性	2(2)	0.2%	0(0)	0.0%	0(0)	0.0%	1(1)	0.9%	0(0)	0.0%	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	110(9)	6.6%	0(0)	0.0%	0(0)	0.0%	2(0)	1.0%	21(4)	4.2%	62(3)	9.8%	25(2)	12.8%	0(0)	0.0%
	男性	87(7)	8.8%	0(0)	0.0%	0(0)	0.0%	1(0)	0.9%	16(3)	5.6%	51(2)	12.6%	19(2)	15.4%	0(0)	0.0%
	女性	23(2)	3.5%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	5(1)	2.4%	11(1)	4.9%	6(0)	8.3%	0(0)	0.0%
⑥円形無 気肺	小計	6(1)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(1)	0.6%	2(0)	1.0%	0(0)	0.0%
	男性	5(1)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(1)	0.7%	2(0)	1.6%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	25(6)	1.5%	0(0)	0.0%	0(0)	0.0%	4(1)	1.9%	10(2)	2.0%	10(2)	1.6%	1(1)	0.5%	0(0)	0.0%
	男性	17(5)	1.7%	0(0)	0.0%	0(0)	0.0%	1(0)	0.9%	7(2)	2.5%	8(2)	2.0%	1(1)	0.8%	0(0)	0.0%
	女性	8(1)	1.2%	0(0)	0.0%	0(0)	0.0%	3(1)	3.0%	3(0)	1.4%	2(0)	0.9%	0(0)	0.0%	0(0)	0.0%
⑧リンパ 節の腫大	小計	12(1)	0.7%	0(0)	0.0%	0(0)	0.0%	3(0)	1.4%	3(0)	0.6%	3(1)	0.5%	3(0)	1.5%	0(0)	0.0%
	男性	6(1)	0.6%	0(0)	0.0%	0(0)	0.0%	2(0)	1.8%	2(0)	0.7%	1(1)	0.2%	1(0)	0.8%	0(0)	0.0%
	女性	6(0)	0.9%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	1(0)	0.5%	2(0)	0.9%	2(0)	2.8%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-1 参考) 年齢階層・性別検査所見((A)実施方法①、平成30年度)

(単位:人)

		合計	40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
CT検査 受診者	合計	1,610	11		80		192		453		668		200		6	
	男性	946	4		45		103		265		401		123		5	
	女性	664	7		35		89		188		267		77		1	
石綿関連 所見(疑い を含む)実 人数	合計	652(37) 40.5%	1(0)	9.1%	8(1)	10.0%	33(6)	17.2%	169(11)	37.3%	322(15)	48.2%	113(4)	56.5%	6(0)	100.0%
	男性	457(24) 48.3%	1(0)	25.0%	4(0)	8.9%	21(5)	20.4%	116(6)	43.8%	232(11)	57.9%	78(2)	63.4%	5(0)	100.0%
	女性	195(13) 29.4%	0(0)	0.0%	4(1)	11.4%	12(1)	13.5%	53(5)	28.2%	90(4)	33.7%	35(2)	45.5%	1(0)	100.0%
①胸水貯 留	小計	6(0) 0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%	2(0)	0.3%	2(0)	1.0%	1(0)	16.7%
	男性	5(0) 0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	2(0)	0.5%	1(0)	0.8%	1(0)	20.0%
	女性	1(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.3%	0(0)	0.0%
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	566(31) 35.2%	0(0)	0.0%	5(0)	6.3%	31(5)	16.1%	149(9)	32.9%	274(13)	41.0%	102(4)	51.0%	5(0)	83.3%
	男性	399(23) 42.2%	0(0)	0.0%	2(0)	4.4%	20(5)	19.4%	101(4)	38.1%	200(11)	49.9%	71(3)	57.7%	5(0)	100.0%
	女性	167(8) 25.2%	0(0)	0.0%	3(0)	8.6%	11(0)	12.4%	48(5)	25.5%	74(2)	27.7%	31(1)	40.3%	0(0)	0.0%
③びまん 性胸膜肥 厚	小計	15(1) 0.9%	0(0)	0.0%	0(0)	0.0%	2(0)	1.0%	2(1)	0.4%	5(0)	0.7%	5(0)	2.5%	1(0)	16.7%
	男性	10(1) 1.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(1)	0.8%	3(0)	0.7%	4(0)	3.3%	1(0)	20.0%
	女性	5(0) 0.8%	0(0)	0.0%	0(0)	0.0%	2(0)	2.2%	0(0)	0.0%	2(0)	0.7%	1(0)	1.3%	0(0)	0.0%
④胸膜腫 瘍(中皮 腫)疑い	小計	2(0) 0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	1.0%	0(0)	0.0%
	男性	1(0) 0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%
	女性	1(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.3%	0(0)	0.0%
⑤肺野の 間質影	小計	101(3) 6.3%	0(0)	0.0%	0(0)	0.0%	2(1)	1.0%	21(1)	4.6%	56(1)	8.4%	21(0)	10.5%	1(0)	16.7%
	男性	74(2) 7.8%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	15(1)	5.7%	42(1)	10.5%	16(0)	13.0%	0(0)	0.0%
	女性	27(1) 4.1%	0(0)	0.0%	0(0)	0.0%	1(1)	1.1%	6(0)	3.2%	14(0)	5.2%	5(0)	6.5%	1(0)	100.0%
⑥円形無 気肺	小計	4(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(0)	0.6%	0(0)	0.0%	0(0)	0.0%
	男性	3(0) 0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.7%	0(0)	0.0%	0(0)	0.0%
	女性	1(0) 0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	24(4) 1.5%	1(0)	9.1%	3(1)	3.8%	0(0)	0.0%	6(1)	1.3%	12(1)	1.8%	2(1)	1.0%	0(0)	0.0%
	男性	17(2) 1.8%	1(0)	25.0%	2(0)	4.4%	0(0)	0.0%	5(1)	1.9%	7(0)	1.7%	2(1)	1.6%	0(0)	0.0%
	女性	7(2) 1.1%	0(0)	0.0%	1(1)	2.9%	0(0)	0.0%	1(0)	0.5%	5(1)	1.9%	0(0)	0.0%	0(0)	0.0%
⑧リンパ 節の腫大	小計	6(1) 0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%	2(0)	0.3%	3(1)	1.5%	0(0)	0.0%
	男性	4(0) 0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	2(0)	0.5%	1(0)	0.8%	0(0)	0.0%
	女性	2(1) 0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(1)	2.6%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-1 参考) 年齢階層・性別検査所見((A)実施方法①、平成31年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
CT検査 受診者	合計	1,610		2		85		210		396		683		225		9	
	男性	955		1		46		124		227		411		137		9	
	女性	655		1		39		86		169		272		88		0	
石綿関連 所見(疑い を含む)実 人数	合計	645(21)	40.1%	0(0)	0.0%	9(0)	10.6%	38(3)	18.1%	150(3)	37.9%	313(10)	45.8%	126(5)	56.0%	9(0)	100.0%
	男性	433(12)	45.3%	0(0)	0.0%	3(0)	6.5%	25(1)	20.2%	91(2)	40.1%	220(5)	53.5%	85(4)	62.0%	9(0)	100.0%
	女性	212(9)	32.4%	0(0)	0.0%	6(0)	15.4%	13(2)	15.1%	59(1)	34.9%	93(5)	34.2%	41(1)	46.6%	0(0)	-
①胸水貯 留	小計	6(1)	0.4%	0(0)	0.0%	0(0)	0.0%	1(1)	0.5%	0(0)	0.0%	1(0)	0.1%	3(0)	1.3%	1(0)	11.1%
	男性	5(0)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%	3(0)	2.2%	1(0)	11.1%
	女性	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%	1(1)	1.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	563(13)	35.0%	0(0)	0.0%	7(0)	8.2%	33(2)	15.7%	139(2)	35.1%	268(5)	39.2%	109(4)	48.4%	7(0)	77.8%
	男性	386(11)	40.4%	0(0)	0.0%	2(0)	4.3%	24(2)	19.4%	84(1)	37.0%	193(4)	47.0%	76(4)	55.5%	7(0)	77.8%
	女性	177(2)	27.0%	0(0)	0.0%	5(0)	12.8%	9(0)	10.5%	55(1)	32.5%	75(1)	27.6%	33(0)	37.5%	0(0)	-
③びまん 性胸膜肥 厚	小計	16(2)	1.0%	0(0)	0.0%	1(0)	1.2%	2(0)	1.0%	0(0)	0.0%	9(2)	1.3%	4(0)	1.8%	0(0)	0.0%
	男性	13(2)	1.4%	0(0)	0.0%	1(0)	2.2%	1(0)	0.8%	0(0)	0.0%	7(2)	1.7%	4(0)	2.9%	0(0)	0.0%
	女性	3(0)	0.5%	0(0)	0.0%	0(0)	0.0%	1(0)	1.2%	0(0)	0.0%	2(0)	0.7%	0(0)	0.0%	0(0)	-
④胸膜腫 瘍(中皮 腫)疑い	小計	2(0)	0.1%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	0(0)	0.0%	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
⑤肺野の 間質影	小計	102(7)	6.3%	0(0)	0.0%	1(0)	1.2%	1(0)	0.5%	8(0)	2.0%	61(5)	8.9%	28(2)	12.4%	3(0)	33.3%
	男性	81(5)	8.5%	0(0)	0.0%	1(0)	2.2%	1(0)	0.8%	5(0)	2.2%	48(3)	11.7%	23(2)	16.8%	3(0)	33.3%
	女性	21(2)	3.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	1.8%	13(2)	4.8%	5(0)	5.7%	0(0)	-
⑥円形無 気肺	小計	10(1)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	5(0)	0.7%	5(1)	2.2%	0(0)	0.0%
	男性	9(1)	0.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	5(0)	1.2%	4(1)	2.9%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.1%	0(0)	-
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	28(8)	1.7%	0(0)	0.0%	1(0)	1.2%	5(1)	2.4%	8(2)	2.0%	10(4)	1.5%	4(1)	1.8%	0(0)	0.0%
	男性	14(3)	1.5%	0(0)	0.0%	0(0)	0.0%	2(0)	1.6%	4(2)	1.8%	6(1)	1.5%	2(0)	1.5%	0(0)	0.0%
	女性	14(5)	2.1%	0(0)	0.0%	1(0)	2.6%	3(1)	3.5%	4(0)	2.4%	4(3)	1.5%	2(1)	2.3%	0(0)	-
⑧リンパ 節の腫大	小計	9(0)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.5%	3(0)	0.4%	4(0)	1.8%	0(0)	0.0%
	男性	4(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	2(0)	0.5%	1(0)	0.7%	0(0)	0.0%
	女性	5(0)	0.8%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	1(0)	0.4%	3(0)	3.4%	0(0)	-

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-2 参考) ばく露分類・性別検査所見((A)実施方法①、平成27年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	1,566		385		148		185		142		706	
	男性	943		333		117		62		102		329	
	女性	623		52		31		123		40		377	
石綿関連 所見(疑い を含む)実 人数	合計	604(68)	38.6%	180(22)	46.8%	60(6)	40.5%	80(6)	43.2%	50(4)	35.2%	234(30)	33.1%
	男性	414(41)	43.9%	165(18)	49.5%	51(5)	43.6%	33(2)	53.2%	39(3)	38.2%	126(13)	38.3%
	女性	190(27)	30.5%	15(4)	28.8%	9(1)	29.0%	47(4)	38.2%	11(1)	27.5%	108(17)	28.6%
①胸水貯 留	小計	9(0)	0.6%	7(0)	1.8%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.3%
	男性	8(0)	0.8%	7(0)	2.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
②胸膜プ ラーク(胸 膜肥厚斑)	小計	515(51)	32.9%	148(17)	38.4%	55(3)	37.2%	70(5)	37.8%	38(4)	26.8%	204(22)	28.9%
	男性	353(29)	37.4%	136(14)	40.8%	46(3)	39.3%	27(2)	43.5%	30(3)	29.4%	114(7)	34.7%
	女性	162(22)	26.0%	12(3)	23.1%	9(0)	29.0%	43(3)	35.0%	8(1)	20.0%	90(15)	23.9%
③びまん 性胸膜肥 厚	小計	19(5)	1.2%	7(0)	1.8%	2(2)	1.4%	2(0)	1.1%	2(0)	1.4%	6(3)	0.8%
	男性	15(4)	1.6%	6(0)	1.8%	2(2)	1.7%	1(0)	1.6%	2(0)	2.0%	4(2)	1.2%
	女性	4(1)	0.6%	1(0)	1.9%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%	2(1)	0.5%
④胸膜腫 瘍(中皮腫 疑い)	小計	7(4)	0.4%	0(0)	0.0%	1(1)	0.7%	0(0)	0.0%	1(1)	0.7%	5(2)	0.7%
	男性	3(2)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	1.0%	2(1)	0.6%
	女性	4(2)	0.6%	0(0)	0.0%	1(1)	3.2%	0(0)	0.0%	0(0)	0.0%	3(1)	0.8%
⑤肺野の 間質影	小計	103(8)	6.6%	34(4)	8.8%	7(0)	4.7%	11(0)	5.9%	11(1)	7.7%	40(3)	5.7%
	男性	78(8)	8.3%	33(4)	9.9%	7(0)	6.0%	6(0)	9.7%	9(1)	8.8%	23(3)	7.0%
	女性	25(0)	4.0%	1(0)	1.9%	0(0)	0.0%	5(0)	4.1%	2(0)	5.0%	17(0)	4.5%
⑥円形無 気肺	小計	7(1)	0.4%	4(0)	1.0%	1(1)	0.7%	0(0)	0.0%	0(0)	0.0%	2(0)	0.3%
	男性	7(1)	0.7%	4(0)	1.2%	1(1)	0.9%	0(0)	0.0%	0(0)	0.0%	2(0)	0.6%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	19(2)	1.2%	8(1)	2.1%	0(0)	0.0%	4(1)	2.2%	1(0)	0.7%	6(0)	0.8%
	男性	10(0)	1.1%	5(0)	1.5%	0(0)	0.0%	3(0)	4.8%	1(0)	1.0%	1(0)	0.3%
	女性	9(2)	1.4%	3(1)	5.8%	0(0)	0.0%	1(1)	0.8%	0(0)	0.0%	5(0)	1.3%
⑧リンパ節 の腫大	小計	35(0)	2.2%	10(0)	2.6%	1(0)	0.7%	4(0)	2.2%	7(0)	4.9%	13(0)	1.8%
	男性	21(0)	2.2%	9(0)	2.7%	1(0)	0.9%	1(0)	1.6%	2(0)	2.0%	8(0)	2.4%
	女性	14(0)	2.2%	1(0)	1.9%	0(0)	0.0%	3(0)	2.4%	5(0)	12.5%	5(0)	1.3%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-2 参考) ばく露分類・性別検査所見((A)実施方法①、平成28年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	1,481		400		146		179		129		627	
	男性	910		341		119		51		99		300	
	女性	571		59		27		128		30		327	
石綿関連 所見(疑い を含む)実 人数	合計	563(47)	38.0%	182(11)	45.4%	57(6)	39.0%	77(5)	43.0%	45(5)	34.9%	202(20)	32.2%
	男性	392(35)	43.1%	167(9)	48.8%	50(6)	42.0%	27(3)	52.9%	38(5)	38.4%	110(12)	36.7%
	女性	171(12)	29.9%	15(2)	25.4%	7(0)	25.9%	50(2)	39.1%	7(0)	23.3%	92(8)	28.1%
①胸水貯 留	小計	4(0)	0.3%	4(0)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	4(0)	0.4%	4(0)	1.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜プ ラーク(胸 膜肥厚斑)	小計	493(42)	33.3%	154(8)	38.4%	51(5)	34.9%	72(5)	40.2%	38(5)	29.5%	178(19)	28.4%
	男性	334(29)	36.7%	140(6)	40.9%	44(5)	37.0%	24(3)	47.1%	31(4)	31.3%	95(11)	31.7%
	女性	159(13)	27.8%	14(2)	23.7%	7(0)	25.9%	48(2)	37.5%	7(1)	23.3%	83(8)	25.4%
③びまん 性胸膜肥 厚	小計	19(4)	1.3%	6(1)	1.5%	4(1)	2.7%	3(0)	1.7%	1(0)	0.8%	5(2)	0.8%
	男性	16(3)	1.8%	6(1)	1.8%	4(1)	3.4%	1(0)	2.0%	1(0)	1.0%	4(1)	1.3%
	女性	3(1)	0.5%	0(0)	0.0%	0(0)	0.0%	2(0)	1.6%	0(0)	0.0%	1(1)	0.3%
④胸膜腫 瘍(中皮腫 疑い)	小計	2(0)	0.1%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	1(0)	0.8%	0(0)	0.0%
	男性	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	107(6)	7.2%	45(2)	11.2%	11(0)	7.5%	9(0)	5.0%	11(1)	8.5%	31(3)	4.9%
	男性	89(5)	9.8%	42(2)	12.3%	11(0)	9.2%	5(0)	9.8%	10(1)	10.1%	21(2)	7.0%
	女性	18(1)	3.2%	3(0)	5.1%	0(0)	0.0%	4(0)	3.1%	1(0)	3.3%	10(1)	3.1%
⑥円形無 気肺	小計	7(1)	0.5%	4(1)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.5%
	男性	6(1)	0.7%	4(1)	1.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.7%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	22(7)	1.5%	8(3)	2.0%	1(1)	0.7%	2(0)	1.1%	3(1)	2.3%	8(2)	1.3%
	男性	18(7)	2.0%	8(3)	2.3%	1(1)	0.8%	1(0)	2.0%	3(1)	3.0%	5(2)	1.7%
	女性	4(0)	0.7%	0(0)	0.0%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%	3(0)	0.9%
⑧リンパ節 の腫大	小計	11(1)	0.7%	4(0)	1.0%	3(1)	2.1%	0(0)	0.0%	0(0)	0.0%	4(0)	0.6%
	男性	8(1)	0.9%	3(0)	0.9%	3(1)	2.5%	0(0)	0.0%	0(0)	0.0%	2(0)	0.7%
	女性	3(0)	0.5%	1(0)	1.7%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.6%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-2 参考) ばく露分類・性別検査所見((A)実施方法①、平成29年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	1,659		406		143		245		156		709	
	男性	993		345		114		76		122		336	
	女性	666		61		29		169		34		373	
石綿関連 所見(疑い を含む)実 人数	合計	618(46)	37.3%	192(15)	47.3%	49(2)	34.3%	92(4)	37.6%	55(7)	35.3%	230(18)	32.4%
	男性	433(35)	43.6%	170(13)	49.3%	40(2)	35.1%	39(3)	51.3%	47(6)	38.5%	137(11)	40.8%
	女性	185(11)	27.8%	22(2)	36.1%	9(0)	31.0%	53(1)	31.4%	8(1)	23.5%	93(7)	24.9%
①胸水貯 留	小計	6(0)	0.4%	4(0)	1.0%	0(0)	0.0%	2(0)	0.8%	0(0)	0.0%	0(0)	0.0%
	男性	5(0)	0.5%	4(0)	1.2%	0(0)	0.0%	1(0)	1.3%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸膜 肥厚斑)	小計	535(38)	32.2%	168(13)	41.4%	42(2)	29.4%	84(4)	34.3%	42(4)	26.9%	198(15)	28.1%
	男性	374(29)	37.7%	150(12)	43.5%	35(2)	30.7%	36(3)	47.4%	36(4)	29.5%	117(8)	34.8%
	女性	161(9)	24.2%	18(1)	29.5%	7(0)	24.1%	48(1)	28.4%	6(0)	17.6%	82(7)	22.0%
③びまん 性胸膜肥 厚	小計	13(1)	0.8%	4(0)	1.0%	0(0)	0.0%	3(0)	1.2%	2(0)	1.3%	4(1)	0.6%
	男性	10(1)	1.0%	3(0)	0.9%	0(0)	0.0%	1(0)	1.3%	2(0)	1.6%	4(1)	1.2%
	女性	3(0)	0.5%	1(0)	1.6%	0(0)	0.0%	2(0)	1.2%	0(0)	0.0%	0(0)	0.0%
④胸膜腫 瘍(中皮腫 疑い)	小計	2(2)	0.1%	2(2)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	2(2)	0.2%	2(2)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	110(9)	6.6%	47(4)	11.6%	8(1)	5.6%	13(0)	5.3%	11(2)	7.1%	31(2)	4.4%
	男性	87(7)	8.8%	43(3)	12.5%	7(1)	6.1%	4(0)	5.3%	10(1)	8.2%	23(2)	6.8%
	女性	23(2)	3.5%	4(1)	6.6%	1(0)	3.4%	9(0)	5.3%	1(1)	2.9%	8(0)	2.1%
⑥円形無 気肺	小計	6(1)	0.4%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	5(1)	0.7%
	男性	5(1)	0.5%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(1)	1.2%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	25(6)	1.5%	4(0)	1.0%	2(0)	1.4%	3(1)	1.2%	3(2)	1.9%	13(3)	1.8%
	男性	17(5)	1.7%	4(0)	1.2%	0(0)	0.0%	1(0)	1.3%	3(2)	2.5%	9(3)	2.7%
	女性	8(1)	1.2%	0(0)	0.0%	2(0)	6.9%	2(1)	1.2%	0(0)	0.0%	4(0)	1.1%
⑧リンパ節 の腫大	小計	12(1)	0.7%	6(1)	1.5%	1(0)	0.7%	2(0)	0.8%	2(0)	1.3%	1(0)	0.1%
	男性	6(1)	0.6%	4(1)	1.2%	1(0)	0.9%	0(0)	0.0%	1(0)	0.8%	0(0)	0.0%
	女性	6(0)	0.9%	2(0)	3.3%	0(0)	0.0%	2(0)	1.2%	1(0)	2.9%	1(0)	0.3%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-2 参考) ばく露分類・性別検査所見((A)実施方法①、平成30年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	1,610		415		161		196		174		664	
	男性	946		355		124		50		112		305	
	女性	664		60		37		146		62		359	
石綿関連 所見(疑い を含む)実 人数	合計	652(37)	40.5%	204(10)	49.2%	79(3)	49.1%	90(6)	45.9%	65(3)	37.4%	214(15)	32.2%
	男性	457(24)	48.3%	189(9)	53.2%	65(3)	52.4%	31(1)	62.0%	47(3)	42.0%	125(8)	41.0%
	女性	195(13)	29.4%	15(1)	25.0%	14(0)	37.8%	59(5)	40.4%	18(0)	29.0%	89(7)	24.8%
①胸水貯 留	小計	6(0)	0.4%	4(0)	1.0%	0(0)	0.0%	1(0)	0.5%	1(0)	0.6%	0(0)	0.0%
	男性	5(0)	0.5%	4(0)	1.1%	0(0)	0.0%	0(0)	0.0%	1(0)	0.9%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	0.7%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸膜 肥厚斑)	小計	566(31)	35.2%	179(10)	43.1%	68(4)	42.2%	83(2)	42.3%	56(2)	32.2%	180(13)	27.1%
	男性	399(23)	42.2%	167(9)	47.0%	57(4)	46.0%	30(1)	60.0%	40(2)	35.7%	105(7)	34.4%
	女性	167(8)	25.2%	12(1)	20.0%	11(0)	29.7%	53(1)	36.3%	16(0)	25.8%	75(6)	20.9%
③びまん 性胸膜肥 厚	小計	15(1)	0.9%	7(0)	1.7%	0(0)	0.0%	4(0)	2.0%	2(1)	1.1%	2(0)	0.3%
	男性	10(1)	1.1%	6(0)	1.7%	0(0)	0.0%	0(0)	0.0%	2(1)	1.8%	2(0)	0.7%
	女性	5(0)	0.8%	1(0)	1.7%	0(0)	0.0%	4(0)	2.7%	0(0)	0.0%	0(0)	0.0%
④胸膜腫 瘍(中皮腫 疑い)	小計	2(0)	0.1%	0(0)	0.0%	0(0)	0.0%	1(0)	0.5%	1(0)	0.6%	0(0)	0.0%
	男性	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.9%	0(0)	0.0%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	1(0)	0.7%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	101(3)	6.3%	37(1)	8.9%	13(0)	8.1%	10(1)	5.1%	11(0)	6.3%	30(1)	4.5%
	男性	74(2)	7.8%	34(1)	9.6%	11(0)	8.9%	2(0)	4.0%	9(0)	8.0%	18(1)	5.9%
	女性	27(1)	4.1%	3(0)	5.0%	2(0)	5.4%	8(1)	5.5%	2(0)	3.2%	12(0)	3.3%
⑥円形無 気肺	小計	4(0)	0.2%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.5%
	男性	3(0)	0.3%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.7%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	24(4)	1.5%	4(0)	1.0%	3(0)	1.9%	1(1)	0.5%	6(1)	3.4%	10(2)	1.5%
	男性	17(2)	1.8%	4(0)	1.1%	2(0)	1.6%	0(0)	0.0%	4(1)	3.6%	7(1)	2.3%
	女性	7(2)	1.1%	0(0)	0.0%	1(0)	2.7%	1(1)	0.7%	2(0)	3.2%	3(1)	0.8%
⑧リンパ節 の腫大	小計	6(1)	0.4%	3(0)	0.7%	2(0)	1.2%	1(1)	0.5%	0(0)	0.0%	0(0)	0.0%
	男性	4(0)	0.4%	2(0)	0.6%	2(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	2(1)	0.3%	1(0)	1.7%	0(0)	0.0%	1(1)	0.7%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-2 参考) ばく露分類・性別検査所見((A)実施方法①、平成31年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
CT検査 受診者	合計	1,610		388		164		201		145		712	
	男性	955		339		126		61		100		329	
	女性	655		49		38		140		45		383	
石綿関連 所見(疑い を含む)実 人数	合計	645(21)	40.1%	194(8)	50.0%	75(0)	45.7%	83(2)	41.3%	52(1)	35.9%	241(10)	33.8%
	男性	433(12)	45.3%	176(6)	51.9%	60(0)	47.6%	27(1)	44.3%	39(1)	39.0%	131(4)	39.8%
	女性	212(9)	32.4%	18(2)	36.7%	15(0)	39.5%	56(1)	40.0%	13(0)	28.9%	110(6)	28.7%
①胸水貯 留	小計	6(1)	0.4%	3(0)	0.8%	0(0)	0.0%	0(0)	0.0%	1(0)	0.7%	2(1)	0.3%
	男性	5(0)	0.5%	3(0)	0.9%	0(0)	0.0%	0(0)	0.0%	1(0)	1.0%	1(0)	0.3%
	女性	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.3%
②胸膜ブ ラーク(胸膜 肥厚斑)	小計	563(13)	35.0%	174(5)	44.8%	66(1)	40.2%	78(1)	38.8%	41(1)	28.3%	204(5)	28.7%
	男性	386(11)	40.4%	160(4)	47.2%	52(1)	41.3%	27(1)	44.3%	30(1)	30.0%	117(4)	35.6%
	女性	177(2)	27.0%	14(1)	28.6%	14(0)	36.8%	51(0)	36.4%	11(0)	24.4%	87(1)	22.7%
③びまん性 胸膜肥厚	小計	16(2)	1.0%	6(1)	1.5%	1(0)	0.6%	3(0)	1.5%	3(1)	2.1%	3(0)	0.4%
	男性	13(2)	1.4%	6(1)	1.8%	1(0)	0.8%	1(0)	1.6%	3(1)	3.0%	2(0)	0.6%
	女性	3(0)	0.5%	0(0)	0.0%	0(0)	0.0%	2(0)	1.4%	0(0)	0.0%	1(0)	0.3%
④胸膜腫 瘍(中皮腫) 疑い	小計	2(0)	0.1%	0(0)	0.0%	1(0)	0.6%	1(0)	0.5%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.2%	0(0)	0.0%	1(0)	0.8%	1(0)	1.6%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の 間質影	小計	102(7)	6.3%	35(4)	9.0%	16(0)	9.8%	7(2)	3.5%	11(0)	7.6%	33(1)	4.6%
	男性	81(5)	8.5%	32(4)	9.4%	15(0)	11.9%	2(1)	3.3%	10(0)	10.0%	22(0)	6.7%
	女性	21(2)	3.2%	3(0)	6.1%	1(0)	2.6%	5(1)	3.6%	1(0)	2.2%	11(1)	2.9%
⑥円形無 気肺	小計	10(1)	0.6%	5(1)	1.3%	1(0)	0.6%	0(0)	0.0%	1(0)	0.7%	3(0)	0.4%
	男性	9(1)	0.9%	5(1)	1.5%	1(0)	0.8%	0(0)	0.0%	1(0)	1.0%	2(0)	0.6%
	女性	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	28(8)	1.7%	9(4)	2.3%	4(0)	2.4%	5(1)	2.5%	0(0)	0.0%	10(3)	1.4%
	男性	14(3)	1.5%	8(3)	2.4%	4(0)	3.2%	0(0)	0.0%	0(0)	0.0%	2(0)	0.6%
	女性	14(5)	2.1%	1(1)	2.0%	0(0)	0.0%	5(1)	3.6%	0(0)	0.0%	8(3)	2.1%
⑧リンパ節 の腫大	小計	9(0)	0.6%	2(0)	0.5%	1(0)	0.6%	1(0)	0.5%	2(0)	1.4%	3(0)	0.4%
	男性	4(0)	0.4%	1(0)	0.3%	1(0)	0.8%	1(0)	1.6%	1(0)	1.0%	0(0)	0.0%
	女性	5(0)	0.8%	1(0)	2.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.2%	3(0)	0.8%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別のCT検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-3 参考) 年齢階層・性別検査所見((B)実施方法②、平成29年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
検査 受診者	合計	198		9		26		35		43		79		6		0	
	男性	129		4		17		24		31		49		4		0	
	女性	69		5		9		11		12		30		2		0	
石綿関連 所見(疑い を含む)実 人数	合計	13(0)	6.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	5(0)	11.6%	8(0)	10.1%	0(0)	0.0%	0(0)	－
	男性	11(0)	8.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(0)	12.9%	7(0)	14.3%	0(0)	0.0%	0(0)	－
	女性	2(0)	2.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	8.3%	1(0)	3.3%	0(0)	0.0%	0(0)	－
①胸水貯 留	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	3(0)	1.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.3%	2(0)	2.5%	0(0)	0.0%	0(0)	－
	男性	3(0)	2.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	3.2%	2(0)	4.1%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
③びまん 性胸膜肥 厚	小計	6(0)	3.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	4.7%	4(0)	5.1%	0(0)	0.0%	0(0)	－
	男性	6(0)	4.7%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	6.5%	4(0)	8.2%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
④胸膜腫 瘍(中皮 腫)疑い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑤肺野の 間質影	小計	3(0)	1.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	3.8%	0(0)	0.0%	0(0)	－
	男性	2(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	4.1%	0(0)	0.0%	0(0)	－
	女性	1(0)	1.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	3.3%	0(0)	0.0%	0(0)	－
⑥円形無 気肺	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	3(0)	1.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	7.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	男性	2(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	6.5%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	女性	1(0)	1.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	8.3%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑧リンパ 節の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-3 参考) 年齢階層・性別検査所見((B)実施方法②、平成30年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
検査 受診者	合計	651		4		67		92		174		275		36		3	
	男性	491		3		52		60		124		218		31		3	
	女性	160		1		15		32		50		57		5		0	
石綿関連 所見(疑い を含む)実 人数	合計	69(0)	10.6%	0(0)	0.0%	2(0)	3.0%	3(0)	3.3%	15(0)	8.6%	42(0)	15.3%	5(0)	13.9%	2(0)	66.7%
	男性	59(0)	12.0%	0(0)	0.0%	0(0)	0.0%	3(0)	5.0%	13(0)	10.5%	38(0)	17.4%	3(0)	9.7%	2(0)	66.7%
	女性	10(0)	6.3%	0(0)	0.0%	2(0)	13.3%	0(0)	0.0%	2(0)	4.0%	4(0)	7.0%	2(0)	40.0%	0(0)	－
①胸水貯 留	小計	2(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.8%	1(0)	0.5%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	30(0)	4.6%	0(0)	0.0%	1(0)	1.5%	1(0)	1.1%	4(0)	2.3%	20(0)	7.3%	2(0)	5.6%	2(0)	66.7%
	男性	24(0)	4.9%	0(0)	0.0%	0(0)	0.0%	1(0)	1.7%	3(0)	2.4%	17(0)	7.8%	1(0)	3.2%	2(0)	66.7%
	女性	6(0)	3.8%	0(0)	0.0%	1(0)	6.7%	0(0)	0.0%	1(0)	2.0%	3(0)	5.3%	1(0)	20.0%	0(0)	－
③びまん 性胸膜肥 厚	小計	21(0)	3.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	8(0)	4.6%	12(0)	4.4%	1(0)	2.8%	0(0)	0.0%
	男性	21(0)	4.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	8(0)	6.5%	12(0)	5.5%	1(0)	3.2%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
④胸膜腫 瘍(中皮 腫)疑い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑤肺野の 間質影	小計	17(0)	2.6%	0(0)	0.0%	0(0)	0.0%	1(0)	1.1%	4(0)	2.3%	10(0)	3.6%	2(0)	5.6%	0(0)	0.0%
	男性	15(0)	3.1%	0(0)	0.0%	0(0)	0.0%	1(0)	1.7%	3(0)	2.4%	10(0)	4.6%	1(0)	3.2%	0(0)	0.0%
	女性	2(0)	1.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.0%	0(0)	0.0%	1(0)	20.0%	0(0)	－
⑥円形無 気肺	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	6(0)	0.9%	0(0)	0.0%	1(0)	1.5%	1(0)	1.1%	0(0)	0.0%	3(0)	1.1%	1(0)	2.8%	0(0)	0.0%
	男性	4(0)	0.8%	0(0)	0.0%	0(0)	0.0%	1(0)	1.7%	0(0)	0.0%	2(0)	0.9%	1(0)	3.2%	0(0)	0.0%
	女性	2(0)	1.3%	0(0)	0.0%	1(0)	6.7%	0(0)	0.0%	0(0)	0.0%	1(0)	1.8%	0(0)	0.0%	0(0)	－
⑧リンパ 節の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	－

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-3 参考) 年齢階層・性別検査所見((B)実施方法②、平成31年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
検査 受診者	合計	161		3		11		18		45		73		11		0	
	男性	107		3		5		10		28		54		7		0	
	女性	54		0		6		8		17		19		4		0	
石綿関連 所見(疑い を含む)実 人数	合計	21(0)	13.0%	0(0)	0.0%	0(0)	0.0%	1(0)	5.6%	2(0)	4.4%	13(0)	17.8%	5(0)	45.5%	0(0)	-
	男性	16(0)	15.0%	0(0)	0.0%	0(0)	0.0%	1(0)	10.0%	1(0)	3.6%	11(0)	20.4%	3(0)	42.9%	0(0)	-
	女性	5(0)	9.3%	0(0)	-	0(0)	0.0%	0(0)	0.0%	1(0)	5.9%	2(0)	10.5%	2(0)	50.0%	0(0)	-
①胸水貯 留	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
②胸膜ブ ラーク(胸 膜肥厚斑)	小計	13(0)	8.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.2%	10(0)	13.7%	2(0)	18.2%	0(0)	-
	男性	9(0)	8.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	8(0)	14.8%	1(0)	14.3%	0(0)	-
	女性	4(0)	7.4%	0(0)	-	0(0)	0.0%	0(0)	0.0%	1(0)	5.9%	2(0)	10.5%	1(0)	25.0%	0(0)	-
③びまん 性胸膜肥 厚	小計	2(0)	1.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.2%	1(0)	1.4%	0(0)	0.0%	0(0)	-
	男性	2(0)	1.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	3.6%	1(0)	1.9%	0(0)	0.0%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
④胸膜腫 瘍(中皮 腫)疑い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
⑤肺野の 間質影	小計	8(0)	5.0%	0(0)	0.0%	0(0)	0.0%	1(0)	5.6%	0(0)	0.0%	6(0)	8.2%	1(0)	9.1%	0(0)	-
	男性	8(0)	7.5%	0(0)	0.0%	0(0)	0.0%	1(0)	10.0%	0(0)	0.0%	6(0)	11.1%	1(0)	14.3%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
⑥円形無 気肺	小計	1(0)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	9.1%	0(0)	-
	男性	1(0)	0.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	14.3%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
⑦肺野の 腫瘤状陰 影(肺がん 等)	小計	1(0)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	9.1%	0(0)	-
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	女性	1(0)	1.9%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	25.0%	0(0)	-
⑧リンパ 節の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-
	女性	0(0)	0.0%	0(0)	-	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	-

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各年齢階層別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-4 参考)ばく露分類・性別検査所見((B)実施方法②、平成29年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
検査 受診者	合計	198		62		40		12		34		50	
	男性	129		57		27		3		19		23	
	女性	69		5		13		9		15		27	
石綿関連所 見(疑いを 含む)実人 数	合計	13(0)	6.6%	8(0)	12.9%	3(0)	7.5%	0(0)	0.0%	1(0)	2.9%	1(0)	2.0%
	男性	11(0)	8.5%	8(0)	14.0%	3(0)	11.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	2(0)	2.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	6.7%	1(0)	3.7%
①胸水貯留	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸膜 肥厚斑)	小計	3(0)	1.5%	3(0)	4.8%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	3(0)	2.3%	3(0)	5.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
③びまん性 胸膜肥厚	小計	6(0)	3.0%	5(0)	8.1%	1(0)	2.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	6(0)	4.7%	5(0)	8.8%	1(0)	3.7%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
④胸膜腫瘍 (中皮腫)疑 い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の間 質影	小計	3(0)	1.5%	0(0)	0.0%	2(0)	5.0%	0(0)	0.0%	1(0)	2.9%	0(0)	0.0%
	男性	2(0)	1.6%	0(0)	0.0%	2(0)	7.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	1.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	6.7%	0(0)	0.0%
⑥円形無気 肺	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑦肺野の腫 瘤状陰影 (肺がん等)	小計	3(0)	1.5%	2(0)	3.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	2.0%
	男性	2(0)	1.6%	2(0)	3.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	1.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	3.7%
⑧リンパ節 の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-4 参考)ばく露分類・性別検査所見((B)実施方法②、平成30年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
検査 受診者	合計	651		255		161		38		112		85	
	男性	491		252		137		9		52		41	
	女性	160		3		24		29		60		44	
石綿関連所 見(疑いを 含む)実人 数	合計	69(0)	10.6%	36(0)	14.1%	17(0)	10.6%	2(0)	5.3%	7(0)	6.3%	7(0)	8.2%
	男性	59(0)	12.0%	36(0)	14.3%	13(0)	9.5%	0(0)	0.0%	6(0)	11.5%	4(0)	9.8%
	女性	10(0)	6.3%	0(0)	0.0%	4(0)	16.7%	2(0)	6.9%	1(0)	1.7%	3(0)	6.8%
①胸水貯留	小計	2(0)	0.3%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.9%	0(0)	0.0%
	男性	2(0)	0.4%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	1.9%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸膜肥 厚斑)	小計	30(0)	4.6%	18(0)	7.1%	6(0)	3.7%	1(0)	2.6%	1(0)	0.9%	4(0)	4.7%
	男性	24(0)	4.9%	18(0)	7.1%	4(0)	2.9%	0(0)	0.0%	1(0)	1.9%	1(0)	2.4%
	女性	6(0)	3.8%	0(0)	0.0%	2(0)	8.3%	1(0)	3.4%	0(0)	0.0%	3(0)	6.8%
③びまん性 胸膜肥厚	小計	21(0)	3.2%	15(0)	5.9%	4(0)	2.5%	0(0)	0.0%	2(0)	1.8%	0(0)	0.0%
	男性	21(0)	4.3%	15(0)	6.0%	4(0)	2.9%	0(0)	0.0%	2(0)	3.8%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
④胸膜腫瘍 (中皮腫)疑 い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の間 質影	小計	17(0)	2.6%	7(0)	2.7%	4(0)	2.5%	1(0)	2.6%	2(0)	1.8%	3(0)	3.5%
	男性	15(0)	3.1%	7(0)	2.8%	3(0)	2.2%	0(0)	0.0%	2(0)	3.8%	3(0)	7.3%
	女性	2(0)	1.3%	0(0)	0.0%	1(0)	4.2%	1(0)	3.4%	0(0)	0.0%	0(0)	0.0%
⑥円形無気 肺	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑦肺野の腫 瘤状陰影 (肺がん等)	小計	6(0)	0.9%	1(0)	0.4%	4(0)	2.5%	0(0)	0.0%	1(0)	0.9%	0(0)	0.0%
	男性	4(0)	0.8%	1(0)	0.4%	3(0)	2.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	2(0)	1.3%	0(0)	0.0%	1(0)	4.2%	0(0)	0.0%	1(0)	1.7%	0(0)	0.0%
⑧リンパ節 の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表5-4 参考)ばく露分類・性別検査所見((B)実施方法②、平成31年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
検査 受診者	合計	161		54		40		8		27		32	
	男性	107		48		32		1		12		14	
	女性	54		6		8		7		15		18	
石綿関連所 見(疑いを 含む)実人 数	合計	21(0)	13.0%	9(0)	16.7%	4(0)	10.0%	0(0)	0.0%	2(0)	7.4%	6(0)	18.8%
	男性	16(0)	15.0%	9(0)	18.8%	4(0)	12.5%	0(0)	0.0%	1(0)	8.3%	2(0)	14.3%
	女性	5(0)	9.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	6.7%	4(0)	22.2%
①胸水貯留	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
②胸膜ブ ラーク(胸膜肥 厚斑)	小計	13(0)	8.1%	6(0)	11.1%	1(0)	2.5%	0(0)	0.0%	1(0)	3.7%	5(0)	15.6%
	男性	9(0)	8.4%	6(0)	12.5%	1(0)	3.1%	0(0)	0.0%	0(0)	0.0%	2(0)	14.3%
	女性	4(0)	7.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	6.7%	3(0)	16.7%
③びまん性 胸膜肥厚	小計	2(0)	1.2%	1(0)	1.9%	1(0)	2.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	1.9%	1(0)	2.1%	1(0)	3.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
④胸膜腫瘍 (中皮腫)疑 い	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑤肺野の間 質影	小計	8(0)	5.0%	4(0)	7.4%	2(0)	5.0%	0(0)	0.0%	1(0)	3.7%	1(0)	3.1%
	男性	8(0)	7.5%	4(0)	8.3%	2(0)	6.3%	0(0)	0.0%	1(0)	8.3%	1(0)	7.1%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑥円形無気 肺	小計	1(0)	0.6%	1(0)	1.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	1(0)	0.9%	1(0)	2.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
⑦肺野の腫 瘤状陰影 (肺がん等)	小計	1(0)	0.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	3.1%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	1.9%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	5.6%
⑧リンパ節 の腫大	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ ①～⑧で複数の所見が出た者は全て計上している。①～⑧の合計は実人数とは一致しない

※ 右欄は各ばく露別の検査受診者に対する有所見者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(3) 石綿関連疾患と診断された者の数

石綿関連疾患と診断された者（石綿によるものかどうかは特定されておらず、疑いも含む）の年齢階層・性別の関係及びばく露・性別の関係は表6に示すとおりである。

表6-1 石綿関連疾患と診断された者の年齢階層・性別分類表(平成27～31年度延べ人数)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
参加者	合計	11,012	77	701	1,370	3,287	4,339	1,194	44
	男性	6,590	36	421	762	1,893	2,697	750	31
	女性	4,422	41	280	608	1,394	1,642	444	13
中皮腫	小計	3(2) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(1) 0.1%	1(1) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	2(1) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(1) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	1(1) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(1) 0.1%	0(0) 0.0%	0(0) 0.0%
肺がん	小計	22(7) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(2) 0.2%	14(5) 0.3%	2(0) 0.2%	1(0) 2.3%
	男性	16(5) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	4(2) 0.2%	10(3) 0.4%	2(0) 0.3%	0(0) 0.0%
	女性	6(2) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	4(2) 0.2%	0(0) 0.0%	1(0) 7.7%
石綿肺	小計	14(2) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(0) 0.1%	9(2) 0.2%	1(0) 0.1%	1(0) 2.3%
	男性	13(2) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(0) 0.2%	9(2) 0.3%	0(0) 0.0%	1(0) 3.2%
	女性	1(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	0(0) 0.0%
びまん性 胸膜肥厚	小計	23(2) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(0) 0.2%	15(2) 0.3%	3(0) 0.3%	0(0) 0.0%
	男性	21(2) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(0) 0.3%	13(2) 0.5%	3(0) 0.4%	0(0) 0.0%
	女性	2(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(0) 0.1%	0(0) 0.0%	0(0) 0.0%
良性 石綿胸水	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
合計	小計	62(13) 0.6%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	15(3) 0.5%	39(10) 0.9%	6(0) 0.5%	2(0) 4.5%
	男性	52(10) 0.8%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	14(3) 0.7%	32(7) 1.2%	5(0) 0.7%	1(0) 3.2%
	女性	10(3) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	7(3) 0.4%	1(0) 0.2%	1(0) 7.7%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

表6-2 石綿関連疾患と診断された者のばく露・性別分類表(平成 27～31 年度延べ人数)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	10,946		2,797		1,246		1,288		1,075		4,540	
	男性	6,552		2,457		965		357		720		2,053	
	女性	4,394		340		281		931		355		2,487	
中皮腫	小計	3(2)	0.0%	1(0)	0.0%	0(0)	0.0%	1(1)	0.1%	0(0)	0.0%	1(1)	0.0%
	男性	2(1)	0.0%	1(0)	0.0%	0(0)	0.0%	1(1)	0.3%	0(0)	0.0%	0(0)	0.0%
	女性	1(1)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.0%
肺がん	小計	22(7)	0.2%	12(3)	0.4%	3(1)	0.2%	0(0)	0.0%	1(1)	0.1%	6(2)	0.1%
	男性	16(5)	0.2%	10(2)	0.4%	1(0)	0.1%	0(0)	0.0%	1(1)	0.1%	4(2)	0.2%
	女性	6(2)	0.1%	2(1)	0.6%	2(1)	0.7%	0(0)	0.0%	0(0)	0.0%	2(0)	0.1%
石綿肺	小計	14(2)	0.1%	6(1)	0.2%	2(0)	0.2%	1(0)	0.1%	2(1)	0.2%	3(0)	0.1%
	男性	13(2)	0.2%	5(1)	0.2%	2(0)	0.2%	1(0)	0.3%	2(1)	0.3%	3(0)	0.1%
	女性	1(0)	0.0%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	23(2)	0.2%	12(1)	0.4%	3(0)	0.2%	1(0)	0.1%	1(0)	0.1%	6(1)	0.1%
	男性	21(2)	0.3%	12(1)	0.5%	2(0)	0.2%	1(0)	0.3%	1(0)	0.1%	5(1)	0.2%
	女性	2(0)	0.0%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	62(13)	0.6%	31(5)	1.1%	8(1)	0.6%	3(1)	0.2%	4(2)	0.4%	16(4)	0.4%
	男性	52(10)	0.8%	28(4)	1.1%	5(0)	0.5%	3(1)	0.8%	4(2)	0.6%	12(3)	0.6%
	女性	10(3)	0.2%	3(1)	0.9%	3(1)	1.1%	0(0)	0.0%	0(0)	0.0%	4(1)	0.2%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 平成 27 年度の参加者の内「石綿ばく露の聴取のみを行った者 66 人」については、ばく露分類が不明のため、参加者数から除外している

※ 小数点以下第2位を四捨五入

(表6-1 参考) 石綿関連疾患と診断された者の年齢階層・性別分類表(平成27年度)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
参加者	合計	2,046	15	141	249	703	748	187	3
	男性	1,180	6	79	130	386	454	123	2
	女性	866	9	62	119	317	294	64	1
中皮腫	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
肺がん	小計	8(0) 0.4%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(0) 0.3%	4(0) 0.5%	1(0) 0.5%	1(0) 33.3%
	男性	5(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	3(0) 0.7%	1(0) 0.8%	0(0) 0.0%
	女性	3(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	1(0) 0.3%	0(0) 0.0%	1(0) 100%
石綿肺	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
びまん性 胸膜肥厚	小計	2(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	1(0) 0.5%	0(0) 0.0%
	男性	2(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	1(0) 0.8%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
良性 石綿胸水	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
合計	小計	10(0) 0.5%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(0) 0.3%	5(0) 0.7%	2(0) 1.1%	1(0) 33.3%
	男性	7(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	4(0) 0.9%	2(0) 1.6%	0(0) 0.0%
	女性	3(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	1(0) 0.3%	0(0) 0.0%	1(0) 100%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-1 参考) 石綿関連疾患と診断された者の年齢階層・性別分類表(平成28年度)

(単位:人)

		合計		40歳未満		40～49歳		50～59歳		60～69歳		70～79歳		80～89歳		90歳以上	
参加者	合計	1,902		12		125		230		611		716		203		5	
	男性	1,171		4		72		127		361		468		136		3	
	女性	731		8		53		103		250		248		67		2	
中皮腫	小計	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
肺がん	小計	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%
	男性	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.2%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
石綿肺	小計	3(1)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%	2(1)	0.3%	0(0)	0.0%	0(0)	0.0%
	男性	3(1)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%	2(1)	0.4%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	3(1)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%	1(1)	0.1%	1(0)	0.5%	0(0)	0.0%
	男性	3(1)	0.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%	1(1)	0.2%	1(0)	0.7%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	8(4)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(1)	0.5%	4(3)	0.6%	1(0)	0.5%	0(0)	0.0%
	男性	8(4)	0.7%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(1)	0.8%	4(3)	0.9%	1(0)	0.7%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-1 参考) 石綿関連疾患と診断された者の年齢階層・性別分類表(平成29年度)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
参加者	合計	2,260	27	156	291	672	860	241	13
	男性	1,332	12	95	164	381	527	146	7
	女性	928	15	61	127	291	333	95	6
中皮腫	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
肺がん	小計	3(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	2(0) 0.2%	0(0) 0.0%	0(0) 0.0%
	男性	2(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	1(0) 0.2%	0(0) 0.0%	0(0) 0.0%
	女性	1(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	0(0) 0.0%	0(0) 0.0%
石綿肺	小計	4(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	2(0) 0.2%	1(0) 0.4%	0(0) 0.0%
	男性	3(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	2(0) 0.4%	0(0) 0.0%	0(0) 0.0%
	女性	1(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 1.1%	0(0) 0.0%
びまん性 胸膜肥厚	小計	5(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(0) 0.4%	2(0) 0.2%	0(0) 0.0%	0(0) 0.0%
	男性	4(0) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(0) 0.8%	1(0) 0.2%	0(0) 0.0%	0(0) 0.0%
	女性	1(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	0(0) 0.0%	0(0) 0.0%
良性 石綿胸水	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
合計	小計	12(0) 0.5%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(0) 0.7%	6(0) 0.7%	1(0) 0.4%	0(0) 0.0%
	男性	9(0) 0.7%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(0) 1.3%	4(0) 0.8%	0(0) 0.0%	0(0) 0.0%
	女性	3(0) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(0) 0.6%	1(0) 1.1%	0(0) 0.0%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-1 参考) 石綿関連疾患と診断された者の年齢階層・性別分類表(平成30年度)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
参加者	合計	2,692	18	170	337	774	1102	279	12
	男性	1,669	10	114	188	469	705	175	8
	女性	1023	8	56	149	305	397	104	4
中皮腫	小計	2(1) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	1(1) 0.1%	0(0) 0.0%	0(0) 0.0%
	男性	1(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	1(1) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(1) 0.3%	0(0) 0.0%	0(0) 0.0%
肺がん	小計	8(6) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(2) 0.3%	6(4) 0.5%	0(0) 0.0%	0(0) 0.0%
	男性	6(4) 0.4%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(2) 0.4%	4(2) 0.6%	0(0) 0.0%	0(0) 0.0%
	女性	2(2) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	2(2) 0.5%	0(0) 0.0%	0(0) 0.0%
石綿肺	小計	5(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	4(0) 0.4%	0(0) 0.0%	0(0) 0.0%
	男性	5(0) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	4(0) 0.6%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
びまん性 胸膜肥厚	小計	11(1) 0.4%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	10(1) 0.9%	0(0) 0.0%	0(0) 0.0%
	男性	10(1) 0.6%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	9(1) 1.3%	0(0) 0.0%	0(0) 0.0%
	女性	1(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.3%	0(0) 0.0%	0(0) 0.0%
良性 石綿胸水	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
合計	小計	26(8) 1.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(2) 0.6%	21(6) 1.9%	0(0) 0.0%	0(0) 0.0%
	男性	22(5) 1.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	5(2) 1.1%	17(3) 2.4%	0(0) 0.0%	0(0) 0.0%
	女性	4(3) 0.4%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	4(3) 1.0%	0(0) 0.0%	0(0) 0.0%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-1 参考) 石綿関連疾患と診断された者の年齢階層・性別分類表(平成31年度)

(単位:人)

		合計	40歳未満	40～49歳	50～59歳	60～69歳	70～79歳	80～89歳	90歳以上
参加者	合計	2,112	5	109	263	527	913	284	11
	男性	1,238	4	61	153	296	543	170	11
	女性	874	1	48	110	231	370	114	0
中皮腫	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -
肺がん	小計	2(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	1(0) 0.4%	0(0) 0.0%
	男性	2(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	1(0) 0.6%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -
石綿肺	小計	2(1) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(1) 0.1%	0(0) 0.0%	1(0) 9.1%
	男性	2(1) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(1) 0.2%	0(0) 0.0%	1(0) 9.1%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -
びまん性 胸膜肥厚	小計	2(0) 0.1%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.1%	1(0) 0.4%	0(0) 0.0%
	男性	2(0) 0.2%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	1(0) 0.2%	1(0) 0.6%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -
良性 石綿胸水	小計	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	男性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -
合計	小計	6(1) 0.3%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(1) 0.3%	2(0) 1.2%	1(0) 9.1%
	男性	6(1) 0.5%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	3(1) 0.6%	2(0) 1.2%	1(0) 9.1%
	女性	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) 0.0%	0(0) -

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各年齢層の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-2 参考) 石綿関連疾患と診断された者のばく露・性別分類表(平成27年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	1,980		480		186		244		176		894	
	男性	1,142		417		144		60		124		397	
	女性	838		63		42		184		52		497	
中皮腫	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
肺がん	小計	8(0)	0.4%	5(0)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	3(0)	0.3%
	男性	5(0)	0.0%	4(0)	1.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
	女性	3(0)	0.0%	1(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.4%
石綿肺	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	2(0)	0.1%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.1%
	男性	2(0)	0.0%	1(0)	0.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	10(0)	0.5%	6(0)	1.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	4(0)	0.4%
	男性	7(0)	0.0%	5(0)	1.2%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.5%
	女性	3(0)	0.0%	1(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	2(0)	0.4%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 平成27年度の参加者の内「石綿ばく露の聴取のみを行った者66人」については、ばく露分類が不明のため、参加者数から除外している

※ 小数点以下第2位を四捨五入

(表6-2 参考) 石綿関連疾患と診断された者のばく露・性別分類表(平成28年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	1,902		489		196		227		153		837	
	男性	1,171		425		157		72		119		398	
	女性	731		64		39		155		34		439	
中皮腫	小計	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	1(1)	0.4%	0(0)	0.0%	0(0)	0.0%
	男性	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	1(1)	1.4%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
肺がん	小計	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.1%
	男性	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.3%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
石綿肺	小計	3(1)	0.2%	2(1)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.1%
	男性	3(1)	0.3%	2(1)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.3%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	3(1)	0.2%	1(0)	0.2%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%	1(1)	0.1%
	男性	3(1)	0.3%	1(0)	0.2%	0(0)	0.0%	1(0)	1.4%	0(0)	0.0%	1(1)	0.3%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	8(4)	0.4%	3(1)	0.6%	0(0)	0.0%	2(1)	0.9%	0(0)	0.0%	3(2)	0.4%
	男性	8(4)	0.7%	3(1)	0.7%	0(0)	0.0%	2(1)	2.8%	0(0)	0.0%	3(2)	0.8%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-2 参考) 石綿関連疾患と診断された者のばく露・性別分類表(平成29年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	2,260		557		238		294		225		946	
	男性	1,332		481		174		86		161		430	
	女性	928		76		64		208		64		516	
中皮腫	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
肺がん	小計	3(0)	0.1%	2(0)	0.4%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.2%	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	0.1%	0(0)	0.0%	1(0)	1.6%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
石綿肺	小計	4(0)	0.2%	3(0)	0.5%	0(0)	0.0%	1(0)	0.3%	0(0)	0.0%	0(0)	0.0%
	男性	3(0)	0.2%	2(0)	0.4%	0(0)	0.0%	1(0)	1.2%	0(0)	0.0%	0(0)	0.0%
	女性	1(0)	0.1%	1(0)	1.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	5(0)	0.2%	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.4%	2(0)	0.2%
	男性	4(0)	0.3%	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%	1(0)	0.6%	1(0)	0.2%
	女性	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	12(0)	0.5%	7(0)	1.3%	1(0)	0.4%	1(0)	0.3%	1(0)	0.4%	2(0)	0.2%
	男性	9(0)	0.7%	6(0)	1.2%	0(0)	0.0%	1(0)	1.2%	1(0)	0.6%	1(0)	0.2%
	女性	3(0)	0.3%	1(0)	1.3%	1(0)	1.6%	0(0)	0.0%	0(0)	0.0%	1(0)	0.2%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-2 参考) 石綿関連疾患と診断された者のばく露・性別分類表(平成30年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	2,692		765		388		277		324		938	
	男性	1,669		691		308		70		186		414	
	女性	1023		74		80		207		138		524	
中皮腫	小計	2(1)	0.1%	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.1%
	男性	1(0)	0.1%	1(0)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	1(1)	0.1%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	1(1)	0.2%
肺がん	小計	8(6)	0.3%	4(3)	0.5%	1(1)	0.3%	0(0)	0.0%	1(1)	0.3%	2(1)	0.2%
	男性	6(4)	0.4%	3(2)	0.4%	0(0)	0.0%	0(0)	0.0%	1(1)	0.5%	2(1)	0.5%
	女性	2(2)	0.2%	1(1)	1.4%	1(1)	1.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
石綿肺	小計	5(0)	0.2%	1(0)	0.1%	1(0)	0.3%	0(0)	0.0%	1(0)	0.3%	2(0)	0.2%
	男性	5(0)	0.3%	1(0)	0.1%	1(0)	0.3%	0(0)	0.0%	1(0)	0.5%	2(0)	0.5%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	11(1)	0.4%	6(1)	0.8%	3(0)	0.8%	0(0)	0.0%	0(0)	0.0%	2(0)	0.2%
	男性	10(1)	0.6%	6(1)	0.9%	2(0)	0.6%	0(0)	0.0%	0(0)	0.0%	2(0)	0.5%
	女性	1(0)	0.1%	0(0)	0.0%	1(0)	1.3%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	26(8)	1.0%	12(4)	1.6%	5(1)	1.3%	0(0)	0.0%	2(1)	0.6%	7(2)	0.7%
	男性	22(5)	1.3%	11(3)	1.6%	3(0)	1.0%	0(0)	0.0%	2(1)	1.1%	6(1)	1.4%
	女性	4(3)	0.4%	1(1)	1.4%	2(1)	2.5%	0(0)	0.0%	0(0)	0.0%	1(1)	0.2%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(表6-2 参考) 石綿関連疾患と診断された者のばく露・性別分類表(平成31年度)

(単位:人)

		合計		ア.主に 直接職歴		イ.主に 間接職歴		ウ.主に 家庭内ばく露		エ.主に 立ち入り等		オ.その他	
参加者	合計	2,112		506		238		246		197		925	
	男性	1,238		443		182		69		130		414	
	女性	874		63		56		177		67		511	
中皮腫	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
肺がん	小計	2(0)	0.1%	1(0)	0.2%	1(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.2%	1(0)	0.2%	1(0)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
石綿肺	小計	2(1)	0.1%	0(0)	0.0%	1(0)	0.4%	0(0)	0.0%	1(1)	0.5%	0(0)	0.0%
	男性	2(1)	0.2%	0(0)	0.0%	1(0)	0.5%	0(0)	0.0%	1(1)	0.8%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
びまん性 胸膜肥厚	小計	2(0)	0.1%	2(0)	0.4%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	2(0)	0.2%	2(0)	0.5%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
良性 石綿胸水	小計	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	男性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%
合計	小計	6(1)	0.3%	3(0)	0.6%	2(0)	0.8%	0(0)	0.0%	1(1)	0.5%	0(0)	0.0%
	男性	6(1)	0.5%	3(0)	0.7%	2(0)	1.1%	0(0)	0.0%	1(1)	0.8%	0(0)	0.0%
	女性	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%	0(0)	0.0%

※ ()内は疑い(内数)

※ 複数の疾患が出た者は全て計上している。各疾病の合計は実人数とは一致しない

※ 石綿によるものかどうかは特定されていない

※ 右欄は各ばく露の参加者に対する石綿関連疾患と診断された者の割合(疑いを含む)

※ 小数点以下第2位を四捨五入

(4) 受診カード

平成 27～30 年度に受診カードを渡した者 3,484 人のうち、2,619 人の状況を確認し、そのうち試行調査でＣＴ検査を受けた者が 1,223 人、何らかの検診等を受けた者が 655 人、試行調査や何らかの検診等を受けなかった者は 741 人であった。

表7 平成 27～30 年度の受診カード等※配布者についての平成 31 年度の状況

		計	試行調査で胸部CT検査を受けた者		肺がん検診のみを受けた者		その他の健診等を受けた者		検診等を受けない者		受診の有無を確認できなかった者	
埼玉県	さいたま市	156	2	1.3%	0	0.0%	19	12.2%	1	0.6%	134	85.9%
東京都	大田区	0	0	—	0	—	0	—	0	—	0	—
神奈川県	横浜市鶴見区	85	44	51.8%	0	0.0%	18	21.2%	11	12.9%	12	14.1%
岐阜県	羽島市	269	116	43.1%	53	19.7%	0	0.0%	0	0.0%	100	37.2%
大阪府	大阪市	66	45	68.2%	0	0.0%	0	0.0%	0	0.0%	21	31.8%
	堺市	116	23	19.8%	0	0.0%	0	0.0%	0	0.0%	93	80.2%
	上記以外	405	174	43.0%	21	5.2%	25	6.2%	28	6.9%	157	38.8%
兵庫県	神戸市	348	6	1.7%	1	0.3%	231	66.4%	99	28.4%	11	3.2%
	尼崎市	981	325	33.1%	150	15.3%	0	0.0%	506	51.6%	0	0.0%
	西宮市	100	13	13.0%	14	14.0%	3	3.0%	0	0.0%	70	70.0%
	芦屋市	20	7	35.0%	0	0.0%	6	30.0%	5	25.0%	2	10.0%
	加古川市	89	0	0.0%	0	0.0%	27	30.3%	17	19.1%	45	50.6%
	宝塚市	0	0	—	0	—	0	—	0	—	0	—
奈良県		414	327	79.0%	0	0.0%	20	4.8%	22	5.3%	45	10.9%
福岡県	北九州市門司区	189	65	34.4%	16	8.5%	26	13.8%	0	0.0%	82	43.4%
佐賀県	鳥栖市	95	18	18.9%	0	0.0%	23	24.2%	49	51.6%	5	5.3%
転居者調査		151	58	38.4%	2	1.3%	0	0.0%	3	2.0%	88	58.3%
計		3,484	1,223	35.1%	257	7.4%	398	11.4%	741	21.3%	865	24.8%

※ 氏人、住所、問い合わせ先、肺がん検診受診勧奨文、肺がん検診実施機関へのお願い、肺がん検診受診歴が記載されているもの