

エチレングリコール（エタン-1,2-ジオール）（CAS no. 107-21-1）

文献信頼性評価結果

示唆された作用							
エストロゲン	抗エストロゲン	アンドロゲン	抗アンドロゲン	甲状腺ホルモン	抗甲状腺ホルモン	脱皮ホルモン	その他*
—	—	—	—	—	—	—	—

○：既存知見から示唆された作用

—：既存知見から示唆されなかった作用

*その他：視床下部—下垂体—生殖腺軸への作用等

エチレングリコールの内分泌かく乱作用に関連する報告では、内分泌かく乱作用に関する試験対象物質として選定する根拠が得られなかった。

参考文献

- Ren L, Meldahl A and Lech JJ (1996) Dimethyl formamide (DMFA) and ethylene glycol (EG) are estrogenic in rainbow trout. *Chemico-Biological Interactions*, 102 (1), 63-67.
- Lamb JC, Maronpot RR, Gulati DK, Russell VS, Hommel-Barnes L and Sabharwal PS (1985) Reproductive and developmental toxicity of ethylene glycol in the mouse. *Toxicology and Applied Pharmacology*, 81 (1), 100-112.
- Carney EW, Tornesi B, Liberacki AB, Markham DA, Weitz KK, Luders TM, Studniski KG, Blessing JC, Gies RA and Corley RA (2011) The impact of dose rate on ethylene glycol developmental toxicity and pharmacokinetics in pregnant CD rats. *Toxicological Sciences*, 119 (1), 178-188.
- Price CJ, Kimmel CA, Tyl RW and Marr MC (1985) The developmental toxicity of ethylene glycol in rats and mice. *Toxicology and Applied Pharmacology*, 81 (1), 113-127.
- Neeper-Bradley TL, Tyl RW, Fisher LC, Kubena MF, Vrbanic MA and Losco PE (1995) Determination of a no-observed-effect level for developmental toxicity of ethylene glycol administered by gavage to CD rats and CD-1 mice. *Fundamental and Applied Toxicology*, 27 (1), 121-130.
- Tyl RW, Fisher LC, Kubena MF, Vrbanic MA and Losco PE (1995) Assessment of the developmental toxicity of ethylene glycol applied cutaneously to CD-1 mice. *Fundamental and Applied Toxicology*, 27 (2), 155-166.
- Tyl RW, Ballantyne B, Fisher LC, Fait DL, Savine TA, Dodd DE, Klonne DR and Pritts IM (1995) Evaluation of the developmental toxicity of ethylene glycol aerosol in the CD rat and CD-1 mouse by whole-body exposure. *Fundamental and Applied Toxicology*, 24 (1), 57-75.
- Tyl RW, Ballantyne B, Fisher LC, Fait DL, Dodd DE, Klonne DR, Pritts IM and Losco PE (1995) Evaluation of the developmental toxicity of ethylene glycol aerosol in CD-1 mice by nose-only exposure. *Fundamental and Applied Toxicology*, 27 (1), 49-62.
- Hong HL, Canipe J, Jameson CW and Boorman GA (1988) Comparative effects of ethylene glycol and ethylene glycol monomethyl ether exposure on hematopoiesis and histopathology in B6C3F1 mice. *Journal of Environmental Pathology, Toxicology and Oncology*, 8 (7 Spec No), 27-38.
- Lee YH, Huang WC, Huang JK and Chang LS (1996) Testosterone enhances whereas estrogen inhibits calcium oxalate stone formation in ethylene glycol treated rats. *Journal of Urology*, 156 (2 Pt 1), 502-505.
- Klug S, Merker HJ and Jackh R (2001) Effects of ethylene glycol and metabolites on *in vitro* development of rat embryos during organogenesis. *Toxicology in Vitro*, 15 (6), 635-642.

(令和元年度第2回 EXTEND2016 化学物質の内分泌かく乱作用に関する検討会 資料1より抜粋)