

OResults of Radioactive Material Monitoring of Aquatic Organisms (Location C along the Uda River)

<Location C along the Uda River: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
C-6	○	○	○	○	○	○

<Location C along the Uda River: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Transparency (cm)
C-6	37.7764°	140.8877°	2023/8/30	10:15	10:31	27.9	28.1	Sand	10YR4/2	None	0.34	>50

<Location C along the Uda River: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
	Latitude	Longitude	Date	Time (water)		(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
C-6	37.7764°	140.8877°	2023/8/30	10:15	7.9	1.2	3.6	9.0	13.4	0.07	1.3	3	2.8	N.D.(0.0015)	0.0045	0.0010

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location C along the Uda River: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
C-6	37.7764°	140.8877°	2023/8/30	10:31	7.8	489	16.8	1.0	0.6	2.680	43.2	33.9	15.0	2.0	2.0	3.9	1.7	9.5	0.76	40	0.18

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location C along the Uda River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
C-6	The main stream of the Uda River	37.7764°	140.8877°	2023/8/26	Algae/plant	-	-	-	-	Sediment deposited on riverbed (Including algae)	-	0.011	-	-	-	46	N.D.(8.7)	46	-
					Algae/plant	Zygnematophyceae	Zygnematales	Zygnemataceae	<i>Spirogyra</i> sp.	Spirogyra	-	0.14	-	-	-	0.81	N.D.(0.43)	0.81	-
					Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Caddisfly	77	0.022	Larva	-	-	7.7	N.D.(1.9)	7.7	-
					Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	Dragonfly	24	0.0070	Larva (Dragonfly larva)	-	-	N.D.	N.D.(4.2)	N.D.(3.4)	-
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Melligomphus viridicostus</i>	Dragonfly									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	Dragonfly									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Davidius</i> sp.	Dragonfly									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Sinogomphus flavolimbatus</i>	Dragonfly									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Asiagomphus melaenops</i>	Dragonfly									
					Arthropoda	Insecta	Odonata	Libellulidae	<i>Orthetrum albistylum speciosum</i>	Common skimmer									
					Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	Dobsonfly	25	0.015	Larva	-	-	2.9	N.D.(2.1)	2.9	-
					Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Lake prawn	25	0.032	Juvenile,Imago	-	-	2.1	N.D.(1.3)	2.1	-
					Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp	218	0.063	Juvenile,Imago	-	-	3.4	N.D.(0.70)	3.4	-
					Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	7	0.071	Juvenile	-	-	5.2	N.D.(0.75)	5.2	-
					Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	1	0.23	Mature fish	Empty stomach	Viscera removed	4.5	N.D.(1.0)	4.5	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Pseudaspius hakonensis</i>	Japanese dace	9	0.24	Immature fish, Mature fish	Obscure digesta	Viscera removed	3.8	N.D.(0.37)	3.8	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	Pale break	145	0.45	Immature fish	-	-	1.9	N.D.(0.31)	1.9	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Candidia temminckii</i>	Dark chub	51	0.15	Immature fish	-	-	2.0	N.D.(0.31)	2.0	-
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Ayu sweetfish	256	2.4	Immature fish	-	-	6.6	N.D.(0.86)	6.6	0.15
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Gymnogobius urotaenia</i>	Floating goby	4	0.032	Immature fish	-	-	3.6	N.D.(1.5)	3.6	-
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius	70	0.12	Immature fish, Mature fish	-	-	3.5	N.D.(0.60)	3.5	-
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius									
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius</i> sp.	Freshwater goby									
					Coarse Particulate Organic Matter	-	-	-	-	Water-bottom leaf litter	-	0.22	-	-	-	8.3	N.D.(1.7)	8.3	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.