OResults of Radioactive Material Monitoring of Aquatic Organisms (Location H in Lake Akimoto)

<Location H in Lake Akimoto: Samples collected>

Items	Genera	al items	Radioactive materials								
Locations	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)					
H-1	0 0		0	0	0	0					
H-2	0	0	0	-	0	-					

<Location H in Lake Akimoto: Site measurement item>

Location II ill Lake A	Cocation it in Lake Akimoto: Site measurement tem?													
Items		longitude of the ation		Survey date and time		Water		Sedi	ment		Ot	ther		
Locations	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)		
H-1(Surface layer)	- 37.6575°	140.1264°	- 2023/12/2	09:36	- 09:52	5.5	6.0	Ooze	7.5Y5/3	Plant pieces	6.9	2.3		
H-1(Bottom layer)	37.0373	140.1204		09:41								2.3		
H-2(Surface layer)	37.6616°	140.1226°		08:57	09:18	6.5	7.2	Ooze	7.5Y5/3	Plant pieces	13.8	2.4		
H-2(Bottom layer)		140.1226		09:01	09:18							2.4		

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Water>

	Items	Latitude and longitude of the location		Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
Loca	eations	Latitude	Longitude	Date	Time (water)		(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
H-1((Surface layer)	37.6575°	140.1264°		09:36	7.1	0.6	2.7	10.3	5.6	0.04	1.3	3	2.8	N.D.(0.0014)	0.0048	=
H-1((Bottom layer)	37.6575°	140.1204	2023/12/2	09:41	7.1	0.9	3.1	10.3	5.8	0.03	1.4	3	3.0	N.D.(0.0014)	0.0033	0.0012
H-2((Surface layer)	37.6616°	140.1226°	2023/12/2	08:57	7.2	0.6	2.9	10.8	5.6	0.03	1.3	2	2.4	N.D.(0.0014)	0.0047	-
H-2((Bottom layer)		140.1226*		09:01	7.1	0.7	2.9	11.1	5.7	0.03	1.2	2	2.3	N.D.(0.0013)	0.0039	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Sediment>

Itame	Latitude and	ongitude of the	Summer d	ate and time										Grain si	ze distribution					Cs-137					
nems	loca	ntion	Survey u	ate and time	pH	Redox potential	Water content	IL	TOC	Soil particle	Gravel	Coarse sand	Medium sand	Fine sand	Silt	Clay	Median grain	Maximum	Cs-134		Sr-90				
Locations	Latitude	Longitude	Date	Time (sediment)		E _{N.H.E}				density	(2-75mm)	(0.85-2mm)	(0.25-0.85mm)	(0.075-0.25mm)	(0.005-0.075mm)	(Less than 0.005mm)	diameter	grain diameter							
Locatons	Latitude	Longitude	Dute	Time (seament)		(mV)	(%)	(%)	(mg/g-dry)	(g/cm ³)	(%)	(%)	(%)	(%)	(%)	(%)	(mm)	(mm)	(Bq/kg-dry)	(Bq/kg-dry)	(Bq/kg-dry)				
H-1	37.6575°	140.1264°	2023/12/2	2023/12/2	2023/12/2	2023/12/2	2023/12/2	09:52	6.8	121	53.3	9.7	35.0	2.530	0.0	0.0	0.0	0.6	50.7	48.7	0.0053	0.85	13	760	1.2
H-2	37.6616°	140.1226°				09:18	6.6	141	53.9	14.0	51.0	2.460	0.0	0.0	0.0	0.1	43.0	56.9	0.0036	0.85	3.5	260	-		

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight		Note			dioactive cesium (Bq/kg-w	Sr-90	
		Latitude	Longitude							-		(kg-wet)	Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	(Bq/kg-wet)
H-1		37.6575°	140.1264°	2023/12/1	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	=	0.030	-	-	-	N.D.	N.D.(1.3)	N.D.(1.3)	-
H-2	In the lake	37.6616°	140.1226°	2023/12/2	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	Pseudaspius hakonensis	Japanese dace	10	2.5	Mature fish	Obscure digesta	Viscera removed	23	N.D.(1.2)	23	-
H-3		37.6653°	140.1329°		Vertebrata	Osteichthyes	Perciformes	Centrarchidae	Micropterus nigricans	Largemouth bass	2	0.40	Immature fish	Palaemon paucidens	Viscera removed	28	N.D.(2.2)	28	-
					Algae/plant	-	-	-	-	Sediment deposited on riverbed (Including algae)	-	0.021	-	-	-	10	N.D.(1.9)	10	-
					Arthropoda	Insecta	Plecoptera	Perlidae	Xanthoneuria sp.	Stonefly	53	0.0050	Larva	-	-	N.D.	N.D.(6.3)	N.D.(5.1)	-
				ŗ	Arthropoda	Insecta	Plecoptera	Perlidae	Kamimuria quadrata	Stonefly									
H-3	Inflowing rivers	37.6653°	140.1329°	2023/12/1	Arthropoda	Insecta	Plecoptera	Perlidae	Calineuria sp.	Stonefly									
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	Oncorhynchus masou masou	Masu salmon	10	0.17	Immature fish	Aquatic insect	Viscera removed	5.7	N.D.(0.80)	5.7	-
					Coarse Particulate Organic Matter	-	-	-	-	Water-bottom leaf litter	-	0.20	-	-	-	2.9	N.D.(0.31)	2.9	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

 $*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40 \mum-mesh).$

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.