OResults of Radioactive Material Monitoring of Aquatic Organisms (Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J)

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Samples collected>

Items	Genera	al items	Radioactive materials								
Locations	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)					
J-1	0	0	0	0	0	0					

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Site measurement item>

Items		longitude of the ation	Survey date and time			Water		Sedi		Other		
Locations	Latitude	Latitude Longitude Date Time (water) Ti		Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)	
J-1(Surface layer)	37.4203°	140.1008°	2023/12/2	12:50	13:30	8.9	9.1	Sand	7.5Y6/3	Shells	3.5	>3.5
J-1(Bottom layer)		140.1008*	2023/12/2	12:56		9.1						73.3

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		rude of the Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
Locations	Latitude	Longitude	Date	Time (water)		(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
J-1(Surface layer)	37.4203°	140.1008°	2023/12/2	12:50	6.9	<0.5	1.3	11.1	12.0	0.06	0.7	<1	0.4	N.D.(0.0015)	0.0043	-
J-1(Bottom layer)		140.1000	2023/12/2	12:56	6.8	<0.5	1.3	11.1	12.1	0.06	0.7	<1	0.4	N.D.(0.0013)	0.0043	0.00076

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: General survey items/Analysis of radioactive materials Sediment>

Itame	Latitude and longitude of the location		Survey date and time								Grain size distribution										
nems					pH	Redox potential	Water content	IL	TOC	Soil particle	Gravel	Coarse sand	Medium sand	Fine sand	Silt	Clay	Median grain	Maximum	Cs-134	Cs-137	Sr-90
Locations	Latitude	Longitude	Dete	Time (sediment)		$E_{N.H.E}$				density	(2-75mm)	(0.85-2mm)	(0.25-0.85mm)	(0.075-0.25mm)	(0.005-0.075mm)	(Less than 0.005mm)	diameter	grain diameter			1
Locations			Date	Time (sediment)		(mV)	(%)	(%)	(mg/g-dry)	(g/cm ³)	(%)	(%)	(%)	(%)	(%)	(%)	(mm)	(mm)	(Bq/kg-dry)	(Bq/kg-dry)	(Bq/kg-dry)
J-1	37.4203°	140.1008°	2023/12/2	13:30	6.0	417	29.0	2.4	5.7	2.680	0.1	0.9	16.9	68.5	8.6	5.0	0.15	4.8	1.2	63	0.14

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

< Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Analysis items Aquatic organisms>

Lake mawasmro (nor	th lakeside) I / Lake ina			Aquatic organisms>															
Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight	Note			Radioactive cesium (Bq/kg-wet)			Sr-90
		Latitude	Longitude									(kg-wet)	Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	(Bq/kg-wet)
I-1 I-2 (north lakeside)	Within the lake and Nagase River	37.5047° 37.4995°	140.1143° 140.1409°	2023/12/2	Coarse Particulate Organic Matter	-	-	-	-	Water-bottom leaf litter	-	0.21	-	-	-	2.0	N.D.(0.27)	2.0	-
					Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.033	-	-	-	N.D.	N.D.(1.1)	N.D.(1.1)	-
					Mollusca	Gastropoda	Discopoda	Pleuroceridae	Semisulcospira libertina	Freshwater snail	16	0.014	Imago	=	Molluscous part	N.D.	N.D.(2.5)	N.D.(2.4)	=
J-1	Within the lake and	37.4203°	140.1008°	2023/12/2	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	Opsariichthys platypus	Pale break	22	0.083	Immature fish	=	-	6.1	N.D.(1.3)	6.1	-
(south lakeside)	around the Oninuma	37.4203	140.1008	2023/12/2	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	Rhodeus ocellatus ocellatus	Rosy bitterling	115	0.093	Mature fish	=	-	6.5	N.D.(0.94)	6.5	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	Pseudorasbora parva	Stone moroko	28	0.023	Immature fish, Mature fish	-	-	2.2	N.D.(1.5)	2.2	-

^{*1:} Organisms were collected in or around the targeted water areas.

^{*2:} When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

^{*3:} For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

^{*4:} Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

 $^{*5:} Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40 \mum-mesh).$

^{*6:} River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

^{*7:} N.D. means to be below the detection limit and figures in parentheses show the detection limit.

^{*8:} Activity concentrations include counting errors, but the details are omitted here.