

**OResults of Radioactive Material Monitoring of Aquatic Organisms (Location H in Lake Akimoto)**

<Location H in Lake Akimoto: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
H-1	○	○	○	○	○	○
H-2	○	○	○	-	○	-

<Location H in Lake Akimoto: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
H-1(Surface layer)	37.6575°	140.1264°	2024/6/19	08:30	08:54	20.6	11.5	Ooze	7.5Y5/2	Plant pieces	12.2	3.4
H-1(Bottom layer)				08:33		12.7						
H-2(Surface layer)	37.6616°	140.1226°		09:09	09:31	20.4	11.7	Ooze	7.5Y5/3	None	4.9	2.9
H-2(Bottom layer)				09:13		17.5						

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
H-1(Surface layer)	37.6575°	140.1264°	2024/6/19	08:30	7.3	1.7	3.6	8.8	5.0	0.03	1.5	7	2.2	N.D.(0.0014)	0.0030	-
H-1(Bottom layer)				08:33	6.9	1.0	3.2	8.2	4.7	0.03	1.5	2	1.3	1.3	N.D.(0.0014)	0.0019
H-2(Surface layer)	37.6616°	140.1226°		09:09	7.2	1.6	3.2	8.9	4.9	0.03	1.3	4	2.4	N.D.(0.0015)	0.0078	-
H-2(Bottom layer)				09:13	7.1	1.7	4.0	9.7	4.7	0.03	1.5	6	3.4	3.4	N.D.(0.0016)	0.011

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>h</sub> (H <sub>2</sub> E) (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
H-1	37.6575°	140.1264°	2024/6/19	08:54	6.6	207	69.9	11.3	37.0	2.490	0.0	0.0	0.1	0.6	68.4	30.9	0.0097	2.0	21	1500	1.1
H-2				09:31	6.5	123	69.5	15.6	53.0	2.430	0.0	0.0	0.0	0.1	70.7	29.2	0.0092	0.85	4.9	350	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)						
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137							
H-1 H-2 H-3	In the lake	37.6575° 37.6616° 37.6653°	140.1264° 140.1226° 140.1329°	2024/6/18	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.034	-	-	-	N.D.	N.D.(1.4)	N.D.(1.4)	-						
				2024/6/19	Arthropoda	Malacostraca	Decapoda	Astacidae	<i>Pacifastacus leniusculus trowbridgii</i>	Signal crayfish	56	1.6	Imago	-	-	-	13	N.D.(1.4)	13	5.4					
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	11	3.2	Mature fish	Obscure digesta	Viscera removed	30	N.D.(1.2)	30	0.67						
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	<i>Carassius auratus langsdorfii</i>	Mature fish	5	1.0	Mature fish	Obscure digesta	Viscera removed	12	N.D.(1.1)	12	0.89					
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp	1	5.5	Mature fish	Obscure digesta	Viscera removed	10	N.D.(1.2)	10	0.86						
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbuis</i>	<i>Hemibarbus barbuis</i>	Immature fish, Mature fish	5	1.8	Immature fish, Mature fish	Obscure digesta	Viscera removed	16	N.D.(1.3)	16	1.0					
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Hypomesus nipponensis</i>	Japanese smelt	55	0.23	Mature fish	-	-	-	6.0	N.D.(0.57)	6.0	-					
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	10	3.5	Mature fish	Japanese smelt	Viscera removed	22	N.D.(1.2)	22	0.27						
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou masou</i>	Seema	3	0.53	Immature fish	Empty stomach	Viscera removed	12	N.D.(1.3)	12	-						
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus salmoides</i>	Largemouth bass	1	0.36	Immature fish	Empty stomach	Viscera removed	29	N.D.(1.8)	29	-						
Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	2	0.51	Immature fish	Empty stomach	Viscera removed	23	N.D.(1.4)	23	-											
H-3	Inflowing rivers	37.6650°	140.1392°	2024/6/18	Algae/plant	-	-	-	-	Riverbed Deposits (Include algae)	-	0.084	-	-	-	1.3	N.D.(0.80)	1.3	-						
					Arthropoda	Insecta	Plecoptera	Perlidae	<i>Calineuria sp.</i>	Calineuria	50	0.011	Larva	-	-	-	N.D.	N.D.(3.4)	N.D.(2.7)	-					
					Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus pollux</i>	Japanese fluvial sculpin	11	0.070	Immature fish	-	-	-	3.0	N.D.(0.92)	3.0	-					
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	14	0.30	Immature fish	-	-	-	3.7	N.D.(0.35)	3.7	-					
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	8	0.16	Immature fish, Mature fish	Obscure digesta	Viscera removed	7.3	N.D.(0.62)	7.3	-						
					Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.23	-	-	-	-	2.4	N.D.(0.33)	2.4	-					
H-4	Within the lake and rivers in the vicinity	37.6551°	140.1181°	2024/6/18	Algae/plant	Monocotyledoneae	Alismatales	Hydrocharitaceae	<i>Elodea nuttallii</i>	Western waterweed	-	0.33	-	-	-	2.7	N.D.(0.28)	2.7	-						
				2024/6/19	Arthropoda	Insecta	Odonata	Corduliidae	<i>Somatochlora uchidai</i>	<i>Somatochlora uchidai</i>	43	0.049	Larva (Dragonfly larva)	-	-	4.7	N.D.(0.87)	4.7	-						
					Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	<i>Anotogaster sieboldii</i>															
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Stylogomphus suzukii</i>	<i>Stylogomphus suzukii</i>															
					Arthropoda	Insecta	Odonata	Aeshnidae	<i>Aeshna juncea</i>	Common hawker	30	0.037	Imago	-	-	Molluscos part	3.5	N.D.(0.99)	3.5	-					
					Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	<i>Semisulcospira libertina</i>															
					Vertebrata	Amphibia	Anura	-	-	Frog											111	0.077	Larva(Tadpole)	-	-
Vertebrata	Amphibia	Caudata	Salamandridae	<i>Cynops pyrrhogaster</i>	<i>Cynops pyrrhogaster</i>	4	0.023	Imago	-	-											-	4.8	N.D.(1.5)	4.8	-

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.