

令和6年度 SO₄²⁻ 月間・年間平均濃度における海塩粒子の寄与率

単位：%

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年間
利尻	19.1	32.9	13.7	22.3	21.8	67.8	70.7	69.4	66.5	57.4	72.3	54.2	51.2
落石岬	*	*	*	*	*	*	*	*	*	*	*	*	*
赤城	*	*	*	*	*	*	5.2	*	*	*	*	*	5.2
小笠原	48.3	46.4	51.4	60.5	87.9	87.8	94.2	88.5	74.2	76.6	68.9	45.9	81.0
佐渡関岬	32.6	60.3	31.2	41.7	47.5	40.3	40.4	77.0	72.6	68.8	77.0	24.2	60.0
八方尾根	3.0	3.2	1.0	2.0	0.7	3.1	2.7	2.3	12.4	8.3	16.8	2.5	5.2
伊自良湖	9.1	9.0	4.0	3.1	3.0	10.0	12.5	33.5	41.2	24.5	*	*	9.8
尼崎	*	*	*	*	*	*	*	*	*	*	*	*	*
隠岐	*	*	*	*	*	*	*	*	*	*	*	*	*
禰原	4.7	7.6	3.1	2.7	4.1	24.3	43.5	21.5	30.1	25.0	17.1	16.9	16.1
対馬	14.3	24.6	8.7	17.6	*	59.5	31.4	52.8	13.8	36.3	25.5	29.4	26.4
えびの	3.5	1.3	6.8	6.9	1.8	10.6	5.9	32.5	13.2	19.3	10.5	7.9	7.2
辺戸岬	*	*	*	*	*	*	*	*	*	*	*	*	*
東京	23.7	15.5	7.1	2.5	13.9	24.5	14.2	23.9	17.7	8.6	4.0	10.1	14.4

網掛けの数値：海塩粒子寄与率が75%を超えたもの。

*：月間全て欠測

令和6年度 Ca²⁺ 月間・年間平均濃度における海塩粒子の寄与率

単位：％

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年間
利尻	9.2	25.3	16.3	46.2	53.3	59.1	68.4	73.9	77.8	69.8	72.7	55.6	55.7
落石岬	*	*	*	*	*	*	*	*	*	*	*	*	*
赤城	*	*	*	*	*	*	6.2	*	*	*	*	*	6.2
小笠原	40.0	29.9	32.9	49.6	88.7	74.4	76.3	90.0	61.4	68.9	68.3	57.6	74.6
佐渡関岬	16.7	60.4	30.8	48.7	63.9	67.8	57.4	86.1	61.8	85.0	67.8	22.7	60.6
八方尾根	0.8	2.5	0.8	4.9	4.1	6.5	3.6	4.9	19.7	7.8	16.3	0.9	4.4
伊自良湖	8.4	12.7	8.4	4.9	2.8	13.3	17.3	38.9	41.4	24.8	*	*	12.6
尼崎	*	*	*	*	*	*	*	*	*	*	*	*	*
隠岐	*	*	*	*	*	*	*	*	*	*	*	*	*
禰原	10.2	14.7	11.6	20.5	8.7	64.0	80.0	31.6	24.9	29.0	26.8	22.4	30.4
対馬	27.2	28.9	28.6	32.1	*	76.1	63.0	73.9	9.8	34.7	30.5	17.1	38.7
えびの	7.5	2.5	6.8	5.6	4.9	6.8	8.7	19.4	3.9	8.9	11.9	9.6	7.9
辺戸岬	*	*	*	*	*	*	*	*	*	*	*	*	*
東京	15.6	12.6	5.9	2.3	15.0	24.5	13.3	21.6	7.0	5.7	5.2	7.9	12.4

網掛けの数値：海塩粒子寄与率が75%を超えたもの。

*：月間全て欠測