

表5(a) 平成19年度 SO₄²⁻濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	50.1	20.6	90.2	25.7	31.5	7.2	29.7	52.1	35.9	43.4	40.4	57.4	28.0	223	1.7
札幌	24.6	19.6	22.9	18.5	21.8	12.6	27.2	17.1	14.0	18.9	27.4	51.1	20.4	102	5.3
落石岬	20.6	20.2	9.8	7.2	14.4	14.1	21.8	43.8	67.2	65.9	15.2	10.6	17.9	313	2.5
竜飛岬	21.2	35.8	31.7	31.6	13.9	20.1	51.6	27.6	29.5	282	69.6	68.6	30.0	605	7.3
八幡平	28.9	21.5	8.1	7.1	16.2	3.2	11.8	14.2	13.5	23.3	26.7	36.7	14.1	238	0.0
籠岳	37.1	19.1	21.2	14.1	12.4	7.1	4.0	14.2	11.5	46.2	26.2	24.7	15.2	308	2.4
尾花沢	28.5	9.3	9.2	8.6	15.6	6.2	4.5	21.8	19.1	15.8	28.3	17.6	15.2	82.0	2.4
筑波	32.5	23.2	12.8	17.3	15.5	8.5	6.2	22.1	14.8	20.9	9.0	29.5	17.3	145	1.2
赤城	25.6	12.7	11.0	10.5	5.1	4.6	4.4	24.2	6.9	6.6	5.5	14.6	8.1	117	1.1
小笠原	15.8	8.0	8.6	17.9	6.1	12.9	9.5	9.9	26.5	14.0	13.3	12.1	12.2	169	1.6
佐渡関岬	62.8	55.9	15.1	11.2	14.5	15.0	25.3	79.3	39.0	59.4	139	31.0	32.4	1070	3.5
新潟巻	27.0	30.1	12.8	12.0	13.0	21.7	17.7	49.7	40.1	48.2	45.0	42.4	27.1	288	2.2
八方尾根	34.0	16.2	8.1	13.8	7.1	3.2	3.4	12.8	13.8	23.3	22.7	26.5	12.4	274	0.0
越前岬	38.9	29.5	11.0	22.0	8.3	9.6	10.0	52.0	56.4	40.3	49.1	32.2	30.6	469	0.0
伊自良湖	36.7	23.6	16.0	13.8	20.1	10.1	14.8	23.4	17.1	12.2	23.4	22.2	17.1	133	2.5
犬山	51.0	17.1	19.9	13.5	13.5	7.4	15.2	27.5	17.7	31.3	15.2	15.4	15.5	212	1.3
潮岬	55.4	23.5	20.2	21.8	16.1	11.1	14.9	22.2	26.2	55.1	38.5	36.6	25.0	297	2.6
京都八幡	44.3	21.8	18.6	14.0	10.4	14.4	10.9	20.9	14.3	19.9	10.9	22.2	16.7	285	1.8
尼崎	53.3	15.2	15.4	16.6	16.5	13.3	15.4	6.0	15.7	17.1	10.9	15.5	17.0	236	2.3
倉橋島	24.1	26.0	21.5	12.6	33.3	13.6	14.7	16.7	15.9	16.7	13.3	28.3	18.6	202	0.0
隠岐	70.6	27.1	5.9	16.7	10.5	32.3	14.0	95.5	93.8	57.4	68.8	51.1	31.3	504	1.1
蟠竜湖	50.9	11.6	14.8	16.7	8.7	14.2	9.8	43.5	96.6	34.3	51.0	41.3	27.9	162	5.6
栲原	19.0	10.5	12.6	4.9	6.6	3.7	6.3	8.0	24.5	13.7	19.8	20.4	9.9	337	0.2
筑後小郡	125	47.5	29.8	10.7	10.1	13.0	12.2	26.4	37.7	21.1	34.4	38.7	27.4	661	3.1
大分久住	26.7	23.7	14.1	8.2	9.1	9.0	4.8	19.0	27.4	12.7	110	18.1	13.1	110	3.3
対馬	44.4	31.0	24.2	22.9	19.7	27.0	14.4	35.6	51.5	27.5	10.6	32.7	28.4	207	0.0
五島	31.1	41.0	13.9	37.5	*	30.5	3.4	34.6	21.7	16.1	34.0	44.7	30.9	434	0.0
えびの	12.4	30.8	16.2	20.0	11.7	7.1	14.7	0.7	23.8	25.8	26.3	22.9	17.7	217	0.0
屋久島	20.5	19.0	18.2	36.7	28.1	9.1	36.6	41.8	31.3	29.3	63.6	35.4	26.7	92.9	5.1
辺戸岬	15.4	14.0	4.4	19.1	5.0	8.8	23.5	18.0	22.2	38.9	34.0	21.7	15.2	2510	0.4
東京	30.3	20.1	27.8	18.4	52.0	12.1	7.9	26.5	13.1	31.1	8.5	21.1	17.3	326	1.3
最大値 ¹⁾	125	55.9	90.2	37.5	52.0	32.3	51.6	95.5	96.6	282	139	68.6	32.4	2510	7.3
最小値 ¹⁾	12.4	8.0	4.4	4.9	5.0	3.2	3.4	0.7	6.9	6.6	5.5	10.6	8.1	82.0	0.0
全国平均	37.4	23.4	18.3	16.8	15.6	12.3	14.9	29.3	30.3	37.7	35.2	30.4	20.5		
標準偏差	21.8	10.8	15.0	7.9	9.9	7.3	10.7	20.6	22.2	48.0	29.9	14.1	7.2		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表6(a) 平成19年度 nss-SO₄²⁻濃度 月・年平均値単位: μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	46.5	19.0	88.0	20.9	28.2	3.8	10.6	16.9	23.0	14.1	17.2	54.3	19.4	132	0.0
札幌	21.6	18.4	22.8	18.2	21.3	9.4	20.4	12.7	12.2	11.3	17.3	47.0	16.1	102	4.8
落石岬	14.1	14.0	8.5	5.3	13.6	6.7	6.8	8.7	29.8	13.3	6.8	4.5	9.4	205	0.2
竜飛岬	16.9	23.1	30.0	18.8	11.6	6.8	22.3	8.7	14.9	27.6	31.0	41.4	15.7	411	0.0
八幡平	28.1	20.8	8.0	7.0	15.9	2.7	10.2	12.7	12.4	12.2	21.9	35.4	13.1	232	0.0
籠岳	36.1	18.6	20.8	13.6	12.1	4.8	3.5	9.5	10.2	43.4	21.3	22.2	14.1	303	1.5
尾花沢	27.7	9.1	9.1	8.5	15.3	4.8	4.1	14.3	15.3	11.3	17.9	16.9	12.1	81.5	2.4
筑波	31.1	22.0	12.1	16.8	15.2	5.9	5.6	20.0	13.7	20.4	8.5	27.2	16.1	143	1.1
赤城	24.9	12.5	10.7	10.3	5.0	4.0	3.7	23.4	6.3	6.5	5.4	13.8	7.7	115	1.0
小笠原	10.5	5.7	4.2	15.1	1.8	1.8	1.1	1.6	7.4	7.3	4.6	6.5	4.8	64.3	0.0
佐渡関岬	50.9	36.6	13.8	9.6	12.2	10.0	11.5	28.3	21.6	28.6	41.6	22.7	20.2	328	0.0
新潟巻	25.9	26.8	12.4	11.6	11.7	17.6	12.4	32.9	24.4	33.3	29.2	36.9	20.1	220	2.1
八方尾根	33.5	16.1	8.1	13.8	7.1	3.1	3.3	12.4	13.3	22.6	21.7	25.6	12.2	253	0.0
越前岬	33.5	24.9	10.5	21.4	7.0	7.8	7.3	32.4	26.6	16.4	23.4	27.1	19.3	411	0.0
伊自良湖	35.0	23.2	15.8	13.6	19.5	9.3	14.3	20.6	14.9	10.7	19.0	20.9	16.3	129	2.4
犬山	47.3	16.5	19.6	13.2	13.0	6.7	14.4	26.5	16.8	30.8	13.9	14.3	15.0	207	1.1
潮岬	42.4	18.7	15.8	10.7	13.2	7.1	10.0	18.4	17.1	49.3	17.6	21.1	17.2	179	0.6
京都八幡	42.6	21.0	18.4	13.8	10.2	14.0	10.3	18.7	13.7	19.4	9.5	20.9	16.1	261	1.7
尼崎	51.3	14.4	14.7	16.3	16.0	12.7	14.7	5.7	15.2	16.7	10.3	14.6	16.3	232	2.2
倉橋島	23.1	25.5	20.9	12.1	28.6	13.3	14.1	16.0	14.8	16.1	12.5	27.0	17.8	188	0.0
隠岐	39.9	20.8	5.0	15.1	6.7	22.5	8.0	26.1	40.0	16.6	22.3	38.4	15.7	266	0.7
蟠竜湖	40.4	11.0	14.2	13.5	7.8	12.3	8.5	32.4	70.2	26.8	36.4	34.8	22.3	156	5.1
栲原	18.4	10.2	12.3	4.8	4.4	3.0	6.0	7.5	20.5	12.6	18.0	19.2	9.0	324	0.1
筑後小郡	118	45.9	28.8	10.2	9.5	12.4	11.3	25.3	35.7	20.4	31.6	36.7	26.1	618	2.9
大分久住	25.8	23.3	13.2	8.1	8.9	8.7	4.5	18.8	25.7	12.4	108	17.8	12.8	108	3.0
対馬	41.7	30.1	23.1	21.7	15.6	24.1	12.4	16.5	48.1	23.6	9.6	30.2	26.3	199	0.0
五島	25.4	36.1	10.8	33.2	*	26.8	2.5	29.4	14.9	14.2	25.5	40.9	26.6	352	0.0
えびの	11.8	30.3	15.8	18.4	11.4	6.4	13.4	0.7	21.3	25.0	25.7	21.5	16.9	194	0.0
屋久島	16.8	16.8	16.4	28.5	12.1	7.3	31.7	29.9	21.5	20.7	54.7	26.5	19.7	90.7	3.2
辺戸岬	10.2	11.0	2.3	4.9	1.7	2.6	13.5	5.1	6.6	16.4	10.6	11.7	6.4	286	0.2
東京	28.4	19.2	26.7	17.9	48.2	7.4	6.9	24.9	12.3	30.5	7.7	18.4	15.2	325	1.3
最大値 ¹⁾	118	45.9	88.0	33.2	48.2	26.8	31.7	32.9	70.2	49.3	108	54.3	26.6	618	5.1
最小値 ¹⁾	10.2	5.7	2.3	4.8	1.7	1.8	1.1	0.7	6.3	6.5	4.6	4.5	4.8	64.3	0.0
全国平均	32.9	20.7	17.2	14.4	13.5	9.2	10.3	18.0	20.7	20.3	22.6	25.7	16.0		
標準偏差	19.7	8.7	14.8	6.5	9.2	6.4	6.4	9.4	13.2	9.9	19.4	11.7	5.4		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表7(a) 平成19年度 NO₃⁻濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	45.7	13.4	86.6	17.5	20.1	4.5	8.8	17.5	16.2	13.4	21.2	39.0	15.8	119	0.0
札幌	24.0	16.6	27.1	17.2	23.6	9.5	22.4	9.8	8.8	13.4	23.9	58.8	17.5	176	1.6
落石岬	10.5	11.0	7.5	5.7	15.4	5.8	7.7	9.2	37.0	12.8	13.9	5.4	9.2	202	0.0
竜飛岬	22.0	34.2	36.4	32.8	13.0	11.6	33.2	10.9	21.9	99.1	59.8	84.7	23.4	781	2.9
八幡平	28.8	15.9	8.6	9.0	13.6	3.5	9.2	11.1	9.9	16.6	38.7	40.1	12.9	247	0.2
籠岳	44.3	16.5	15.8	16.1	18.4	5.5	5.9	7.1	9.6	50.2	25.2	30.7	15.5	439	1.4
尾花沢	27.5	11.1	10.2	8.1	13.8	4.2	5.8	9.9	11.8	15.8	23.1	27.9	12.8	55.4	2.3
筑波	51.5	18.4	23.7	20.5	35.9	11.3	9.5	20.6	18.3	37.6	16.8	34.9	22.6	227	2.6
赤城	37.4	14.5	29.5	27.5	20.8	9.2	24.0	35.5	10.7	18.5	9.4	45.0	19.1	489	0.0
小笠原	9.7	3.9	6.2	11.3	2.0	2.1	0.3	2.6	6.1	6.6	5.0	4.9	4.4	153	0.0
佐渡関岬	53.1	47.3	16.4	13.1	14.2	15.0	17.7	31.4	27.7	52.2	129	44.8	27.0	714	2.8
新潟巻	24.0	32.8	13.7	14.1	12.9	19.1	14.3	29.2	21.7	49.0	34.1	51.6	21.7	470	1.9
八方尾根	27.9	13.4	7.2	12.4	6.3	4.4	4.3	8.2	7.8	20.0	22.7	28.0	10.6	215	1.1
越前岬	34.1	21.1	12.3	20.1	9.4	12.3	13.2	45.7	34.8	18.8	29.0	30.2	23.1	324	0.0
伊自良湖	62.8	20.0	20.0	17.2	27.6	18.0	26.9	21.7	20.1	18.8	33.5	30.1	21.8	306	3.4
犬山	75.9	14.6	22.3	17.3	22.5	15.8	25.0	42.8	21.3	29.6	27.9	14.7	20.0	270	0.7
潮岬	37.7	10.3	15.4	8.5	9.2	6.9	12.6	30.7	18.9	26.2	17.7	18.6	14.3	178	0.5
京都八幡	66.2	23.9	23.4	21.0	20.3	25.6	13.1	20.5	18.3	26.7	22.6	28.3	23.6	390	3.5
尼崎	76.3	13.2	22.3	20.1	23.3	22.7	22.3	8.5	18.7	22.3	25.1	16.5	22.0	177	3.1
倉橋島	25.9	15.7	17.0	10.0	20.5	14.3	15.7	31.7	22.0	17.8	20.5	27.8	17.1	364	0.0
隠岐	55.1	25.7	6.3	12.7	8.1	31.4	10.5	49.4	53.0	15.5	30.0	42.7	19.3	482	0.4
蟠竜湖	36.7	9.9	15.8	12.6	10.9	14.8	8.8	35.5	139	33.9	60.5	35.4	28.8	251	4.7
栲原	13.0	7.0	13.6	3.3	2.4	3.2	7.1	7.2	18.0	10.7	28.9	14.1	7.8	431	0.0
筑後小郡	67.2	36.8	31.7	11.0	17.0	10.4	11.7	32.7	25.5	20.4	53.1	32.4	23.1	710	3.9
大分久住	16.4	13.5	14.8	4.2	6.7	5.3	3.0	8.5	21.4	6.9	49.4	14.7	8.6	99.6	1.5
対馬	35.9	17.1	20.2	19.5	20.6	34.0	14.0	24.2	67.2	20.5	19.1	35.0	25.9	439	0.0
五島	21.9	19.2	7.7	34.2	*	49.6	3.6	35.8	12.2	23.2	35.5	28.6	24.2	408	0.0
えびの	7.8	17.0	11.1	17.6	5.5	6.0	12.4	1.5	13.1	11.7	15.9	19.4	11.4	606	0.5
屋久島	9.9	11.7	14.5	33.9	6.6	5.2	48.7	36.1	13.4	21.3	46.1	13.4	18.0	103	1.5
辺戸岬	8.5	10.2	3.1	20.2	2.4	3.2	15.4	5.7	6.3	13.4	5.9	17.1	6.3	900	0.2
東京	42.6	19.2	43.1	18.3	77.1	10.9	13.3	23.4	18.1	58.2	21.8	20.7	20.2	580	2.6
最大値 ¹⁾	76.3	47.3	86.6	34.2	77.1	49.6	48.7	49.4	139	99.1	129	84.7	28.8	900	4.7
最小値 ¹⁾	7.8	3.9	3.1	3.3	2.0	2.1	0.3	1.5	6.1	6.6	5.0	4.9	4.4	55.4	0.0
全国平均	35.5	17.9	19.5	16.4	16.7	12.7	14.2	21.4	24.2	25.8	31.1	30.2	17.7		
標準偏差	20.3	9.3	15.6	7.9	14.0	10.7	9.9	13.8	25.1	19.1	23.1	16.4	6.5		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表8(a) 平成19年度 C l⁻濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	63.0	29.0	50.9	84.9	58.9	61.3	374	665	232	528	412	54.2	157	3530	2.7
札幌	56.3	23.6	7.1	5.9	13.3	62.8	131	85.4	38.7	142	185	80.8	81.9	460	3.0
落石岬	121	120	24.3	35.9	17.8	152	295	739	699	2030	158	117	167	4950	6.6
竜飛岬	78.2	233	29.3	236	42.9	262	557	373	267	5090	723	542	277	7480	10.0
八幡平	17.1	15.0	2.6	3.0	7.1	9.2	30.3	27.9	20.6	224	83.0	22.0	20.3	986	0.0
籠岳	23.4	11.1	7.9	8.6	5.8	47.6	8.6	102	25.9	53.9	94.7	47.9	21.7	311	0.8
尾花沢	13.9	4.0	2.3	2.5	6.3	27.0	7.1	142	72.6	88.5	202	14.1	61.0	354	0.9
筑波	28.5	23.7	15.8	11.6	9.1	51.5	13.9	37.7	19.2	16.3	13.0	42.1	24.0	453	0.9
赤城	14.0	4.3	5.9	5.4	3.1	11.8	11.2	14.9	10.3	3.6	5.0	17.1	8.5	281	0.0
小笠原	99.9	41.6	82.8	54.5	83.4	242	161	172	368	133	170	111	146	2790	3.9
佐渡関岬	226	369	23.0	26.9	43.9	93.3	243	983	331	600	1830	162	231	14300	5.3
新潟巻	21.6	65.0	7.4	9.8	26.3	80.5	102	326	306	287	304	106	136	3400	1.0
八方尾根	10.7	3.6	1.3	2.0	2.2	1.9	2.0	9.3	9.9	12.8	19.3	13.2	5.3	313	0.0
越前岬	97.6	87.3	8.5	9.2	20.7	32.2	48.9	359	576	431	489	85.6	213	3440	0.5
伊自良湖	31.4	7.5	4.7	5.2	14.0	14.8	10.6	55.7	43.5	30.6	86.2	25.4	17.2	282	0.9
犬山	63.4	11.1	6.5	5.8	9.0	12.9	13.6	18.0	13.9	8.6	23.9	21.2	10.9	201	1.2
潮岬	240	90.6	82.7	214	53.1	72.0	88.0	66.8	169	108	390	287	148	2180	6.5
京都八幡	34.0	14.9	7.1	5.6	5.1	9.6	12.3	41.9	11.7	11.7	28.4	21.1	12.1	451	1.8
尼崎	41.4	15.6	14.4	8.8	11.6	20.6	16.0	6.6	9.5	7.8	12.0	15.4	14.1	152	1.4
倉橋島	17.9	12.6	14.0	11.6	91.3	12.3	13.2	15.7	21.3	12.9	17.6	22.6	18.5	234	1.8
隠岐	573	124	16.4	31.5	72.2	190	116	1280	1000	762	898	233	293	8830	4.0
蟠竜湖	198	11.5	12.5	59.6	18.1	38.2	27.7	212	488	144	270	119	105	2070	1.5
栲原	11.1	5.8	6.9	2.6	43.4	12.7	7.1	8.4	78.3	22.9	31.4	22.6	17.6	738	0.0
筑後小郡	89.7	32.9	24.1	10.6	10.7	8.8	14.1	25.3	32.1	14.2	41.3	61.0	25.8	941	1.9
大分久住	12.7	6.1	12.7	2.1	2.4	4.2	4.2	4.6	30.8	7.1	32.4	6.6	6.0	98.4	1.2
対馬	41.8	17.7	21.4	23.1	70.1	50.9	36.0	371	61.2	66.0	20.0	45.4	37.5	745	0.0
五島	112	92.1	55.7	74.3	*	65.9	18.8	88.4	131	35.0	151	69.0	78.0	1830	0.0
えびの	9.4	9.7	9.0	28.7	8.6	11.6	25.3	0.9	45.5	18.7	15.4	21.0	14.3	878	0.0
屋久島	77.2	42.1	34.0	159	315	36.1	95.6	248	196	175	228	183	141	728	13.8
辺戸岬	97.8	53.3	39.1	265	62.4	113	179	241	288	416	437	183	164	42900	2.0
東京	37.1	17.4	20.1	11.3	73.0	93.6	15.1	32.1	15.3	18.2	12.4	58.0	41.2	679	0.5
最大値 ¹⁾	573	369	82.8	265	315	262	557	1280	1000	5090	1830	542	293	42900	13.8
最小値 ¹⁾	9.4	3.6	1.3	2.0	2.2	1.9	2.0	0.9	9.5	3.6	5.0	6.6	5.3	98.4	0.0
全国平均	82.5	51.5	21.0	45.6	40.0	61.4	86.4	218	181	371	238	90.6	86.9		
標準偏差	110	77.1	21.3	72.3	58.8	67.6	128	310	238	959	368	110	86.1		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表9(a) 平成19年度 NH₄⁺濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	59.1	23.8	115	33.4	40.0	3.1	14.9	30.3	27.8	10.6	14.8	64.5	25.3	167	0.0
札幌	31.7	22.8	31.2	31.9	30.8	15.7	30.1	19.7	16.6	18.3	28.8	65.8	24.3	181	6.2
落石岬	6.8	10.1	7.8	4.7	10.5	5.4	5.9	1.6	32.9	0.0	6.5	1.9	7.4	198	0.0
竜飛岬	22.1	27.1	33.5	25.2	11.0	9.1	23.6	14.0	8.0	5.7	46.8	54.4	18.3	645	0.0
八幡平	33.9	21.8	12.3	9.2	24.3	4.2	12.2	12.7	11.4	15.4	31.8	50.8	16.3	342	0.0
籠岳	44.7	19.1	25.0	17.5	18.9	4.0	5.6	12.5	12.0	45.2	35.2	26.2	17.2	240	0.7
尾花沢	32.5	11.6	13.7	14.5	22.2	7.4	6.9	15.5	13.6	12.2	22.5	25.9	15.0	129	3.5
筑波	52.5	31.4	30.1	30.5	43.9	12.6	7.8	25.9	17.0	23.9	11.5	43.7	27.7	193	2.9
赤城	44.0	18.4	21.6	37.8	18.3	9.8	13.7	39.5	5.4	9.1	5.0	39.3	19.7	286	0.0
小笠原	16.3	7.5	5.6	26.7	12.3	4.8	3.3	2.0	11.8	4.5	5.6	9.4	7.4	234	0.0
佐渡閨岬	58.8	46.6	16.3	10.8	14.0	11.2	13.1	30.4	24.0	37.6	94.6	34.9	24.5	757	2.3
新潟巻	27.1	28.2	16.2	14.1	14.1	16.2	12.7	28.2	17.3	36.4	35.8	40.9	20.4	289	2.6
八方尾根	37.6	15.0	7.9	15.0	7.7	2.9	2.2	8.3	5.7	15.7	24.0	18.9	10.9	121	0.0
越前岬	43.3	28.8	12.2	24.0	7.7	12.2	9.9	42.7	34.1	23.6	41.5	33.4	25.0	380	0.9
伊自良湖	50.0	28.9	19.5	17.2	25.8	15.1	13.0	15.8	9.9	8.7	20.5	21.7	19.3	208	1.9
犬山	44.8	19.8	25.2	19.5	27.1	15.9	18.8	25.6	15.8	24.9	17.0	14.8	20.4	939	2.3
潮岬	38.4	15.7	9.1	7.3	4.5	6.3	3.9	9.4	11.3	34.6	12.3	16.7	11.8	176	0.3
京都八幡	63.7	24.3	25.1	18.1	14.0	18.6	9.2	19.8	12.7	21.7	16.9	26.1	20.4	396	2.9
尼崎	70.8	15.2	20.7	22.0	15.5	20.5	13.3	7.3	14.0	20.4	18.2	16.9	20.0	232	3.2
倉橋島	17.7	21.6	19.0	11.2	24.4	7.2	10.6	19.9	8.5	9.2	12.2	31.2	15.5	299	0.0
隠岐	36.0	13.0	4.1	13.2	3.1	13.2	7.6	25.2	39.8	12.2	26.9	44.0	13.9	430	0.0
蟠竜湖	34.3	8.2	16.0	14.2	5.7	8.3	6.6	28.6	80.1	21.1	42.6	32.7	21.4	140	1.9
栲原	15.2	7.8	12.0	4.5	4.1	3.1	3.9	7.4	15.7	7.8	23.3	17.1	8.0	544	0.0
筑後小郡	94.3	57.4	53.0	13.5	13.8	10.5	14.5	36.6	37.0	32.6	44.9	35.6	29.6	375	3.8
大分久住	28.5	22.5	25.9	9.5	14.2	9.7	2.9	11.9	17.3	8.3	73.0	23.1	14.8	133	2.9
対馬	49.6	28.4	31.3	27.5	13.9	27.3	17.0	9.9	83.9	24.7	9.6	43.1	32.5	467	0.4
五島	24.9	32.8	11.7	32.9	*	31.7	3.4	20.1	8.6	8.8	22.6	29.1	24.3	150	0.0
えびの	12.9	33.6	15.1	17.9	11.8	14.0	18.8	3.1	17.6	16.3	23.7	25.2	17.8	676	0.4
屋久島	12.9	13.1	15.8	28.3	9.1	3.3	22.7	29.2	11.1	10.1	39.3	13.6	14.6	69.3	1.1
辺戸岬	14.4	11.8	2.8	12.8	3.1	3.7	14.8	5.3	5.6	9.9	8.1	16.0	6.9	640	0.5
東京	53.3	30.0	56.9	27.6	87.9	17.2	15.0	39.8	24.9	89.3	23.8	35.4	29.0	631	3.2
最大値 ¹⁾	94.3	57.4	115	37.8	87.9	31.7	30.1	42.7	83.9	89.3	94.6	65.8	32.5	939	6.2
最小値 ¹⁾	6.8	7.5	2.8	4.5	3.1	2.9	2.2	1.6	5.4	0.0	5.0	1.9	6.9	69.3	0.0
全国平均	37.8	22.5	23.0	19.1	18.5	11.1	11.5	19.3	21.0	20.0	27.1	30.7	18.7		
標準偏差	19.6	11.2	21.1	9.1	16.6	7.1	6.8	11.8	18.7	16.9	19.5	15.4	6.8		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つもない）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表10(a) 平成19年度 Na⁺濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	59.5	26.0	36.5	79.4	55.2	57.2	317	583	214	486	384	51.6	143	3170	2.6
札幌	50.1	19.2	2.9	4.1	7.1	53.3	113	73.5	29.9	126	167	66.7	71.2	396	1.6
落石岬	109	104	20.7	31.7	14.0	124	248	583	621	873	139	101	141	4150	4.8
竜飛岬	71.1	210	26.9	212	37.9	242	486	314	242	4220	641	451	243	6000	3.5
八幡平	14.1	12.8	1.7	1.4	5.6	7.7	26.5	25.3	18.3	185	78.6	21.3	17.4	814	0.2
籠岳	16.8	8.4	6.2	9.0	3.9	39.8	7.7	78.6	22.1	45.2	82.7	41.5	18.1	241	0.0
尾花沢	13.2	2.9	1.2	2.0	5.2	23.2	6.0	124	62.4	73.5	173	11.3	52.1	311	0.3
筑波	22.2	19.9	11.4	8.0	4.6	43.5	10.7	33.9	17.9	9.3	8.3	37.9	19.5	385	0.5
赤城	11.2	4.0	4.7	2.4	1.4	9.3	10.4	13.3	10.4	2.1	2.8	14.3	6.5	301	0.0
小笠原	88.5	37.8	73.7	46.7	70.6	210	138	145	317	111	143	93.5	125	2440	3.8
佐渡関岬	198	319	21.3	26.7	38.9	83.1	230	847	288	510	1620	137	203	12200	3.7
新潟巻	18.4	54.8	5.5	6.9	22.1	67.8	86.9	280	261	248	261	91.1	116	2880	0.5
八方尾根	8.0	2.4	1.0	0.6	1.1	1.4	1.6	7.0	7.6	10.9	16.3	15.1	4.0	357	0.0
越前岬	89.6	77.6	8.1	8.5	20.4	29.2	43.9	325	498	397	426	84.9	189	2910	0.5
伊自良湖	28.4	6.7	3.4	3.5	11.1	12.3	8.7	47.0	36.6	24.6	72.5	22.0	14.2	235	0.9
犬山	61.0	10.3	4.4	3.9	6.8	10.7	12.3	16.4	14.4	7.0	21.4	18.9	9.1	189	0.8
潮岬	216	79.7	73.3	184	47.8	65.2	81.4	63.1	152	96.4	347	257	131	1950	4.1
京都八幡	27.5	12.7	3.6	2.8	3.4	7.6	9.8	36.3	8.6	9.3	23.4	22.3	9.6	395	0.7
尼崎	34.0	13.8	11.5	4.8	7.5	10.6	11.9	4.4	8.9	5.9	9.7	14.3	11.0	129	0.9
倉橋島	17.0	8.7	8.7	8.1	77.3	7.9	9.3	11.3	18.0	8.5	12.6	22.2	14.7	223	0.7
隠岐	510	105	14.9	26.5	63.5	162	99.1	1150	892	677	771	211	258	7540	2.1
蟠竜湖	174	9.7	9.8	53.1	14.7	31.5	22.6	184	438	125	242	108	92.5	1820	0.5
栲原	10.5	5.4	5.7	1.6	36.0	11.1	4.6	8.4	66.2	16.7	29.1	20.2	14.8	802	0.4
筑後小郡	119	25.9	17.3	8.4	9.7	9.9	14.2	18.9	32.1	11.9	47.1	33.2	23.2	734	4.0
大分久住	15.4	7.1	13.8	2.3	2.4	4.5	4.6	3.2	28.0	4.2	31.8	4.9	6.0	79	1.0
対馬	45.3	14.0	17.3	20.2	67.9	48.0	33.3	317	57.0	66.3	16.6	40.8	34.9	635	0.0
五島	95.8	82.3	51.7	70.2	*	61.2	15.6	86.4	113	30.3	142	62.8	70.9	1500	0.0
えびの	9.7	9.4	6.5	26.4	4.3	11.5	22.2	0.8	41.3	13.2	11.0	22.8	12.5	986	0.3
屋久島	61.6	37.3	29.7	135	265	30.6	81.0	197	162	143	148	147	115	613	10.9
辺戸岬	85.3	48.2	34.4	236	55.0	103	165	213	260	373	388	166	146	36900	2.0
東京	30.2	14.3	18.9	7.4	62.4	78.7	16.4	26.3	12.7	10.4	12.4	45.4	34.3	729	0.0
最大値 ¹⁾	510	319	73.7	236	265	242	486	1150	892	4220	1620	451	258	36900	10.9
最小値 ¹⁾	8.0	2.4	1.0	0.6	1.1	1.4	1.6	0.8	7.6	2.1	2.8	4.9	4.0	79	0.0
全国平均	74.5	44.8	17.6	39.8	34.1	53.5	75.4	188	160	288	209	78.6	75.6		
標準偏差	98.1	67.5	19.1	64.2	50.3	60.1	111	271	211	762	325	92.9	75.8		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表11(a) 平成19年度 K⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	3.0	1.5	5.4	2.4	2.0	2.5	8.2	15.0	4.9	8.8	8.2	2.8	4.0	80.2	0.0
札幌	1.9	1.1	1.4	1.1	1.0	1.6	3.4	2.4	1.2	2.3	5.0	3.0	2.3	11.9	0.0
落石岬	2.7	2.7	1.0	0.9	0.9	3.4	6.6	11.8	16.3	20.3	3.8	2.3	3.6	109	0.0
竜飛岬	2.2	5.4	2.8	5.0	1.2	4.9	15.2	6.5	5.3	90.7	15.2	12.4	6.0	130	0.4
八幡平	1.6	1.3	0.5	0.3	1.5	0.1	0.9	1.0	0.8	4.5	2.7	2.5	1.0	18.8	0.0
籠岳	1.4	0.9	0.7	0.4	0.6	3.7	1.7	2.2	0.9	3.5	2.7	1.7	1.4	21.0	0.0
尾花沢	1.6	0.4	0.3	0.4	0.4	0.3	0.3	3.1	1.7	2.0	4.8	0.8	1.5	9.5	0.1
筑波	2.7	1.5	1.4	1.0	0.7	1.7	1.1	1.9	1.6	1.7	0.7	2.1	1.5	17.8	0.1
赤城	3.4	0.9	1.3	1.1	0.4	1.3	0.8	2.8	1.3	0.9	0.5	2.3	1.2	26.3	0.0
小笠原	2.5	1.2	1.8	5.5	1.6	4.3	4.0	2.9	6.5	3.1	3.9	3.1	3.2	54.4	0.2
佐渡関岬	6.6	9.0	1.5	1.2	1.4	2.0	5.0	18.6	7.7	18.4	34.4	4.5	5.5	271	0.0
新潟巻	1.8	2.7	0.6	0.5	0.8	1.7	2.5	7.6	6.5	7.7	7.6	3.4	3.4	63.4	0.0
八方尾根	1.9	1.0	0.4	0.5	0.3	0.1	0.1	0.7	0.6	1.4	1.6	1.5	0.6	12.4	0.0
越前岬	3.5	3.3	1.1	1.0	0.9	1.2	1.8	8.8	12.6	9.8	12.1	3.4	5.4	74.5	0.1
伊自良湖	2.8	1.3	0.9	0.5	0.6	0.5	0.4	1.8	1.3	0.9	2.4	1.1	0.9	16.4	0.1
犬山	7.1	2.1	1.3	0.6	1.0	0.6	0.9	1.6	1.0	1.9	1.3	1.4	1.1	16.9	0.2
潮岬	7.9	2.3	1.7	3.6	1.0	3.0	1.1	1.7	4.2	4.8	8.9	6.1	3.3	73.0	0.1
京都八幡	9.7	2.5	1.5	1.5	1.4	1.1	1.1	1.9	1.2	1.3	1.5	1.8	1.7	42.9	0.6
尼崎	5.1	1.3	2.8	0.7	0.1	0.8	1.1	1.0	0.5	1.2	1.2	2.4	1.5	27.1	0.0
倉橋島	2.0	1.2	1.4	1.0	3.1	1.1	1.3	0.9	1.8	2.2	1.1	1.4	1.4	17.5	0.0
隠岐	17.3	4.8	1.2	1.9	2.8	5.5	3.2	44.0	26.4	15.5	39.7	7.0	9.5	1280	0.3
蟠竜湖	5.8	0.9	0.8	1.8	0.7	0.9	0.9	5.8	14.8	3.7	8.1	3.9	3.2	41.7	0.4
栲原	0.9	0.4	0.6	0.2	1.1	0.4	0.2	0.5	2.0	0.8	1.6	1.1	0.7	25.3	0.0
筑後小郡	14.3	6.1	2.8	2.2	5.1	1.5	1.6	2.9	4.8	2.4	3.6	3.6	3.8	144	0.6
大分久住	1.4	1.5	0.9	0.2	0.3	0.5	0.3	0.9	1.7	0.5	4.8	0.7	0.6	8.6	0.0
対馬	4.5	1.3	6.9	4.7	2.0	1.6	1.2	6.0	3.4	1.7	0.1	2.2	3.2	29.0	0.0
五島	3.3	2.4	1.2	5.4	*	2.9	0.6	24.0	2.8	0.6	4.4	3.7	3.7	107	0.0
えびの	0.9	1.8	1.5	3.4	0.8	1.0	1.5	0.1	1.7	1.5	1.4	1.8	1.4	65.5	0.0
屋久島	1.7	1.2	0.8	7.2	5.7	0.6	2.1	4.7	3.0	2.3	3.9	3.1	2.5	11.8	0.0
辺戸岬	2.7	1.6	1.0	7.1	1.4	2.6	5.2	5.0	5.8	8.5	8.8	5.0	3.6	880	0.1
東京	2.4	1.1	2.9	0.7	4.0	1.7	0.7	1.8	1.0	2.4	1.7	2.3	1.5	22.2	0.0
最大値 ¹⁾	17.3	9.0	6.9	7.2	5.7	5.5	15.2	44.0	26.4	90.7	39.7	12.4	9.5	1280	0.6
最小値 ¹⁾	0.9	0.4	0.3	0.2	0.1	0.1	0.1	0.1	0.5	0.5	0.1	0.7	0.6	8.6	0.0
全国平均	4.1	2.1	1.6	2.1	1.5	1.8	2.4	6.1	4.7	7.3	6.4	3.0	2.7		
標準偏差	3.8	1.9	1.4	2.1	1.4	1.4	3.1	9.0	5.7	16.3	9.0	2.3	2.0		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表12(a) 平成19年度 Ca²⁺濃度 月・年平均値単位: $\mu\text{mol L}^{-1}$

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	6.2	3.8	29.1	3.1	2.3	2.4	12.5	16.6	9.6	12.9	12.9	7.1	5.7	99.4	0.3
札幌	9.4	5.9	2.7	2.5	2.6	2.7	7.1	4.5	2.2	6.4	9.0	10.7	5.2	63.1	1.1
落石岬	12.0	6.4	1.7	1.6	2.2	3.6	6.9	16.4	19.1	34.0	7.9	5.0	5.6	140	0.9
竜飛岬	6.9	21.7	6.3	8.4	1.8	6.8	15.4	10.0	25.2	117	23.9	29.0	11.2	331	0.5
八幡平	9.4	9.6	1.4	1.1	1.5	0.8	2.4	2.4	4.1	7.4	10.7	9.2	3.9	59.3	0.0
籠岳	14.3	10.2	2.4	1.9	1.4	3.4	1.1	3.2	3.6	8.1	9.8	4.2	4.0	355	0.5
尾花沢	9.2	2.7	0.6	0.6	0.9	0.5	0.5	4.0	3.4	3.8	8.7	1.9	3.1	21.4	0.1
筑波	19.3	7.4	3.4	3.1	5.8	3.2	2.3	4.6	11.5	5.5	2.8	12.5	6.3	70.1	0.5
赤城	9.8	3.6	6.5	2.1	1.0	1.2	2.1	7.3	4.2	3.0	2.0	8.6	2.6	115	0.0
小笠原	4.7	2.3	2.9	3.5	2.8	5.3	3.7	4.5	8.9	5.1	5.7	3.3	4.3	99.4	0.7
佐渡関岬	25.0	28.9	3.2	3.0	3.1	4.7	7.6	25.4	10.5	23.2	52.2	14.5	11.0	723	1.1
新潟巻	12.6	13.1	1.3	1.4	1.1	3.1	3.1	8.4	9.0	13.1	14.9	13.6	6.1	125	0.2
八方尾根	17.7	10.1	0.6	0.8	0.4	0.3	0.2	1.4	2.2	5.6	5.1	16.3	3.2	338	0.0
越前岬	13.8	10.8	0.9	1.0	0.8	1.1	1.8	10.3	15.7	10.7	15.0	15.7	8.1	551	0.0
伊自良湖	19.4	7.8	1.5	0.7	1.9	1.6	1.2	2.7	5.2	2.3	6.7	5.0	3.1	130	0.2
犬山	41.0	9.0	2.6	1.1	2.2	1.6	2.0	5.2	7.4	4.3	4.3	3.9	3.3	114	0.2
潮岬	21.7	7.7	2.2	4.6	1.3	1.9	2.2	3.5	8.6	5.1	9.4	10.3	5.3	144	0.4
京都八幡	27.6	12.0	2.9	1.3	2.0	2.6	1.9	5.2	8.4	5.5	6.1	11.8	5.5	296	0.0
尼崎	27.1	7.6	3.7	2.8	3.8	5.4	3.1	2.4	7.5	5.7	5.8	7.4	6.1	119	0.7
倉橋島	10.0	11.5	1.9	1.1	3.9	1.2	3.7	3.2	5.0	2.7	2.2	13.9	4.7	213	0.6
隠岐	44.5	17.9	1.2	1.6	2.1	5.8	3.7	42.2	32.5	18.3	23.7	15.3	10.5	346	0.5
蟠竜湖	30.3	4.0	1.1	2.0	1.1	1.3	1.1	10.5	30.3	5.9	23.2	23.8	8.9	78.0	0.6
栲原	9.7	2.5	0.9	0.3	1.2	0.5	0.4	0.8	7.3	1.7	5.9	6.0	2.2	158	0.0
筑後小郡	57.9	31.9	7.4	2.3	5.5	2.3	1.2	6.8	8.7	4.0	13.4	28.7	11.5	513	0.7
大分久住	12.0	17.1	1.5	0.0	1.0	0.8	1.0	1.5	8.5	1.1	9.4	5.0	3.4	83.7	0.0
対馬	12.9	5.5	1.5	1.4	2.6	3.0	2.5	8.3	7.9	4.9	3.2	10.2	4.6	119	0.0
五島	8.6	8.5	3.0	3.9	*	6.9	1.9	7.8	5.0	2.9	6.9	8.6	5.1	254	0.0
えびの	5.7	9.4	3.6	14.1	0.7	0.8	1.4	0.3	5.4	1.6	3.6	10.1	5.0	397	0.1
屋久島	6.3	8.1	1.3	6.0	5.9	0.8	3.1	6.2	6.7	3.9	10.9	6.9	4.9	39.2	0.5
辺戸岬	5.6	3.0	1.5	9.2	1.8	3.1	5.1	5.6	7.4	10.3	10.9	9.2	4.8	935	0.5
東京	7.8	5.8	11.1	4.0	20.9	3.8	2.3	4.5	6.9	9.5	3.7	8.8	5.3	245	0.4
最大値 ¹⁾	57.9	31.9	29.1	14.1	20.9	6.9	15.4	42.2	32.5	117	52.2	29.0	11.5	935	1.1
最小値 ¹⁾	4.7	2.3	0.6	0.0	0.4	0.3	0.2	0.3	2.2	1.1	2.0	1.9	2.2	21.4	0.0
全国平均	16.7	9.9	3.6	2.9	2.8	2.7	3.4	7.6	9.6	11.1	10.6	10.9	5.6		
標準偏差	12.7	7.1	5.2	3.0	3.7	1.9	3.4	8.3	7.5	20.8	9.8	6.7	2.6		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表13(a) 平成19年度 nss-Ca²⁺濃度 月・年平均値

単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	4.9	3.2	28.3	1.4	1.1	1.1	5.6	4.2	5.0	2.5	4.6	6.0	2.6	32.5	0.0
札幌	8.3	5.5	2.7	2.4	2.4	1.6	4.7	2.9	1.6	3.7	5.4	9.3	3.7	62.3	0.9
落石岬	9.6	4.2	1.2	0.9	1.9	1.0	1.5	3.8	5.7	15.2	4.9	2.8	2.5	97.4	0.0
竜飛岬	5.4	17.2	5.7	3.9	1.0	1.7	4.9	3.2	20.0	25.6	10.1	19.3	5.9	261	0.0
八幡平	9.1	9.3	1.4	1.1	1.4	0.7	1.8	1.9	3.7	3.4	9.0	8.7	3.5	57.1	0.0
籠岳	13.9	10.0	2.3	1.7	1.3	2.6	0.9	1.5	3.1	7.1	8.0	3.3	3.6	354	0.4
尾花沢	9.0	2.7	0.6	0.6	0.7	0.1	0.3	1.4	2.1	2.2	5.0	1.7	2.0	14.7	0.0
筑波	18.8	7.0	3.2	2.9	5.7	2.3	2.1	3.9	11.1	5.3	2.6	11.7	5.8	68.5	0.5
赤城	9.6	3.5	6.4	2.0	1.0	1.0	1.9	7.0	4.0	3.0	2.0	8.3	2.4	114	0.0
小笠原	2.8	1.5	1.3	2.5	1.3	0.7	0.7	1.3	2.2	2.7	2.6	1.3	1.6	60.9	0.0
佐渡関岬	20.7	22.0	2.7	2.4	2.2	2.9	2.7	7.1	4.3	12.2	17.2	11.6	6.6	459	0.5
新潟巻	12.2	11.9	1.2	1.3	0.6	1.6	1.2	2.4	3.4	7.8	9.2	11.6	3.6	107	0.0
八方尾根	17.5	10.1	0.6	0.8	0.3	0.3	0.1	1.2	2.1	5.4	4.7	16.0	3.1	331	0.0
越前岬	11.8	9.1	0.7	0.8	0.4	0.5	0.9	3.2	4.9	2.1	5.8	13.9	4.0	530	0.0
伊自良湖	18.8	7.6	1.4	0.6	1.6	1.4	1.0	1.7	4.4	1.8	5.1	4.6	2.8	128	0.2
犬山	39.7	8.8	2.5	1.0	2.0	1.4	1.8	4.8	7.1	4.1	3.9	3.5	3.1	113	0.1
潮岬	17.0	6.0	0.6	0.6	0.3	0.5	0.5	2.2	5.3	3.0	1.9	4.8	2.5	131	0.0
京都八幡	27.0	11.7	2.8	1.2	2.0	2.4	1.7	4.4	8.2	5.3	5.6	11.3	5.3	287	0.0
尼崎	26.4	7.3	3.4	2.7	3.6	5.1	2.9	2.3	7.3	5.5	5.6	7.1	5.9	118	0.7
倉橋島	9.7	11.4	1.7	0.9	2.2	1.0	3.5	3.0	4.6	2.5	1.9	13.4	4.4	208	0.4
隠岐	33.5	15.6	0.8	1.0	0.7	2.3	1.6	17.3	13.2	3.7	7.0	10.8	5.0	303	0.0
蟠竜湖	26.6	3.8	0.9	0.9	0.8	0.6	0.6	6.5	20.8	3.2	18.0	21.5	6.9	65.0	0.3
栲原	9.5	2.4	0.8	0.3	0.4	0.3	0.3	0.6	5.9	1.4	5.3	5.6	1.9	153	0.0
筑後小郡	55.3	31.3	7.0	2.1	5.3	2.1	0.9	6.4	8.0	3.7	12.4	28.0	11.0	500	0.1
大分久住	11.6	17.0	1.2	0.0	0.9	0.7	0.9	1.4	7.9	1.0	8.7	4.9	3.2	83.2	0.0
対馬	11.9	5.2	1.3	1.1	1.2	1.9	1.8	1.5	6.7	3.4	2.9	9.3	3.9	116	0.0
五島	6.6	6.7	1.9	2.4	*	5.6	1.6	5.9	2.6	2.2	3.8	7.2	3.6	225	0.0
えびの	5.5	9.2	3.4	13.5	0.7	0.6	0.9	0.3	4.5	1.4	3.4	9.6	4.7	379	0.1
屋久島	4.9	7.3	0.7	3.1	0.2	0.1	1.3	2.0	3.2	0.8	7.7	3.7	2.5	37.7	0.0
辺戸岬	3.8	1.9	0.7	4.2	0.6	0.9	1.5	1.0	1.8	2.3	2.5	5.6	1.7	159	0.1
東京	7.2	5.4	10.7	3.8	19.5	2.1	1.9	3.9	6.6	9.2	3.4	7.9	4.6	242	0.4
最大値 ¹⁾	<u>55.3</u>	<u>31.3</u>	<u>28.3</u>	<u>13.5</u>	19.5	<u>5.6</u>	<u>5.6</u>	17.3	<u>20.8</u>	<u>25.6</u>	18.0	28.0	11.0	530	0.9
最小値 ¹⁾	2.8	1.5	0.6	<u>0.0</u>	<u>0.2</u>	<u>0.1</u>	0.1	0.3	1.6	<u>0.8</u>	1.9	1.3	1.6	14.7	0.0
全国平均	15.1	8.9	3.2	2.1	2.1	1.5	1.7	3.6	6.2	4.9	6.1	9.2	4.0		
標準偏差	11.7	6.4	5.2	2.4	3.5	1.3	1.3	3.2	4.6	5.0	4.0	6.0	2.0		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表14(a) 平成19年度 Mg²⁺濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	7.2	3.3	13.0	9.3	6.5	7.3	37.4	70.1	26.8	59.2	44.5	6.9	17.3	372	0.3
札幌	6.0	2.6	0.8	1.0	1.0	6.4	13.5	8.4	3.3	14.1	19.2	8.6	8.3	44.6	0.3
落石岬	13.6	12.9	2.7	3.7	2.2	14.8	27.4	65.9	71.0	100	16.3	12.4	16.5	499	0.8
竜飛岬	8.5	25.3	4.0	23.1	4.3	25.7	54.6	36.5	29.0	512	72.0	53.3	27.8	729	0.4
八幡平	3.0	3.1	0.6	0.5	1.2	0.9	3.3	3.1	2.3	21.9	10.1	4.1	2.6	92.5	0.0
籠岳	5.3	2.5	1.1	0.8	0.6	5.0	1.0	9.7	2.8	6.4	9.7	5.1	2.6	40.7	0.0
尾花沢	3.0	0.7	0.2	0.3	0.6	2.1	0.5	13.5	7.1	8.5	20.0	1.6	5.9	36.2	0.0
筑波	4.8	2.8	1.4	1.4	1.5	5.9	1.8	4.5	2.4	2.1	1.3	6.8	3.1	52.9	0.0
赤城	3.6	1.1	1.7	0.7	0.8	1.2	1.7	3.4	1.5	1.4	0.7	3.0	1.3	102	0.0
小笠原	10.1	4.2	8.5	5.9	7.9	22.2	15.1	16.4	36.1	13.2	15.3	10.3	13.9	269	1.0
佐渡関岬	26.2	38.6	2.5	2.9	4.4	9.4	25.3	94.0	33.9	59.9	181	17.0	23.5	1450	0.0
新潟巻	4.3	8.4	0.8	1.3	2.5	8.0	9.7	31.5	30.3	29.3	30.7	11.6	13.6	326	0.1
八方尾根	2.7	1.5	0.1	0.1	0.1	0.1	0.1	0.8	1.1	1.8	2.6	3.4	0.8	36.1	0.0
越前岬	11.5	9.8	1.0	1.0	2.2	3.3	4.8	38.1	57.3	45.2	49.5	10.3	21.9	338	0.0
伊自良湖	6.4	1.8	0.5	0.4	1.4	1.5	1.1	5.4	4.6	2.9	8.6	2.9	1.9	27.4	0.2
犬山	12.5	2.1	0.8	0.6	0.9	1.3	1.6	2.1	2.3	1.4	2.7	2.6	1.4	26.4	0.0
潮岬	28.8	9.7	7.7	20.6	5.3	7.1	9.1	7.3	17.2	11.3	39.4	29.2	14.9	219	0.3
京都八幡	6.6	2.9	0.6	0.4	0.6	1.0	1.6	4.1	2.2	1.7	3.4	3.4	1.7	35.1	0.0
尼崎	8.5	2.4	2.0	0.9	1.8	2.0	1.7	1.2	1.3	1.1	1.4	2.4	2.0	24.3	0.2
倉橋島	3.6	3.0	1.6	1.3	9.3	1.3	1.5	1.9	2.8	1.7	2.0	3.9	2.5	40.5	0.3
隠岐	61.2	14.5	1.9	3.0	6.3	19.2	11.5	130	101	75.6	88.8	26.1	29.5	850	0.5
蟠竜湖	24.2	1.8	1.2	6.0	1.7	3.6	2.6	21.7	53.4	14.9	29.0	14.6	11.4	206	0.4
栲原	2.5	1.0	0.8	0.1	4.2	1.5	0.7	0.8	8.5	2.1	3.8	2.7	1.9	102	0.0
筑後小郡	19.5	5.6	2.7	1.7	1.7	1.3	1.9	9.5	8.6	2.8	7.8	6.6	4.4	176	0.6
大分久住	5.3	3.4	1.5	0.0	0.4	0.5	0.7	0.7	3.8	0.6	5.1	1.3	1.2	12.3	0.0
対馬	7.4	3.1	1.7	2.7	7.2	6.1	3.4	34.1	9.6	7.7	2.0	6.0	4.7	72.9	0.0
五島	11.4	10.4	5.4	6.9	*	9.0	4.0	10.1	12.7	3.8	15.0	8.7	8.0	179	0.0
えびの	2.7	2.5	1.1	2.2	0.6	1.5	2.7	0.1	5.3	1.9	1.9	4.4	2.1	143	0.1
屋久島	8.0	5.4	3.7	17.3	29.7	3.5	9.0	23.3	19.3	16.6	19.2	18.0	13.6	70.4	1.2
辺戸岬	10.0	5.8	4.0	28.3	6.3	12.0	19.5	24.2	29.5	42.5	43.8	19.4	16.7	4050	0.2
東京	4.0	2.4	3.1	1.2	9.5	9.0	1.6	3.3	1.9	2.4	1.4	6.1	4.2	63.9	0.0
最大値 ¹⁾	61.2	38.6	13.0	28.3	29.7	25.7	54.6	130	101	512	181	53.3	29.5	4050	1.2
最小値 ¹⁾	2.5	0.7	0.1	0.0	0.1	0.1	0.1	0.1	1.1	0.6	0.7	1.3	0.8	12.3	0.0
全国平均	10.7	6.3	2.5	4.7	4.1	6.2	8.7	21.8	19.0	34.4	24.1	10.1	9.1		
標準偏差	11.7	7.9	2.8	7.4	5.6	6.6	12.5	30.4	24.0	92.2	36.4	10.7	8.5		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表15(a) 平成19年度 H⁺濃度 月・年平均値単位：μmol L⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	62.0	21.5	77.6	27.0	44.6	7.3	7.8	20.4	22.7	19.1	23.0	59.5	26.0	151	0.6
札幌	25.7	24.4	47.5	28.5	44.2	17.6	31.5	16.6	25.0	18.7	32.3	52.8	27.1	123	8.7
落石岬	12.3	18.0	14.2	11.1	27.6	14.8	14.2	21.0	49.9	6.1	11.2	8.0	16.2	83.2	1.6
竜飛岬	22.8	19.7	50.9	38.4	22.6	17.1	37.8	18.8	16.4	20.0	66.9	58.2	26.0	195	0.7
八幡平	31.1	14.3	11.9	12.7	19.6	5.8	15.8	17.9	15.3	16.4	25.4	37.9	15.6	214	0.3
籠岳	42.7	16.0	30.3	23.3	19.4	5.9	6.3	9.1	13.5	83.5	16.2	46.1	20.1	219	1.5
尾花沢	26.2	13.2	17.3	14.2	29.4	12.1	12.3	25.9	22.0	18.4	15.2	31.9	19.0	63.1	4.0
筑波	26.5	21.0	16.6	24.3	20.2	9.8	10.2	28.6	11.0	43.1	17.0	21.0	19.3	180	0.2
赤城	14.6	12.8	15.5	18.4	15.1	11.5	18.8	42.9	9.7	17.7	6.6	24.8	14.8	102	0.1
小笠原	9.5	10.5	10.0	16.0	4.4	5.3	2.8	8.1	13.7	17.1	15.4	13.9	10.3	107	0.2
佐渡関岬	48.3	40.6	23.4	12.8	22.2	20.8	26.1	47.9	47.1	49.0	10.0	44.5	31.0	174	0.3
新潟巻	23.2	32.8	22.1	20.8	24.0	38.0	23.8	57.7	43.1	54.0	33.2	57.6	33.3	263	0.1
八方尾根	20.4	10.3	13.6	21.2	10.2	6.1	9.5	24.5	24.1	34.2	30.4	25.9	16.7	145	0.1
越前岬	34.5	25.5	22.4	39.1	15.8	20.1	18.3	47.6	52.4	36.5	34.2	32.2	32.8	245	0.0
伊自良湖	31.9	19.7	28.5	26.7	36.3	19.5	38.3	42.4	30.5	29.0	40.6	38.8	28.8	100	3.2
犬山	23.6	11.2	27.3	23.3	20.2	12.4	29.9	57.4	22.3	50.6	31.2	19.4	22.7	295	0.3
潮岬	34.3	18.0	35.2	21.5	27.3	13.4	22.4	43.2	27.6	79.2	33.1	27.9	28.7	219	0.6
京都八幡	23.7	13.9	31.2	28.8	26.3	33.9	21.4	25.8	19.0	30.0	18.2	18.4	25.2	229	0.6
尼崎	45.2	11.8	25.0	28.5	31.6	24.6	33.5	8.1	21.6	25.1	13.5	13.1	23.3	245	1.0
倉橋島	25.8	23.6	38.8	22.4	46.7	31.5	23.6	26.3	26.0	33.4	30.7	30.4	27.9	191	0.1
隠岐	23.0	19.5	7.7	20.8	10.1	43.7	12.3	28.9	53.8	26.5	27.5	35.9	20.3	198	0.1
蟠竜湖	14.6	10.7	21.4	18.8	15.3	24.6	14.6	46.0	121	51.0	38.5	21.8	29.6	201	1.7
栲原	16.4	13.3	24.4	9.6	8.1	9.3	14.1	13.7	42.1	29.3	30.8	30.9	16.6	380	0.3
筑後小郡	18.7	14.6	25.2	12.0	6.7	13.4	9.1	25.3	34.8	19.0	18.1	12.6	15.1	148	0.3
大分久住	16.7	12.2	18.4	14.1	13.0	13.9	8.7	28.5	29.0	26.0	129	19.2	16.2	129	0.6
対馬	29.8	29.0	26.3	26.5	28.0	26.3	17.5	29.0	74.0	37.0	22.2	26.7	30.8	295	0.0
五島	27.0	29.2	16.1	22.9	*	42.7	4.9	43.2	19.0	31.1	37.7	52.8	25.3	219	0.0
えびの	12.7	26.4	18.4	25.4	20.2	8.9	16.9	1.9	24.4	47.8	39.9	18.3	20.5	302	0.1
屋久島	26.6	15.3	28.7	33.9	15.2	14.4	82.3	64.0	35.7	55.4	143	43.5	38.3	257	2.6
辺戸岬	11.4	20.9	5.6	21.1	5.4	5.9	21.4	8.4	9.1	27.4	12.8	13.2	10.4	447	0.3
東京	32.4	22.1	18.0	20.4	33.4	9.7	8.8	30.2	10.1	23.0	10.0	23.9	16.9	331	0.1
最大値 ¹⁾	62.0	40.6	77.6	39.1	46.7	43.7	82.3	64.0	121	83.5	143	59.5	38.3	447	8.7
最小値 ¹⁾	9.5	10.3	5.6	9.6	4.4	5.3	2.8	1.9	9.1	6.1	6.6	8.0	10.3	63.1	0.0
全国平均	26.2	19.1	24.8	22.1	22.1	17.4	19.8	29.3	31.2	34.0	32.7	31.0	22.7		
標準偏差	11.7	7.3	14.3	7.4	11.4	11.0	15.0	16.2	22.7	17.9	30.2	14.9	7.1		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表16(a) 平成19年度 pH 月・年平均値

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	4.21	4.67	4.11	4.57	4.35	5.14	5.11	4.69	4.64	4.72	4.64	4.23	4.59	6.23	3.82
札幌	4.59	4.61	4.32	4.54	4.35	4.75	4.50	4.78	4.60	4.73	4.49	4.28	4.57	5.06	3.91
落石岬	4.91	4.74	4.85	4.95	4.56	4.83	4.85	4.68	4.30	5.22	4.95	5.10	4.79	5.80	4.08
竜飛岬	4.64	4.71	4.29	4.42	4.65	4.77	4.42	4.73	4.78	4.70	4.17	4.23	4.58	6.18	3.71
八幡平	4.51	4.84	4.93	4.90	4.71	5.24	4.80	4.75	4.81	4.79	4.60	4.42	4.81	6.47	3.67
籠岳	4.37	4.80	4.52	4.63	4.71	5.23	5.20	5.04	4.87	4.08	4.79	4.34	4.70	5.82	3.66
尾花沢	4.58	4.88	4.76	4.85	4.53	4.92	4.91	4.59	4.66	4.74	4.82	4.50	4.72	5.40	4.20
筑波	4.58	4.68	4.78	4.61	4.69	5.01	4.99	4.54	4.96	4.37	4.77	4.68	4.71	6.70	3.74
赤城	4.84	4.89	4.81	4.74	4.82	4.94	4.73	4.37	5.01	4.75	5.18	4.61	4.83	6.87	3.99
小笠原	5.02	4.98	5.00	4.80	5.36	5.28	5.55	5.09	4.86	4.77	4.81	4.86	4.99	6.81	3.97
佐渡関岬	4.32	4.39	4.63	4.89	4.65	4.68	4.58	4.32	4.33	4.31	5.00	4.35	4.51	6.48	3.76
新潟巻	4.64	4.48	4.66	4.68	4.62	4.42	4.62	4.24	4.37	4.27	4.48	4.24	4.48	6.92	3.58
八方尾根	4.69	4.99	4.87	4.67	4.99	5.21	5.02	4.61	4.62	4.47	4.52	4.59	4.78	6.83	3.84
越前岬	4.46	4.59	4.65	4.41	4.80	4.70	4.74	4.32	4.28	4.44	4.47	4.49	4.48	7.93	3.61
伊自良湖	4.50	4.71	4.55	4.57	4.44	4.71	4.42	4.37	4.52	4.54	4.39	4.41	4.54	5.50	4.00
犬山	4.63	4.95	4.56	4.63	4.69	4.91	4.52	4.24	4.65	4.30	4.51	4.71	4.64	6.52	3.53
潮岬	4.47	4.75	4.45	4.67	4.56	4.87	4.65	4.36	4.56	4.10	4.48	4.55	4.54	6.23	3.66
京都八幡	4.63	4.86	4.51	4.54	4.58	4.47	4.67	4.59	4.72	4.52	4.74	4.74	4.60	6.23	3.64
尼崎	4.34	4.93	4.60	4.55	4.50	4.61	4.48	5.09	4.67	4.60	4.87	4.88	4.63	5.98	3.61
倉橋島	4.59	4.63	4.41	4.65	4.33	4.50	4.63	4.58	4.59	4.48	4.51	4.52	4.55	6.88	3.72
隠岐	4.64	4.71	5.11	4.68	4.99	4.36	4.91	4.54	4.27	4.58	4.56	4.44	4.69	7.07	3.70
蟠竜湖	4.83	4.97	4.67	4.73	4.81	4.61	4.84	4.34	3.92	4.29	4.41	4.66	4.53	5.76	3.70
栲原	4.79	4.88	4.61	5.02	5.09	5.03	4.85	4.86	4.38	4.53	4.51	4.51	4.78	6.54	3.42
筑後小郡	4.73	4.84	4.60	4.92	5.17	4.87	5.04	4.60	4.46	4.72	4.74	4.90	4.82	6.59	3.83
大分久住	4.78	4.91	4.74	4.85	4.89	4.86	5.06	4.55	4.54	4.59	3.89	4.72	4.79	6.25	3.89
対馬	4.53	4.54	4.58	4.58	4.55	4.58	4.76	4.54	4.13	4.43	4.65	4.57	4.51	8.18	3.53
五島	4.57	4.53	4.79	4.64	*	4.37	5.31	4.37	4.72	4.51	4.42	4.28	4.60	11.0	3.66
えびの	4.90	4.58	4.74	4.60	4.69	5.05	4.77	5.71	4.61	4.32	4.40	4.74	4.69	7.02	3.52
屋久島	4.57	4.81	4.54	4.47	4.82	4.84	4.08	4.19	4.45	4.26	3.85	4.36	4.42	5.59	3.59
辺戸岬	4.94	4.68	5.25	4.68	5.27	5.23	4.67	5.07	5.04	4.56	4.89	4.88	4.98	6.46	3.35
東京	4.49	4.65	4.74	4.69	4.48	5.02	5.05	4.52	5.00	4.64	5.00	4.62	4.77	6.95	3.48
最大値 ¹⁾	5.02	4.99	5.25	5.02	5.36	5.28	5.55	5.71	5.04	5.22	5.18	5.10	4.99	11.0	4.20
最小値 ¹⁾	4.21	4.39	4.11	4.41	4.33	4.36	4.08	4.19	3.92	4.08	3.85	4.23	4.42	5.06	3.35
全国平均	4.62	4.75	4.67	4.68	4.72	4.84	4.80	4.62	4.59	4.53	4.60	4.56	4.67		
標準偏差	0.19	0.16	0.23	0.15	0.26	0.27	0.30	0.33	0.27	0.23	0.30	0.22	0.14		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。

最大値²⁾、最小値²⁾：サンプルの最大値、最小値。

(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。

表17(a) 平成19年度 電気伝導率 月・年平均値

単位：mS m⁻¹

	4月	5月	6月	7月	8月	9月	10月	11月	12月	1月	2月	3月	年平均値	最大値 ²⁾	最小値 ²⁾
利尻	4.72	1.81	6.45	2.90	3.13	1.26	5.40	10.3	5.03	8.75	7.24	4.86	3.74	50.5	0.28
札幌	2.54	1.87	2.49	1.72	2.41	1.68	3.66	2.24	1.79	3.35	4.47	5.04	2.71	9.28	0.47
落石岬	2.53	2.67	1.20	1.15	1.77	2.84	4.85	10.3	12.6	14.9	2.61	2.16	3.17	63.6	0.61
竜飛岬	2.66	4.74	3.10	5.26	1.86	4.39	9.78	6.02	4.99	74.3	12.5	10.6	5.04	74.3	0.94
八幡平	2.31	1.56	0.80	0.83	1.46	0.51	1.41	1.62	1.38	4.34	2.89	4.75	1.42	18.4	0.15
籠岳	3.46	1.38	1.88	1.49	1.32	1.14	0.53	2.03	1.27	5.47	2.90	3.32	1.59	16.1	0.28
尾花沢	2.20	0.94	0.97	0.81	1.61	1.04	0.77	3.41	2.49	2.61	4.45	2.27	2.08	7.84	0.41
筑波	2.68	1.88	1.43	1.70	1.91	1.41	0.83	2.45	1.42	2.58	1.12	2.63	1.75	13.7	0.18
赤城	1.97	1.05	1.35	1.45	0.89	0.88	1.38	3.24	0.93	1.17	0.62	2.09	1.16	12.8	0.18
小笠原	2.33	1.16	1.63	1.96	1.55	3.43	2.38	2.72	5.58	2.42	3.05	2.22	2.52	24.9	0.44
佐渡関岬	4.78	4.42	1.78	1.22	2.00	2.41	4.13	14.5	6.80	9.15	8.75	4.41	4.10	79.6	0.59
新潟巻	2.11	3.07	1.39	1.28	1.68	3.01	2.66	7.36	6.37	6.94	5.96	4.71	3.68	40.6	0.28
八方尾根	2.19	1.01	0.88	1.40	0.72	0.40	0.46	1.40	1.28	2.05	2.14	2.14	1.12	13.1	0.18
越前岬	3.83	3.03	1.30	2.24	1.17	1.50	1.74	7.91	10.8	8.23	9.17	3.57	4.91	50.0	0.13
伊自良湖	3.28	1.66	1.68	1.51	2.27	1.34	2.14	2.97	2.25	1.86	3.42	2.55	1.90	10.1	0.34
犬山	3.96	1.27	1.92	1.50	1.65	1.07	1.95	3.19	1.74	3.08	2.06	1.43	1.64	18.0	0.26
潮岬	6.39	2.59	3.04	4.04	2.31	1.81	2.59	3.34	4.05	5.91	7.31	5.87	3.74	39.0	0.57
京都八幡	3.07	1.56	1.90	1.60	1.52	2.03	1.29	1.71	1.38	1.96	1.30	1.78	1.70	15.3	0.26
尼崎	4.29	1.22	1.87	1.83	1.91	1.65	1.98	0.74	1.64	1.70	1.30	1.33	1.74	14.4	0.31
倉橋島	2.18	2.06	2.50	1.51	4.02	1.90	1.47	1.60	2.04	2.08	2.13	2.70	2.05	15.7	0.28
隠岐	11.0	3.47	0.81	1.88	1.73	5.22	2.51	18.9	16.9	12.7	15.0	6.48	5.56	112	0.31
蟠竜湖	4.80	1.01	1.53	2.05	1.16	1.91	1.27	5.81	14.3	4.86	6.78	3.90	3.45	31.5	0.52
栲原	1.37	0.92	1.45	0.56	1.02	0.60	0.77	0.85	2.95	1.63	2.16	1.90	1.10	24.3	0.16
筑後小郡	5.30	2.79	2.17	0.99	0.89	1.02	1.06	2.31	2.96	1.70	2.65	2.41	1.82	26.8	0.22
大分久住	1.72	1.37	1.38	0.78	0.86	0.87	0.58	1.64	2.39	1.42	8.39	1.43	1.12	8.39	0.43
対馬	3.36	2.33	1.96	1.98	2.48	2.85	1.55	6.67	5.88	3.25	1.57	2.80	2.65	17.0	0.30
五島	3.34	3.50	1.76	2.36	*	3.36	0.62	4.37	3.00	2.27	4.46	4.12	2.65	26.6	0.25
えびの	1.04	2.13	1.28	2.20	1.24	0.78	1.51	0.17	2.23	2.51	2.30	1.78	1.52	19.8	0.11
屋久島	2.52	1.75	2.00	4.54	5.12	1.27	5.26	6.52	4.40	4.72	8.95	4.60	3.82	15.0	0.87
辺戸岬	2.03	1.78	0.87	4.93	1.16	2.07	4.01	3.67	4.36	7.52	7.02	3.74	2.87	340	0.17
東京	2.97	1.74	2.05	1.60	4.17	2.02	0.87	2.45	1.32	2.63	1.13	2.27	1.83	18.2	0.32
最大値 ¹⁾	11.0	4.74	6.45	5.26	5.12	5.22	9.78	18.9	16.9	74.3	15.0	10.6	5.56	340	0.94
最小値 ¹⁾	1.04	0.92	0.80	0.56	0.72	0.40	0.46	0.17	0.93	1.17	0.62	1.33	1.10	7.84	0.11
全国平均	3.32	2.06	1.83	1.98	1.90	1.86	2.30	4.59	4.40	6.71	4.71	3.41	2.59		
標準偏差	1.87	1.00	1.04	1.19	1.03	1.13	1.98	4.21	4.06	13.0	3.59	1.92	1.24		

網掛けの数値：参考値（月間値または年間値が有効判定基準で棄却されたもの）を表す。

「*」：欠測（該当する月間に測定データが1つも無い）を表す。

最大値¹⁾、最小値¹⁾：月・年平均値の最大値、最小値。最大値²⁾、最小値²⁾：サンプルの最大値、最小値。(注)最大値¹⁾、最小値¹⁾、全国平均、標準偏差は参考値(下線で示す)を含む。