

Survey point's No.	Names of species	Specimen collection site	Gender (M : Male, F : Female)	Age (A : Adult AL : In lactation Pr : Pregnant S : Sub-adult J : Juvenile N : Newly born y : Years old	Specimen Lipid (%)	No.	SPEED'98 No.	object substance-1	object substance-2	data	unit	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	16	-	-	Monobutyltin	<500	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	16	-	-	Monobutyltin	<200	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	16	-	-	Monobutyltin	<500	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	16	-	-	Monobutyltin	<200	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	16	-	-	Monobutyltin	<200	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	16	-	-	Monobutyltin	<200	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	16	-	-	Monobutyltin	<200	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	16	-	-	Monobutyltin	<200	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	16	-	-	Monobutyltin	<200	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	16	-	-	Monobutyltin	<500	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	16	-	-	Monobutyltin	<200	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	16	-	-	Monobutyltin	<200	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	16	-	-	Monobutyltin	<200	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	16	-	-	Monobutyltin	300	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	16	-	-	Monobutyltin	<200	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	16	-	-	Monobutyltin	<200	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	16	-	-	Monobutyltin	<200	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	16	-	-	Monobutyltin	<200	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	16	-	-	Monobutyltin	<200	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	16	-	-	Monobutyltin	<200	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	16	-	-	Monobutyltin	<200	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	16	-	-	Monobutyltin	200	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	16	-	-	Monobutyltin	<200	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	16	-	-	Monobutyltin	<200	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	16	-	-	Monobutyltin	<200	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	16	-	-	Monobutyltin	<200	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	15	-	-	Dibutyltin	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	15	-	-	Dibutyltin	<100	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	15	-	-	Dibutyltin	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	15	-	-	Dibutyltin	300	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	15	-	-	Dibutyltin	200	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	15	-	-	Dibutyltin	100	ug/kg-wet

7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	15	-	-	Dibutyltin	<100	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	15	-	-	Dibutyltin	400	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	15	-	-	Dibutyltin	200	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	15	-	-	Dibutyltin	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	15	-	-	Dibutyltin	400	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	15	-	-	Dibutyltin	200	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	15	-	-	Dibutyltin	<100	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	15	-	-	Dibutyltin	800	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	15	-	-	Dibutyltin	<100	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	15	-	-	Dibutyltin	<100	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	15	-	-	Dibutyltin	100	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	15	-	-	Dibutyltin	200	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	15	-	-	Dibutyltin	200	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	15	-	-	Dibutyltin	600	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	15	-	-	Dibutyltin	<100	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	15	-	-	Dibutyltin	200	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	15	-	-	Dibutyltin	100	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	15	-	-	Dibutyltin	1,100	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	15	-	-	Dibutyltin	400	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	15	-	-	Dibutyltin	<100	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	14	34	-	Triphenyltin	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	14	34	-	Triphenyltin	20	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	14	34	-	Triphenyltin	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	14	34	-	Triphenyltin	60	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	14	34	-	Triphenyltin	40	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	14	34	-	Triphenyltin	40	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	14	34	-	Triphenyltin	<20	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	14	34	-	Triphenyltin	30	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	14	34	-	Triphenyltin	<20	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	14	34	-	Triphenyltin	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	14	34	-	Triphenyltin	40	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	14	34	-	Triphenyltin	50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	14	34	-	Triphenyltin	<20	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	14	34	-	Triphenyltin	40	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	14	34	-	Triphenyltin	<20	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	14	34	-	Triphenyltin	<20	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	14	34	-	Triphenyltin	20	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	14	34	-	Triphenyltin	<20	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	14	34	-	Triphenyltin	<20	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	14	34	-	Triphenyltin	20	ug/kg-wet

21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	14	34	-	Triphenyltin	<20	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	14	34	-	Triphenyltin	20	ug/kg-wet	
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	14	34	-	Triphenyltin	<20	ug/kg-wet	
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	14	34	-	Triphenyltin	50	ug/kg-wet	
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	14	34	-	Triphenyltin	<20	ug/kg-wet	
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	14	34	-	Triphenyltin	<20	ug/kg-wet	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	13	33	-	Tributyltin	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	13	33	-	Tributyltin	<20	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	13	33	-	Tributyltin	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	13	33	-	Tributyltin	60	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	13	33	-	Tributyltin	40	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	13	33	-	Tributyltin	30	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	13	33	-	Tributyltin	<20	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	13	33	-	Tributyltin	40	ug/kg-wet	
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	13	33	-	Tributyltin	<20	ug/kg-wet	
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	13	33	-	Tributyltin	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	13	33	-	Tributyltin	50	ug/kg-wet	
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	13	33	-	Tributyltin	90	ug/kg-wet	
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	13	33	-	Tributyltin	<20	ug/kg-wet	
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	13	33	-	Tributyltin	330	ug/kg-wet	
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	13	33	-	Tributyltin	<20	ug/kg-wet	
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	13	33	-	Tributyltin	30	ug/kg-wet	
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	13	33	-	Tributyltin	30	ug/kg-wet	
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	13	33	-	Tributyltin	40	ug/kg-wet	
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	13	33	-	Tributyltin	50	ug/kg-wet	
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	13	33	-	Tributyltin	110	ug/kg-wet	
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	13	33	-	Tributyltin	30	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	13	33	-	Tributyltin	150	ug/kg-wet	
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	13	33	-	Tributyltin	40	ug/kg-wet	
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	13	33	-	Tributyltin	130	ug/kg-wet	
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	13	33	-	Tributyltin	140	ug/kg-wet	
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	13	33	-	Tributyltin	40	ug/kg-wet	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet	

9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	12	43	-	Benzo(a)pyrene	<5	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	11	26	Heptachlir epoxide	<10	ug/kg-wet	
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	11	26	Heptachlir epoxide	60	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	11	26	Heptachlir epoxide	20	ug/kg-wet	
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	11	26	Heptachlir epoxide	110	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	11	26	Heptachlir epoxide	50	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	11	26	Heptachlir epoxide	50	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	11	26	Heptachlir epoxide	20	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	11	26	Heptachlir epoxide	40	ug/kg-wet	
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	11	26	Heptachlir epoxide	30	ug/kg-wet	
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	11	26	Heptachlir epoxide	20	ug/kg-wet	
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	11	26	Heptachlir epoxide	30	ug/kg-wet	
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	11	26	Heptachlir epoxide	30	ug/kg-wet	
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	11	26	Heptachlir epoxide	30	ug/kg-wet	
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	11	26	Heptachlir epoxide	<10	ug/kg-wet	
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	11	26	Heptachlir epoxide	20	ug/kg-wet	
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	11	26	Heptachlir epoxide	60	ug/kg-wet	
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	11	26	Heptachlir epoxide	10	ug/kg-wet	
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	11	26	Heptachlir epoxide	40	ug/kg-wet	
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	11	26	Heptachlir epoxide	200	ug/kg-wet	
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	11	26	Heptachlir epoxide	10	ug/kg-wet	
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	11	26	Heptachlir epoxide	70	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	11	26	Heptachlir epoxide	50	ug/kg-wet	

23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	11	26	Heptachlir epoxide	20	ug/kg-wet		
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	11	26	Heptachlir epoxide	220	ug/kg-wet		
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	11	26	Heptachlir epoxide	30	ug/kg-wet		
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	11	26	Heptachlir epoxide	<10	ug/kg-wet		
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	10	25	-	Heptachlor	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	10	25	-	Heptachlor	<5	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	10	25	-	Heptachlor	<5	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	10	25	-	Heptachlor	<5	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	10	25	-	Heptachlor	<5	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	10	25	-	Heptachlor	<5	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	10	25	-	Heptachlor	<5	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	10	25	-	Heptachlor	<5	ug/kg-wet	
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	10	25	-	Heptachlor	<5	ug/kg-wet	
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	10	25	-	Heptachlor	<5	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	10	25	-	Heptachlor	<5	ug/kg-wet	
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	10	25	-	Heptachlor	<5	ug/kg-wet	
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	10	25	-	Heptachlor	<5	ug/kg-wet	
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	10	25	-	Heptachlor	<5	ug/kg-wet	
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	10	25	-	Heptachlor	<5	ug/kg-wet	
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	10	25	-	Heptachlor	<5	ug/kg-wet	
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	10	25	-	Heptachlor	<5	ug/kg-wet	
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	10	25	-	Heptachlor	<5	ug/kg-wet	
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	10	25	-	Heptachlor	<5	ug/kg-wet	
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	10	25	-	Heptachlor	<5	ug/kg-wet	
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	10	25	-	Heptachlor	<5	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	10	25	-	Heptachlor	<5	ug/kg-wet	
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	10	25	-	Heptachlor	<5	ug/kg-wet	
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	10	25	-	Heptachlor	<5	ug/kg-wet	
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	10	25	-	Heptachlor	<5	ug/kg-wet	
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	10	25	-	Heptachlor	<5	ug/kg-wet	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	9	23	-	Dieldrin	<10	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	9	23	-	Dieldrin	190	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	9	23	-	Dieldrin	50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	9	23	-	Dieldrin	240	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	9	23	-	Dieldrin	110	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	9	23	-	Dieldrin	120	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	9	23	-	Dieldrin	60	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	9	23	-	Dieldrin	120	ug/kg-wet	
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	9	23	-	Dieldrin	90	ug/kg-wet	
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	9	23	-	Dieldrin	40	ug/kg-wet

11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	9	23	-	Dieldrin	90	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	9	23	-	Dieldrin	100	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	9	23	-	Dieldrin	80	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	9	23	-	Dieldrin	20	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	9	23	-	Dieldrin	70	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	9	23	-	Dieldrin	110	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	9	23	-	Dieldrin	40	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	9	23	-	Dieldrin	70	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	9	23	-	Dieldrin	430	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	9	23	-	Dieldrin	40	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	9	23	-	Dieldrin	180	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	9	23	-	Dieldrin	140	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	9	23	-	Dieldrin	50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	9	23	-	Dieldrin	1,930	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	9	23	-	Dieldrin	230	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	9	23	-	Dieldrin	<10	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	8	19	DDE and DDD	p,p'-DDD	20	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	8	19	DDE and DDD	p,p'-DDD	940	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	8	19	DDE and DDD	p,p'-DDD	987	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	8	19	DDE and DDD	p,p'-DDD	2,440	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	8	19	DDE and DDD	p,p'-DDD	1,820	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	8	19	DDE and DDD	p,p'-DDD	1,150	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	8	19	DDE and DDD	p,p'-DDD	711	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	8	19	DDE and DDD	p,p'-DDD	1,380	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	8	19	DDE and DDD	p,p'-DDD	912	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	8	19	DDE and DDD	p,p'-DDD	611	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	8	19	DDE and DDD	p,p'-DDD	1,010	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	8	19	DDE and DDD	p,p'-DDD	887	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	8	19	DDE and DDD	p,p'-DDD	714	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	8	19	DDE and DDD	p,p'-DDD	194	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	8	19	DDE and DDD	p,p'-DDD	610	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	8	19	DDE and DDD	p,p'-DDD	246	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	8	19	DDE and DDD	p,p'-DDD	147	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	8	19	DDE and DDD	p,p'-DDD	136	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	8	19	DDE and DDD	p,p'-DDD	1,760	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	8	19	DDE and DDD	p,p'-DDD	221	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	8	19	DDE and DDD	p,p'-DDD	361	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	8	19	DDE and DDD	p,p'-DDD	530	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	8	19	DDE and DDD	p,p'-DDD	71	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	8	19	DDE and DDD	p,p'-DDD	4,780	ug/kg-wet

25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	8	19	DDE and DDD	p,p'-DDD	211	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	8	19	DDE and DDD	p,p'-DDD	20	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	8	19	DDE and DDD	o,p'-DDD	7	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	8	19	DDE and DDD	o,p'-DDD	136	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	8	19	DDE and DDD	o,p'-DDD	52	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	8	19	DDE and DDD	o,p'-DDD	257	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	8	19	DDE and DDD	o,p'-DDD	147	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	8	19	DDE and DDD	o,p'-DDD	110	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	8	19	DDE and DDD	o,p'-DDD	30	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	8	19	DDE and DDD	o,p'-DDD	132	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	8	19	DDE and DDD	o,p'-DDD	83	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	8	19	DDE and DDD	o,p'-DDD	53	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	8	19	DDE and DDD	o,p'-DDD	104	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	8	19	DDE and DDD	o,p'-DDD	116	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	8	19	DDE and DDD	o,p'-DDD	169	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	8	19	DDE and DDD	o,p'-DDD	34	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	8	19	DDE and DDD	o,p'-DDD	109	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	8	19	DDE and DDD	o,p'-DDD	111	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	8	19	DDE and DDD	o,p'-DDD	31	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	8	19	DDE and DDD	o,p'-DDD	37	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	8	19	DDE and DDD	o,p'-DDD	348	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	8	19	DDE and DDD	o,p'-DDD	88	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	8	19	DDE and DDD	o,p'-DDD	63	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	8	19	DDE and DDD	o,p'-DDD	79	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	8	19	DDE and DDD	o,p'-DDD	13	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	8	19	DDE and DDD	o,p'-DDD	392	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	8	19	DDE and DDD	o,p'-DDD	21	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	8	19	DDE and DDD	o,p'-DDD	<5	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	8	19	DDE and DDD	p,p'-DDE	192	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	8	19	DDE and DDD	p,p'-DDE	1,820	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	8	19	DDE and DDD	p,p'-DDE	5,090	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	8	19	DDE and DDD	p,p'-DDE	13,600	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	8	19	DDE and DDD	p,p'-DDE	6,390	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	8	19	DDE and DDD	p,p'-DDE	7,840	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	8	19	DDE and DDD	p,p'-DDE	2,490	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	8	19	DDE and DDD	p,p'-DDE	5,770	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	8	19	DDE and DDD	p,p'-DDE	3,780	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	8	19	DDE and DDD	p,p'-DDE	3,410	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	8	19	DDE and DDD	p,p'-DDE	2,820	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	8	19	DDE and DDD	p,p'-DDE	2,220	ug/kg-wet

13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	8	19	DDE and DDD	p,p'-DDE	3,330	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	8	19	DDE and DDD	p,p'-DDE	1,330	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	8	19	DDE and DDD	p,p'-DDE	2,830	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	8	19	DDE and DDD	p,p'-DDE	6,280	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	8	19	DDE and DDD	p,p'-DDE	618	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	8	19	DDE and DDD	p,p'-DDE	1,140	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	8	19	DDE and DDD	p,p'-DDE	10,800	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	8	19	DDE and DDD	p,p'-DDE	4,910	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	8	19	DDE and DDD	p,p'-DDE	1,310	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	8	19	DDE and DDD	p,p'-DDE	1,410	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	8	19	DDE and DDD	p,p'-DDE	195	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	8	19	DDE and DDD	p,p'-DDE	30,300	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	8	19	DDE and DDD	p,p'-DDE	984	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	8	19	DDE and DDD	p,p'-DDE	60	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	8	19	DDE and DDD	o,p'-DDE	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	8	19	DDE and DDD	o,p'-DDE	57	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	8	19	DDE and DDD	o,p'-DDE	89	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	8	19	DDE and DDD	o,p'-DDE	243	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	8	19	DDE and DDD	o,p'-DDE	170	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	8	19	DDE and DDD	o,p'-DDE	180	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	8	19	DDE and DDD	o,p'-DDE	71	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	8	19	DDE and DDD	o,p'-DDE	124	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	8	19	DDE and DDD	o,p'-DDE	92	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	8	19	DDE and DDD	o,p'-DDE	68	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	8	19	DDE and DDD	o,p'-DDE	82	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	8	19	DDE and DDD	o,p'-DDE	87	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	8	19	DDE and DDD	o,p'-DDE	213	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	8	19	DDE and DDD	o,p'-DDE	25	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	8	19	DDE and DDD	o,p'-DDE	75	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	8	19	DDE and DDD	o,p'-DDE	182	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	8	19	DDE and DDD	o,p'-DDE	22	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	8	19	DDE and DDD	o,p'-DDE	44	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	8	19	DDE and DDD	o,p'-DDE	236	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	8	19	DDE and DDD	o,p'-DDE	84	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	8	19	DDE and DDD	o,p'-DDE	38	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	8	19	DDE and DDD	o,p'-DDE	43	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	8	19	DDE and DDD	o,p'-DDE	9.0	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	8	19	DDE and DDD	o,p'-DDE	351	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	8	19	DDE and DDD	o,p'-DDE	13	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	8	19	DDE and DDD	o,p'-DDE	<5	ug/kg-wet

1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	7	18	DDT	p,p'-DDT	92	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	7	18	DDT	p,p'-DDT	554	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	7	18	DDT	p,p'-DDT	2,660	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	7	18	DDT	p,p'-DDT	6,610	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	7	18	DDT	p,p'-DDT	2,870	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	7	18	DDT	p,p'-DDT	3,500	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	7	18	DDT	p,p'-DDT	1,750	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	7	18	DDT	p,p'-DDT	4,170	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	7	18	DDT	p,p'-DDT	2,400	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	7	18	DDT	p,p'-DDT	1,670	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	7	18	DDT	p,p'-DDT	1,900	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	7	18	DDT	p,p'-DDT	1,540	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	7	18	DDT	p,p'-DDT	1,570	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	7	18	DDT	p,p'-DDT	557	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	7	18	DDT	p,p'-DDT	1,340	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	7	18	DDT	p,p'-DDT	251	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	7	18	DDT	p,p'-DDT	143	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	7	18	DDT	p,p'-DDT	165	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	7	18	DDT	p,p'-DDT	1,520	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	7	18	DDT	p,p'-DDT	360	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	7	18	DDT	p,p'-DDT	480	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	7	18	DDT	p,p'-DDT	780	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	7	18	DDT	p,p'-DDT	88	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	7	18	DDT	p,p'-DDT	3,090	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	7	18	DDT	p,p'-DDT	104	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	7	18	DDT	p,p'-DDT	20	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	7	18	DDT	o,p'-DDT	47	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	7	18	DDT	o,p'-DDT	345	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	7	18	DDT	o,p'-DDT	370	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	7	18	DDT	o,p'-DDT	1,030	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	7	18	DDT	o,p'-DDT	610	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	7	18	DDT	o,p'-DDT	652	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	7	18	DDT	o,p'-DDT	355	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	7	18	DDT	o,p'-DDT	775	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	7	18	DDT	o,p'-DDT	528	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	7	18	DDT	o,p'-DDT	251	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	7	18	DDT	o,p'-DDT	478	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	7	18	DDT	o,p'-DDT	514	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	7	18	DDT	o,p'-DDT	670	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	7	18	DDT	o,p'-DDT	204	ug/kg-wet

15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	7	18	DDT	o,p'-DDT	492	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	7	18	DDT	o,p'-DDT	1,020	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	7	18	DDT	o,p'-DDT	119	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	7	18	DDT	o,p'-DDT	242	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	7	18	DDT	o,p'-DDT	1,850	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	7	18	DDT	o,p'-DDT	591	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	7	18	DDT	o,p'-DDT	292	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	7	18	DDT	o,p'-DDT	368	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	7	18	DDT	o,p'-DDT	48	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	7	18	DDT	o,p'-DDT	2,270	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	7	18	DDT	o,p'-DDT	34	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	7	18	DDT	o,p'-DDT	12	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	6	16	-	cis-Nonachlor	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	6	16	-	cis-Nonachlor	94	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	6	16	-	cis-Nonachlor	41	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	6	16	-	cis-Nonachlor	227	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	6	16	-	cis-Nonachlor	136	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	6	16	-	cis-Nonachlor	139	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	6	16	-	cis-Nonachlor	51	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	6	16	-	cis-Nonachlor	134	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	6	16	-	cis-Nonachlor	87	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	6	16	-	cis-Nonachlor	56	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	6	16	-	cis-Nonachlor	113	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	6	16	-	cis-Nonachlor	114	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	6	16	-	cis-Nonachlor	81	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	6	16	-	cis-Nonachlor	40	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	6	16	-	cis-Nonachlor	99	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	6	16	-	cis-Nonachlor	112	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	6	16	-	cis-Nonachlor	132	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	6	16	-	cis-Nonachlor	106	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	6	16	-	cis-Nonachlor	820	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	6	16	-	cis-Nonachlor	106	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	6	16	-	cis-Nonachlor	115	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	6	16	-	cis-Nonachlor	200	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	6	16	-	cis-Nonachlor	23	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	6	16	-	cis-Nonachlor	1,640	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	6	16	-	cis-Nonachlor	610	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	6	16	-	cis-Nonachlor	13	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	6	16	-	trans-Nonachlor	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	6	16	-	trans-Nonachlor	473	ug/kg-wet

3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	6	16	-	trans-Nonachlor	462	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	6	16	-	trans-Nonachlor	1,260	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	6	16	-	trans-Nonachlor	967	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	6	16	-	trans-Nonachlor	793	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	6	16	-	trans-Nonachlor	384	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	6	16	-	trans-Nonachlor	706	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	6	16	-	trans-Nonachlor	406	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	6	16	-	trans-Nonachlor	593	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	6	16	-	trans-Nonachlor	568	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	6	16	-	trans-Nonachlor	537	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	6	16	-	trans-Nonachlor	650	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	6	16	-	trans-Nonachlor	281	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	6	16	-	trans-Nonachlor	648	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	6	16	-	trans-Nonachlor	920	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	6	16	-	trans-Nonachlor	441	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	6	16	-	trans-Nonachlor	579	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	6	16	-	trans-Nonachlor	2,900	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	6	16	-	trans-Nonachlor	754	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	6	16	-	trans-Nonachlor	801	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	6	16	-	trans-Nonachlor	786	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	6	16	-	trans-Nonachlor	130	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	6	16	-	trans-Nonachlor	7,570	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	6	16	-	trans-Nonachlor	1,200	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	6	16	-	trans-Nonachlor	32	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	5	15	-	Oxychlorane	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	5	15	-	Oxychlorane	79	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	5	15	-	Oxychlorane	62	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	5	15	-	Oxychlorane	320	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	5	15	-	Oxychlorane	218	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	5	15	-	Oxychlorane	237	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	5	15	-	Oxychlorane	71	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	5	15	-	Oxychlorane	152	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	5	15	-	Oxychlorane	90	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	5	15	-	Oxychlorane	89	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	5	15	-	Oxychlorane	116	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	5	15	-	Oxychlorane	108	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	5	15	-	Oxychlorane	187	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	5	15	-	Oxychlorane	39	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	5	15	-	Oxychlorane	135	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	5	15	-	Oxychlorane	92	ug/kg-wet

17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	5	15	-	Oxychlordane	30	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	5	15	-	Oxychlordane	86	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	5	15	-	Oxychlordane	495	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	5	15	-	Oxychlordane	64	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	5	15	-	Oxychlordane	111	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	5	15	-	Oxychlordane	136	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	5	15	-	Oxychlordane	17	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	5	15	-	Oxychlordane	1,190	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	5	15	-	Oxychlordane	200	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	5	15	-	Oxychlordane	6	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	4	14	Chlordane	trans-Chlordane	8	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	4	14	Chlordane	trans-Chlordane	15	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	4	14	Chlordane	trans-Chlordane	10	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	4	14	Chlordane	trans-Chlordane	10	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	4	14	Chlordane	trans-Chlordane	8	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	4	14	Chlordane	trans-Chlordane	13	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	4	14	Chlordane	trans-Chlordane	10	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	4	14	Chlordane	trans-Chlordane	10	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	4	14	Chlordane	trans-Chlordane	10	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	4	14	Chlordane	trans-Chlordane	14	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	4	14	Chlordane	trans-Chlordane	11	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	4	14	Chlordane	trans-Chlordane	12	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	4	14	Chlordane	trans-Chlordane	20	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	4	14	Chlordane	trans-Chlordane	21	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	4	14	Chlordane	trans-Chlordane	<10	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	4	14	Chlordane	trans-Chlordane	45	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	4	14	Chlordane	trans-Chlordane	17	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	4	14	Chlordane	trans-Chlordane	12	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	4	14	Chlordane	trans-Chlordane	11	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	4	14	Chlordane	trans-Chlordane	7	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	4	14	Chlordane	trans-Chlordane	<5	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	4	14	Chlordane	cis-Chlordane	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	4	14	Chlordane	cis-Chlordane	43	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	4	14	Chlordane	cis-Chlordane	16	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	4	14	Chlordane	cis-Chlordane	54	ug/kg-wet

5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	4	14	Chlordane	cis-Chlordane	29	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	4	14	Chlordane	cis-Chlordane	27	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	4	14	Chlordane	cis-Chlordane	18	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	4	14	Chlordane	cis-Chlordane	42	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	4	14	Chlordane	cis-Chlordane	39	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	4	14	Chlordane	cis-Chlordane	22	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	4	14	Chlordane	cis-Chlordane	32	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	4	14	Chlordane	cis-Chlordane	51	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	4	14	Chlordane	cis-Chlordane	33	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	4	14	Chlordane	cis-Chlordane	17	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	4	14	Chlordane	cis-Chlordane	43	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	4	14	Chlordane	cis-Chlordane	132	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	4	14	Chlordane	cis-Chlordane	109	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	4	14	Chlordane	cis-Chlordane	80	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	4	14	Chlordane	cis-Chlordane	459	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	4	14	Chlordane	cis-Chlordane	106	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	4	14	Chlordane	cis-Chlordane	156	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	4	14	Chlordane	cis-Chlordane	162	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	4	14	Chlordane	cis-Chlordane	41	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	4	14	Chlordane	cis-Chlordane	180	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	4	14	Chlordane	cis-Chlordane	89	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	4	14	Chlordane	cis-Chlordane	14	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	3	12		HCH total*	#REF!	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	3	12		HCH total*	#REF!	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	3	12		HCH total*	#REF!	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	3	12		HCH total*	#REF!	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	3	12		HCH total*	#REF!	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	3	12		HCH total*	#REF!	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	3	12		HCH total*	#REF!	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	3	12		HCH total*	#REF!	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	3	12		HCH total*	#REF!	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	3	12		HCH total*	#REF!	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	3	12		HCH total*	#REF!	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	3	12		HCH total*	#REF!	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	3	12		HCH total*	#REF!	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	3	12		HCH total*	#REF!	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	3	12		HCH total*	#REF!	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	3	12		HCH total*	#REF!	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	3	12		HCH total*	#REF!	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	3	12		HCH total*	#REF!	ug/kg-wet

19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	3	12	HCH total*	#REF!	ug/kg-wet	
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	3	12	HCH total*	#REF!	ug/kg-wet	
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	3	12	HCH total*	#REF!	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	3	12	HCH total*	#REF!	ug/kg-wet	
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	3	12	HCH total*	#REF!	ug/kg-wet	
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	3	12	HCH total*	#REF!	ug/kg-wet	
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	3	12	HCH total*	#REF!	ug/kg-wet	
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	3	12	HCH total*	#REF!	ug/kg-wet	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	3	12	-HCH	<10	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	3	12	-HCH	<10	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	3	12	-HCH	<10	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	3	12	-HCH	<10	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	3	12	-HCH	<10	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	3	12	-HCH	<10	ug/kg-wet	
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	3	12	-HCH	<10	ug/kg-wet	
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	3	12	-HCH	<10	ug/kg-wet	
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	3	12	-HCH	<10	ug/kg-wet	
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	3	12	-HCH	<10	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	3	12	-HCH	<10	ug/kg-wet	
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	3	12	-HCH	<10	ug/kg-wet	
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	3	12	-HCH	<10	ug/kg-wet	
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	3	12	-HCH	<10	ug/kg-wet	
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	3	12	-HCH	<10	ug/kg-wet	
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	3	12	-HCH	<10	ug/kg-wet	
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	3	12	-HCH	<10	ug/kg-wet	
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	3	12	-HCH	<10	ug/kg-wet	
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	3	12	-HCH	<10	ug/kg-wet	
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	3	12	-HCH	<10	ug/kg-wet	
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	3	12	-HCH	<10	ug/kg-wet	
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	3	12	-HCH	<10	ug/kg-wet	
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	3	12	-HCH	<10	ug/kg-wet	
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	3	12	-HCH	<10	ug/kg-wet	
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	3	12	-HCH	<10	ug/kg-wet	
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	3	12	-HCH	<10	ug/kg-wet	
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	3	12	-HCH	<10	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	3	12	-HCH	30	ug/kg-wet	
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	3	12	-HCH	<10	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	3	12	-HCH	<10	ug/kg-wet	
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	3	12	-HCH	<10	ug/kg-wet	
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	3	12	-HCH	<10	ug/kg-wet	

7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	3	12	-HCH	<10	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	3	12	-HCH	<10	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	3	12	-HCH	<10	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	3	12	-HCH	<10	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	3	12	-HCH	<10	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	3	12	-HCH	<10	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	3	12	-HCH	<10	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	3	12	-HCH	<10	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	3	12	-HCH	<10	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	3	12	-HCH	<10	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	3	12	-HCH	<10	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	3	12	-HCH	<10	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	3	12	-HCH	30	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	3	12	-HCH	<10	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	3	12	-HCH	30	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	3	12	-HCH	30	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	3	12	-HCH	20	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	3	12	-HCH	10	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	3	12	-HCH	<10	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	3	12	-HCH	<10	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	3	12	-HCH	<10	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	3	12	-HCH	1,550	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	3	12	-HCH	390	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	3	12	-HCH	1,680	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	3	12	-HCH	660	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	3	12	-HCH	870	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	3	12	-HCH	460	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	3	12	-HCH	810	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	3	12	-HCH	640	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	3	12	-HCH	260	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	3	12	-HCH	540	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	3	12	-HCH	520	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	3	12	-HCH	70	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	3	12	-HCH	30	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	3	12	-HCH	60	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	3	12	-HCH	270	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	3	12	-HCH	30	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	3	12	-HCH	140	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	3	12	-HCH	1,020	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	3	12	-HCH	30	ug/kg-wet

21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	3	12	-HCH	320	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	3	12	-HCH	230	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	3	12	-HCH	110	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	3	12	-HCH	2,330	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	3	12	-HCH	130	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	3	12	-HCH	10	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	3	12	Hexachlorocyclohexane	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	3	12	Hexachlorocyclohexane	72	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	3	12	Hexachlorocyclohexane	30	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	3	12	Hexachlorocyclohexane	50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	3	12	Hexachlorocyclohexane	20	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	3	12	Hexachlorocyclohexane	62	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	3	12	Hexachlorocyclohexane	40	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	3	12	Hexachlorocyclohexane	64	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	3	12	Hexachlorocyclohexane	40	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	3	12	Hexachlorocyclohexane	30	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	3	12	Hexachlorocyclohexane	40	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	3	12	Hexachlorocyclohexane	70	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	3	12	Hexachlorocyclohexane	20	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	3	12	Hexachlorocyclohexane	<10	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	3	12	Hexachlorocyclohexane	20	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	3	12	Hexachlorocyclohexane	24	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	3	12	Hexachlorocyclohexane	<10	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	3	12	Hexachlorocyclohexane	10	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	3	12	Hexachlorocyclohexane	117	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	3	12	Hexachlorocyclohexane	<10	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	3	12	Hexachlorocyclohexane	166	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	3	12	Hexachlorocyclohexane	192	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	3	12	Hexachlorocyclohexane	75	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	3	12	Hexachlorocyclohexane	17	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	3	12	Hexachlorocyclohexane	21	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	3	12	Hexachlorocyclohexane	<5	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	2	4	Hexachlorobenzene (HCB)	<5	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	2	4	Hexachlorobenzene (HCB)	448	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	2	4	Hexachlorobenzene (HCB)	200	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	2	4	Hexachlorobenzene (HCB)	546	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	2	4	Hexachlorobenzene (HCB)	342	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	2	4	Hexachlorobenzene (HCB)	319	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	2	4	Hexachlorobenzene (HCB)	290	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	2	4	Hexachlorobenzene (HCB)	348	ug/kg-wet

9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	2	4	Hexachlorobenzene (HCB)	299	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	2	4	Hexachlorobenzene (HCB)	236	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	2	4	Hexachlorobenzene (HCB)	277	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	2	4	Hexachlorobenzene (HCB)	282	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	2	4	Hexachlorobenzene (HCB)	395	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	2	4	Hexachlorobenzene (HCB)	68	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	2	4	Hexachlorobenzene (HCB)	300	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	2	4	Hexachlorobenzene (HCB)	116	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	2	4	Hexachlorobenzene (HCB)	61	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	2	4	Hexachlorobenzene (HCB)	301	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	2	4	Hexachlorobenzene (HCB)	549	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	2	4	Hexachlorobenzene (HCB)	73	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	2	4	Hexachlorobenzene (HCB)	285	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	2	4	Hexachlorobenzene (HCB)	323	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	2	4	Hexachlorobenzene (HCB)	113	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	2	4	Hexachlorobenzene (HCB)	218	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	2	4	Hexachlorobenzene (HCB)	84	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	2	4	Hexachlorobenzene (HCB)	31	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet

23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C PCB total*	#REF!	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Decichloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet

11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	240	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Nonachloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	920	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	70	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	450	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	120	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	410	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl	4,740	ug/kg-wet

25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl <50	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Octachloro biphenyl <50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl <50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl <50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 8,370	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 550	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 520	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 430	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 560	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 590	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 450	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 550	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 190	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 160	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 630	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 160	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 410	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 1,330	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 1,510	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 1,300	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 1,730	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 2,560	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl <50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 180	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl <50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 33,300	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl 1,820	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Heptachloro biphenyl <50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl <50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 660	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 22,300	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 2,820	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 2,480	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 2,360	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 1,550	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 2,330	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 1,290	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 1,790	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 1,170	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl 1,000	ug/kg-wet

13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	3,290	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	550	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	2,060	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	3,620	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	1,550	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	5,770	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	6,510	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	5,620	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	640	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	1,050	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	170	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	57,000	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	4,250	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Hexachloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	530	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	13,300	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,860	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,800	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,630	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,090	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,440	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	890	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	930	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	990	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	900	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	3,330	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	300	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,920	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	810	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	420	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	3,050	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	3,010	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,990	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	500	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	720	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	17,100	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	1,910	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Pentachloro biphenyl	<50	ug/kg-wet

1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	170	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	3,750	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	600	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	480	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	380	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	230	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	380	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	260	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	170	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	210	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	150	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	1,300	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	570	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	220	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	180	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	1,200	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	710	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	420	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	150	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	120	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	8,220	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	1,180	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	1	2	Polychlorinated biphenyls (PC Tetrachloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat	25	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver	42	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	310	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	90	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	1	2	Polychlorinated biphenyls (PC Trichloro biphenyl	<50	ug/kg-wet

15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	70	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	60	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	100	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	80	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Trichloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat 88	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver 86	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver 77	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver 81	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat 78	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver 64	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver 89	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver 82	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver 71	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver 83	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver 67	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver 62	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver 20	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver 65	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver -	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver 90	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver 78	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver 88	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver 70	1	2	Polychlorinated biphenyls (P C Dichloro biphenyl	<50	ug/kg-wet
1	Fin whales	Yamaguchi Pref.	F	A	Fat 25	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
2	Minke whales	Niigata Pref.	F	J	Fat, liver 42	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet

3	Mesoplodon stejnegeri	Niigata Pref.	M	A	Fat	88	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
4	Mesoplodon stejnegeri	Yamagata Pref.	M	S	Fat, liver	78	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
5	Mesoplodon stejnegeri	Niigata Pref.	M	S	Fat, liver	86	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
6	Mesoplodon stejnegeri	Aomori Pref.	M	S	Fat, liver	77	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
7	Mesoplodon stejnegeri	Ishikawa Pref.	F	A	Fat, liver	81	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
8	Mesoplodon stejnegeri	Tottori Pref.	F	A	Fat, liver	70	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
9	Mesoplodon stejnegeri	Hokkaido	F	AL	Fat, liver	81	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
10	Mesoplodon stejnegeri	Niigata Pref.	F	Pr	Fat	78	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
11	Mesoplodon stejnegeri	Yamagata Pref.	F	S	Fat, liver	64	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
12	Mesoplodon stejnegeri	Niigata Pref.	F	J	Fat, liver	89	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
13	Mesoplodon carlhubbsi	Iwate Pref.	F	N	Fat, liver	82	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
14	Blainville's beaked whale	Aichi Pref.	F	AL	Fat, liver	71	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
15	Blainville's beaked whale	Aichi Pref.	F	N	Fat, liver	83	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
16	Feral dolphin	Iwate Pref.	M	S	Fat, liver	67	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
17	Feral dolphin	Kanagawa Pref.	F	Pr	Fat, liver	62	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
18	Pacific white-sided dolphin	Ibaraki Pref.	M	N	Fat, liver	20	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
19	Pacific white-sided dolphin	Niigata Pref.	F	S	Fat, liver	65	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
20	Melon-headed whale	Shizuoka Pref.	M	A	Fat, liver	-	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
21	Harbor porpoise	Iwate Pref.	M	A	Fat, liver	90	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
22	Harbor porpoise	Hokkaido	M	S	Fat, liver	88	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
23	Harbor porpoise	Hokkaido	F	A	Fat, liver	90	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
24	Finless porpoise	Yamaguchi Pref.	M	A	Fat, liver	78	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
25	Finless porpoise	Yamaguchi Pref.	M	J	Fat, liver	88	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet
26	Finless porpoise	Chiba Pref.	F	N	Fat, liver	70	1	2	Polychlorinated biphenyls (P C Chlorinated biphenyl	<50	ug/kg-wet