

Results of Radioactive Material Monitoring of Aquatic Organisms (Location L off Soma City)

<Location L off Soma City: Samples collected>

Items	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
L-1	-	○	-	-	○	-
L-2	○	○	○	○	○	○
L-3	○	○	○	-	○	-

<Location L off Soma City: Site measurement item>

Items	Latitude and longitude of the location		Survey date and time			Water	Sediment			Other		
	Latitude	Longitude	Date	Time (water)	Time (sediment)		Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)
L-1	37.8210°	140.9610°	2016/8/26	-	10:11	-	25.8	Silt with sand	10Y3/1	None	-	-
L-2	37.8155°	140.9763°		08:05	09:36	25.0	25.5	Sand with silt	7.5Y5/2	Shells, Plant pieces	1.8	>1.8
L-3	37.8217°	140.9765°		08:24	09:52	25.1	25.6	Silt with sand	10Y4/1	None	1.4	>1.4

<Location L off Soma City: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity (mg/L)	TOC (FNU)	SS (mg/L)	Turbidity (Bq/L)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
L-2	37.8155°	140.9763°	2016/8/26	08:05	8.0	<0.5	2.8	6.3	4640	30.16	1.1	6	3.4	0.0049	0.032	0.0010
L-3	37.8217°	140.9765°		08:24	8.0	<0.5	3.4	6.7	4470	28.90	1.1	5	2.9	0.0031	0.019	-

<Location L off Soma City: General survey items/Analysis of radioactive materials Sediment>

Items	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution						Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)	
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)		
L-1	37.8210°	140.9610°	2016/8/26	10:11	7.9	-9	32.9	2.1	4.7	2.688	1.7	5.8	27.3	42.2	11.9	0.19	9.5	11	90	-
L-2	37.8155°	140.9763°		09:36	7.7	241	23.6	1.1	3.2	2.726	1.7	8.0	76.3	9.4	1.1	3.5	0.42	9.5	54	N.D.(0.13)
L-3	37.8217°	140.9765°		09:52	7.8	248	25.2	1.1	2.2	2.705	0.0	0.4	43.9	47.7	2.0	6.0	0.23	2.0	8.7	36

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location L off Soma City: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
L-1 L-2 L-3	Matsukawaura	37.8210° 37.8155° 37.8217°	140.9610° 140.9763° 140.9765°	2016/8/26	Algae/plant	-	-	-	Plankton (Planktonic algae)		-	0.020	-	-	-	19.8	2.8	17	-
					Algae/plant	Monocotyledoneae	Najadales	Zosteraceae	Zostera marina	Eel grass	-	1.4	-	-	-	0.59	0.13	0.46	-
				2016/8/20	Arthropoda	Malacostraca	Mysida	Mysidae		Mysidae	-	0.29	Imago	-	-	4.06	0.66	3.4	-
				2016/8/24	Arthropoda	Malacostraca	Decapoda	Varunidae	Hemigrapsus sp.	Hemigrapsus	110	0.23	Imago	-	-	2.37	0.37	2.0	-
				2016/8/20	Mollusca	Bivalvia	Ostreidae	Crassostrea gigas	Oyster		18	0.27	Imago	-	-	1.2	N.D.(0.36)	1.2	-
					Mollusca	Bivalvia	Veneridae	Ruditapes philippinarum	Japanese littleneck		90	0.31	Imago	-	-	3.53	0.53	3.0	-
					Vertebrata	Osteichthyes	Perciformes	Mugilidae	Mugil cephalus	Flathead mullet	12	0.18	Immature fish	Obscure digesta	Viscera removed	1.7	N.D.(0.57)	1.7	-
					Vertebrata	Osteichthyes	Clupeiformes	Clupeidae	Konosirus punctatus	Dotted gizzard shad	119	0.50	Immature fish	-	-	3.62	0.62	3.0	-
					Vertebrata	Osteichthyes	Tetraodontiformes	Tetraodontidae	Takifugu niphobles	Takifugu niphobles	2	0.19	Mature fish	Obscure digesta	Viscera removed	2.1	N.D.(0.44)	2.1	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.