### **FY2016 Radioactive Material Monitoring of Aquatic Organisms (December)**

#### 1. Survey Overview

Samples of aquatic organisms (algae, aquatic insects, crustaceans, shellfishes, fishes, and amphibians, etc.) were collected mainly in Fukushima Prefecture and concentrations of radioactive cesiums and radioactive strontium in the samples were measured (survey period: December 1 to 8, 2016).

In order to clarify the environment of the water areas where aquatic organisms live, surveys were also conducted on general items concerning water and sediments and activity concentrations in these water areas.

The following water areas were selected based on the results of the past Radioactive Material Monitoring of Aquatic Organisms and Radioactive Material Monitoring in the Water Environment in and around Fukushima Prefecture, as well as the results of the measurement of radioactive materials in fisheries products conducted by other relevant organizations and interviews with local fishermen.

- (i) Rivers: Abukuma River, Uda River, Mano River, Niida River, and Ota River
- (ii) Lakes: Lake Hayama, Lake Akimoto, Lake Inawashiro
- (iii) Sea areas: Off the mouth of the Abukuma River, off Soma City, off Iwaki City

# o Survey locations and dates

| Aı         | rea    | Targeted water areas | Zone   | Item                       | Survey dates               | Remarks   |  |  |
|------------|--------|----------------------|--|----------------------------|----------------------------|---|--|--|
|            | A      |                      | Shinfuna Bridge to the<br>Iinoentei Dam;Harase River (a        | Aquatic organisms sampling | December 2, 2016           | Algae/Plants, Aquatic insects, Crustanceans, Shellfishes, Fishes, Amphibians, Fallen leaves, etc. |  |  |
|            |        |                      | tributary)   | Water/sediment sampling    | December 9, 2016           | (Water sampling) A-1,A-2 (Sediment sampling) A-1,A-2  |  |  |
|            | В      |                      | Confluence with the Matsukawa<br>River (a tributary) to Taisho | Aquatic organisms sampling | December 2, and 8,<br>2016 | Algae/Plants, Aquatic insects, Crustanceans, Fishes, Amphibians, Fallen leaves, etc.              |  |  |
|            |        |                      | Bridge;Surikami River (a<br>tributary)                         | Water/sediment sampling    | December 9, 2016           | (Water sampling) B-1-B-3 (Sediment sampling) B-1-B-3  |  |  |
| _          | С      | Uda River            | Kawadaira Bridge to Horisaka<br>Bridge;Around Tamano           | Aquatic organisms sampling | December 4, 2016           | Algae/Plants, Aquatic insects, Crustanceans, Fishes, Fallen leaves, etc.                          |  |  |
| River area | C      |                      | Bridge   | Water/sediment sampling    | December 8, 2016           | (Water sampling) C-1-C-6 (Sediment sampling) C-1,C-2,C-4-C-6                                      |  |  |
| area       | D      | Mano River           | Zennami Bridge to Ochiai                                       | Aquatic organisms sampling | December 7, 2016           | Algae/Plants, Aquatic insects, Fishes, Fallen leaves, etc.  |  |  |
|            |        |                      | Bridge   | Water/sediment sampling    | December 5, 2016           | (Water sampling) D-1-D-5 (Sediment sampling) D-1-D-3,D-4a,D-5                                     |  |  |
|            | Е      |                      | Kayanoki Bridge to Sugauchi                                    | Aquatic organisms sampling | December 8, 2016           | Algae/Plants, Aquatic insects, Fishes, Fallen leaves, etc.  |  |  |
|            |        |                      | Bridge   | Water/sediment sampling    | December 7, 2016           | (Water sampling) E-1-E-5 (Sediment sampling) E-1,E-2a,E-3-E-5                                     |  |  |
|            | F      | Ota River            | Yaigomesaka Bridge to  | Aquatic organisms sampling | December 6, 2016           | Algae/Plants, Aquatic insects, Fishes, Fallen leaves, etc.  |  |  |
|            |        | Ota Kivei            | Memezawa district  | Water/sediment sampling    | · ·                        | (Water sampling) F-1-F-6 (Sediment sampling) F-1-F-5  |  |  |
|            | G      | Lake Hayama          |  | Aquatic organisms sampling | December 5, and 8,<br>2016 | Algae/Plants, Aquatic insects, Fishes, Amphibians, Fallen leaves, etc.                            |  |  |
|            |        |                      |  | Water/sediment sampling    | December 5, 2016           | (Water sampling) G-1,G-3,G-5 (Sediment sampling) G-1-G-5  |  |  |
|            | Н      | Lake Akimoto         |  | Aquatic organisms sampling | December 1, 2016           | Algae/Plants, Aquatic insects, Crustanceans, Fishes   |  |  |
| Lake area  |        |                      |  | Water/sediment sampling    | December 1, 2016           | (Water sampling) H-1,H-3,H-5 (Sediment sampling) H-1-H-5  |  |  |
| area       | I<br>J | Lake Inawashiro      | North lakeside   | Aquatic organisms sampling | -                          | -   |  |  |
|            |        |                      | North fakeside   | Water/sediment sampling    | December 3, 2016           | (Water sampling) I-1,I-3 (Sediment sampling) I-1-I-4  |  |  |
|            |        |                      | South lakeside   | Aquatic organisms sampling | December 2, and 3,<br>2016 | Algae/Plants, Shellfishes   |  |  |
|            |        |                      |  | Water/sediment sampling    | December 3, 2016           | (Water sampling) J-1 (Sediment sampling) J-1  |  |  |
|            | K      | the Abukuma          | Sea area in front of the                                       | Aquatic organisms sampling | December 7, 2016           | Crustanceans, Fishes  |  |  |
|            |        |                      | Abukuma River Estuary  | Water/sediment sampling    |                            | (Water sampling) K-2 (Sediment sampling) K-1-K-3  |  |  |
| Sea area   | L      | Off Soma City        | Matsukawaura Lagoon  | Aquatic organisms sampling | December 4, and 8,<br>2016 | Seaweeds/Algae, Polychaetes, Shellfishes  |  |  |
| area       | L      | on John City         |  | Water/sediment sampling    | December 8, 2016           | (Water sampling) L-2,L-3 (Sediment sampling) L-1-L-3  |  |  |
|            | М      | Off Iwaki City       | Offshore of Hisanohama   | Aquatic organisms sampling | December 3, 2016           | Seaweeds/Algae, Sea urchins, Starfishes, Shellfishes, Fishes                                      |  |  |
|            |        |                      | Onshore of thisahohania  | Water/sediment sampling    | December 3, 2016           | (Water sampling) M-2 (Sediment sampling) M-1-M-3  |  |  |

### 2. Survey Items and Locations, etc.

### 2.1 Survey Items

For all samples of aquatic organisms, analysis of radioactive cesiums (Cs-134, Cs-137) was conducted. Additionally, for samples of large fish, etc. analysis of radioactive strontium (Sr-90) was also conducted.

With regard to surveys of water and sediments, locations where aquatic organism samples were scheduled to be collected and where clay particles and coarse particulate organic matters (Fallen leaves at the bottom, etc.) are supposed to accumulate due to inflows from the surrounding environment, etc. were selected for the analysis of radioactive materials and general survey items.

Survey items and samples for aquatic organisms, water, and sediments are as shown in the following table.


### Survey targets and items

| Target    |               | Measurement item                    | Analyzed samples   |  |  |  |
|-----------|---------------|-------------------------------------|--|--|--|--|
| Aquatic   | Radioactive   | Radioactive cesiums (Cs-134,Cs-137) | All samples  |  |  |  |
| Organisms | materials     | Radioactive strontium (Sr-90)       | Large fish, etc.   |  |  |  |
|           | Radioactive   | Radioactive cesiums (Cs-134,Cs-137) | Samples collected at one to six locations for each water area    |  |  |  |
|           | materials     | Radioactive strontium (Sr-90)       | Samples collected at one location for each water area            |  |  |  |
|           |               | pH                                  |  |  |  |  |
|           |               | BOD (Biological oxygen demand)      |  |  |  |  |
| Water     |               | COD (Chemical oxygen demand)        |  |  |  |  |
|           |               | DO (Dissolved oxygen level)         | Samples collected at one to six locations for each               |  |  |  |
|           | General items | Electric conductivity               | water area   |  |  |  |
|           |               | Salinity                            |  |  |  |  |
|           |               | TOC (Total organic carbon)          |  |  |  |  |
|           |               | SS (Suspended solids)               |  |  |  |  |
|           |               | Turbidity                           |  |  |  |  |
|           | Radioactive   | Radioactive cesiums (Cs-134,Cs-137) | Samples collected at three to five locations for each water area |  |  |  |
|           | materials     | Radioactive strontium (Sr-90)       | Samples collected at one location for each water area            |  |  |  |
|           |               | pH                                  |  |  |  |  |
| Sediments |               | Oxidation-reduction potential       |  |  |  |  |
|           |               | Water content                       | Samples collected at three to five locations for each            |  |  |  |
|           | General items | IL (Ignition loss)                  | water area   |  |  |  |
|           |               | TOC (Total organic carbon)          | water area   |  |  |  |
|           |               | Soil particle density               |  |  |  |  |
|           |               | Grain size distribution             |  |  |  |  |


#### 2.2 Survey Locations at Respective Water Areas

(1) Tributaries to the Abukuma River (Location A along the Abukuma River; Location B along the Abukuma River; Location K off the mouth of the Abukuma River).

As water areas where clay particles and CPOMs (dead leaves at the bottom, etc.) are supposed to accumulate topographically, Location A along the Abukuma River was set from the Harase River (a tributary to the Abukuma River) and Shinfuna Bridge (Nihonmatsu City, Fukushima Prefecture) to the Iino Dam, and Location B along the Abukuma River was set from the confluence with the Matsukawa River to Taisho Bridge (Date City, Fukushima Prefecture) as well as the zone where a tributary to the Surikami River inflows. Additionally, Location K was set off the mouth of the Abukuma River in order to survey the sea area in front of the mouth of the Abukuma River, where the outflow of radioactive materials through the Abukuma River is suspected.




Detailed map showing Location K off the mouth of the Abukuma River



Map showing Location A and Location B along the Abukuma River

# (2) Location C along the Uda River


Surveys were started in the autumn term of FY2012 for the location from Kawadaira Bridge to Horisaka Bridge, where water flows into the Matsugafusa Dam (Lake Udagawa), and around Tamano Bridge, where water flows into the Tamano Reservoir (a tributary to the Tamano River).




Detailed map showing Location C along the Uda River

(3) Tributaries to the Mano River (Location D along the Mano River; Location G in Lake Hayama)

Surveys were conducted at Location D along the Mano River, which covers from Zennami Bridge to Ochiai Bridge (Kashima Ward, Minamisoma City, Fukushima Prefecture), and at Location G in Lake Hayama (Mano Dam), which covers the lake as a whole and inflow points.




Detailed map showing Location D along the Mano River



Detailed map showing Location G in Lake Hayama (Mano Dam)

# (4) Location E along the Niida River


Surveys were conducted from Kayanoki Bridge to Sugauchi Bridge.



Detailed map showing Location E along the Niida River

# (5) Location F along the Ota River


Surveys were started in the autumn term of FY2012 for the location from Yaigomesaka Bridge to Memezawa District.



Detailed map showing Location F along the Ota River

## (6) Location H in Lake Akimoto


Surveys were conducted in the whole area of Lake Akimoto, and around Lake Akimoto.



Detailed map showing Location H in Lake Akimoto

## (7) Location I (North Lakeside) and Location J (South Lakeside) in Lake Inawashiro

Surveys were conducted at around the point where the Nagase River inflows into Lake Inawashiro, and at around the point where lake water flows out into the Nippashi River (at the north lakeside), and at the south lakeside.




Detailed map showing Location I (north lakeside) and Location J (south lakeside) in Lake Inawashiro

## (8) Location L off Soma City

Surveys were conducted within the Matsukawaura Lagoon, centering on the estuary region of the Uda River.


Sampling point in Location L-2 was expanded to the south in the FY2015 survey because sampling was impossible at the conventional point due to bank protection work.



Detailed map showing Location L off Soma City (Matsukawaura Lagoon)

# (9) Location M off Iwaki City

Surveys were conducted at offshore of the Hisanohama Fishing Port and coastal areas in Hisanohama.



Detailed map showing Location M off Iwaki City

### 3. Results

Survey results are shown in the table.

The outline of the measurement results of radioactive cesiums (the total of Cs-134 and Cs-137).

# (i) Rivers and lakes

Unit:Bq/kg-wet

|                  |                    |                      |                            |                            |                            |                                    |                             |                           | Unit:Bq/kg-wet                    |
|------------------|--------------------|----------------------|----------------------------|----------------------------|----------------------------|------------------------------------|-----------------------------|---------------------------|-----------------------------------|
| Water area       |                    | Time                 | Algae,<br>Plants           | Aquatic insects            | Crustaceans                | Shellfishes<br>(Molluscan<br>body) | Fishes                      | Amphibians                | CPOMs<br>(fallen leaves,<br>etc.) |
| Abukuma<br>River | Abukuma<br>River A | FY2016<br>Dec.       | 180                        | 17.8 , 26.3<br>(2 species) | 16.6                       | 16.9                               | 9.3 , 10.9<br>(2 species)   | 20.6                      | 26.7                              |
|                  |                    | FY2016<br>Oct.       | 201                        | 14.7                       | 19.4                       | 15.7                               | 10 - 16.2<br>(5 species)    | 3.5 - 135<br>(3 species)  | 16.5                              |
|                  |                    | FY2016<br>Aug.       | -                          | 32.9                       | 31.3                       | 26.0                               | 7.1 , 8.3<br>(2 species)    | 104                       | 62                                |
|                  |                    | FY2016<br>Jun.       | 278                        | 11.9 , 30.5<br>(2 species) | 26.1 , 26.9<br>(2 species) | 11                                 | 6.7 - 21.1<br>(9 species)   | 10.6 - 177<br>(3 species) | 80                                |
| System           |                    | FY2016<br>Dec.       | 125                        | N.D 30.5<br>(3 species)    | 9.2                        | -                                  | 4.15 - 15.1<br>(4 species)  | 5.2                       | 28.8                              |
|                  | Abukuma            | FY2016<br>Sep Oct.   | 113                        | 2.2 , 20.7<br>(2 species)  | 5.9                        | 1                                  | N.D 32.5<br>(10 species)    | 46.0                      | 37.0                              |
|                  | River B            | FY2016<br>Aug Sep.   | 97                         | 2.0 - 44.0<br>(3 species)  | 13.0                       | ı                                  | 4.30 - 52.1<br>(10 species) | -                         | 12.8                              |
|                  |                    | FY2016<br>Jun Jul.   | 46.5 , 193<br>(2 species)  | 13.7 - 41.6<br>(3 species) | 5.4 , 15.2<br>(2 species)  | i                                  | 2.5 - 89<br>(16 species)    | 7.8 , 50.1<br>(2 species) | 29.8                              |
|                  |                    | FY2016<br>Dec.       | 188                        | N.D 57.4<br>(4 species)    | 8.4 , 16.2<br>(2 species)  | -                                  | 6.4 - 15<br>(4 species)     | -                         | 7.5                               |
| Uda              | Piver C            | FY2016<br>Oct.       | 16.0 , 84.7<br>(2 species) | -                          | 19.4                       | -                                  | 5.5 - 19.2<br>(7 species)   | -                         | 12.6                              |
| Oua              | Uda River C        |                      | 1                          | 10                         | 10.3 , 18.1<br>(2 species) | ī                                  | N.D 22.9<br>(9 species)     | 85                        | 49.3                              |
|                  |                    |                      | 147                        | 7.2 - 68<br>(3 species)    | 9.3 , 15.3<br>(2 species)  | -                                  | N.D 64.5<br>(10 species)    | -                         | 52.1                              |
|                  | Lake<br>Hayama G   | FY2016<br>Dec.       | 2.4 , 222<br>(2 species)   | 4.3 - 103<br>(3 species)   | -                          | -                                  | 28.3 - 400<br>(4 species)   | 18.4                      | 45.5                              |
|                  |                    | FY2016<br>Oct.       | 4.6 , 660<br>(2 species)   | 326                        | -                          | -                                  | 12.8 - 659<br>(5 species)   | -                         | 75                                |
|                  |                    | FY2016<br>Aug.       | 6.6 , 120<br>(2 species)   | 2.10 , 22<br>(2 species)   | 64                         | -                                  | 15 - 521<br>(6 species)     | -                         | 285                               |
| Mano<br>River    |                    | FY2016<br>May - Jun. | 11.1, 235<br>(2 species)   | 11 - 117<br>(3 species)    | 49.7                       | -                                  | 37.8 - 189<br>(8 species)   | -                         | 310                               |
| System           | Mano River<br>D    | FY2016<br>Dec.       | 63.0 , 365<br>(2 species)  | 4.0 - 89<br>(4 species)    | -                          | -                                  | 15.9 - 38.1<br>(3 species)  | -                         | 20.3                              |
|                  |                    | FY2016<br>Sep Oct.   | 153 , 282<br>(2 species)   | -                          | 22.4 , 33.1<br>(2 species) | 94, 364<br>(2 species)             | 14.4 - 54.4<br>(4 species)  | -                         | 21.4                              |
|                  |                    | FY2016<br>Aug.       | 660                        | 37.5                       | 45 - 76<br>(3 species)     | 36.4, 248<br>(2 species)           | 14 - 28.4<br>(6 species)    | 6.3                       | 21.6                              |
|                  |                    | FY2016<br>Jun.       | 170                        | 22 - 47.8<br>(3 species)   | 31.5 - 144<br>(3 species)  | -                                  | 14.8 - 74<br>(8 species)    | 444                       | 38.4                              |
| Niida River E    |                    | FY2016<br>Dec.       | 15.0 , 72<br>(2 species)   | 5.8 - 185<br>(4 species)   | -                          | -                                  | 40.9 - 81<br>(3 species)    | -                         | 85                                |
|                  |                    | FY2016<br>Oct.       | 192                        | 88.3                       | 40.2 , 80<br>(2 species)   | -                                  | 19.2 - 59.0<br>(5 species)  | 48.1 , 359<br>(1 species) | 186                               |
|                  |                    | FY2016<br>Aug.       | 194                        | 108                        | 49.1 - 72<br>(4 species)   | -                                  | 27.1 - 426<br>(7 species)   | 37.3 , 507<br>(2 species) | 295                               |
|                  |                    | FY2016<br>Jun Jul.   | 291                        | 27 - 205<br>(4 species)    | 32.1 - 83<br>(4 species)   | -                                  | 32.8 - 204<br>(14 species)  | -                         | 1050                              |
|                  |                    |                      |                            |                            |                            |                                    |                             |                           |                                   |

<sup>\*</sup> ND means to be below the detection limit.

<sup>\*</sup> Organisms were collected in or around the targeted water areas.

<sup>\*</sup> Basically, measurement was conducted for all targeted samples, not limited to edible parts.

<sup>\*</sup>Since the autumn term of FY2012, sampling and analysis of aquatic insects had been conducted separately for four categories (Plecoptera, Trichoptera, Odonata, and Megaloptera) (by feeding habit and type). Since the FY2014 June-July Survey, Ephemeroptera was added and sampling and analysis were conducted for five categories.

Unit:Bq/kg-wet

| Water area         |   | Time                 | Algae,<br>Plants           | Aquatic insects           | Crustaceans              | Shellfishes<br>(Molluscan<br>body) | Fishes                      | Amphibians                | CPOMs<br>(fallen leaves,<br>etc.) |
|--------------------|---|----------------------|----------------------------|---------------------------|--------------------------|------------------------------------|-----------------------------|---------------------------|-----------------------------------|
| Ota River F        |   | FY2016<br>Dec.       | 7.7 , 2950<br>(2 species)  | 65.0 - 930<br>(4 species) | -                        | -                                  | 223 , 412<br>(2 species)    | -                         | 160                               |
|                    |   | FY2016<br>Oct.       | 1000 - 2120<br>(3 species) | 176                       | 292                      | 267                                | 224 - 431<br>(4 species)    | 2000                      | 141                               |
|                    |   | FY2016<br>Aug.       | 1900                       | 249, 389<br>(2 species)   | 392 - 531<br>(3 species) | -                                  | 128 - 630<br>(8 species)    | -                         | 86                                |
|                    |   | FY2016<br>May - Jul. | 1320                       | 111, 211<br>(2 species)   | 333 - 780<br>(3 species) | -                                  | 45.7 - 2860<br>(12 species) | 840                       | 2040                              |
|                    |   | FY2016<br>Dec.       | N.D.                       | 22.4                      | 22.9                     | -                                  | 16.8 - 109<br>(8 species)   | -                         | -                                 |
| Lake A             | Lake Akimoto H                              |                      | N.D., 3.85<br>(2 species)  | 10.9                      | 18.9                     | 8.2                                | 10.2 - 66.8<br>(10 species) | -                         | 17.8                              |
| Euro 1             |   |                      | 41.3 , 54.1<br>(2 species) | 7.2                       | 18, 36.3<br>(2 species)  | -                                  | 5.6 - 78<br>(9 species)     | 9.0 - 21.0<br>(3 species) | 56.5                              |
|                    |   |                      | 1.4, 9.4<br>(2 species)    | N.D., 26.0<br>(2 species) | 39.8                     | -                                  | N.D 99<br>(14 species)      | 13.4 - 259<br>(4 species) | 58.7                              |
|                    | Lake<br>Inawashiro<br>I (north<br>lakeside) | FY2016<br>Dec.       | -                          | -                         | -                        | -                                  | -                           | -                         | -                                 |
|                    |   | FY2016<br>Oct.       | -                          | 1                         | -                        | -                                  | 9.2 - 53.8<br>(9 species)   | -                         | 1.40                              |
|                    |   | FY2016<br>Aug.       | -                          | -                         | -                        | -                                  | 6.69 - 69<br>(4 species)    | -                         | 132                               |
|                    |   | FY2016<br>Jun.       | -                          | -                         | -                        | -                                  | 15.5 - 58.6<br>(5 species)  | -                         | 31.3                              |
| Lake<br>Inawashiro | Lake<br>Inawashiro<br>J (south<br>lakeside) | FY2016<br>Dec.       | N.D.                       | ı                         | -                        | N.D.                               | -                           | -                         | -                                 |
|                    |   | FY2016<br>Oct.       | N.D 2.72<br>(4 species)    | -                         | -                        | N.D 3.3<br>(3 species)             | 1.5 - 50.1<br>(10 species)  | 1.4 - 15.3<br>(3 species) | -                                 |
|                    |   | FY2016<br>Aug.       | 0.78 - 1.97<br>(3 species) | -                         | -                        | 2.3                                | 1.9 - 131<br>(7 species)    | N.D., 2.5<br>(2 species)  | -                                 |
|                    |   | FY2016<br>Jun.       | 1.7 , 2.4<br>(2 species)   | -                         | 7.7                      | N.D.                               | N.D 101<br>(8 species)      | 3.44 , 4.8<br>(2 species) | -                                 |

<sup>\*</sup> ND means to be below the detection limit.

<sup>\*</sup> Organisms were collected in or around the targeted water areas.

<sup>\*</sup> Basically, measurement was conducted for all targeted samples, not limited to edible parts.

<sup>\*</sup>Since the autumn term of FY2012, sampling and analysis of aquatic insects had been conducted separately for four categories (Plecoptera, Trichoptera, Odonata, and Megaloptera) (by feeding habit and type). Since the FY2014 June-July Survey, Ephemeroptera was added and sampling and analysis were conducted for five categories.

# (ii) Sea areas

Unit:Bq/kg-wet

| Water area                      | Time           | Seaweeds,<br>Algae         | Polychaetes | Sea urchins,<br>Starfishes,<br>Trepangs | Crustaceans                | Shellfishes<br>(Molluscan<br>body) | Squids,<br>Octopuses | Fishes                     |
|---------------------------------|----------------|----------------------------|-------------|---|----------------------------|------------------------------------|----------------------|----------------------------|
|                                 | FY2016<br>Dec. | -                          | -           | -                                       | 2.10                       | -                                  | -                    | 0.67 - 2.36<br>(3 species) |
| Location K off the mouth of the | FY2016<br>Oct. | -                          | 1           | -                                       | 0.48                       | -                                  | -                    | 0.48 - 1.56<br>(4 species) |
| Abukuma River                   | FY2016<br>Sep. | -                          | ı           | ٠                                       | -                          | 1                                  | -                    | 0.47 - 0.86<br>(3 species) |
|                                 | FY2016<br>Jun. | -                          | •           | -                                       | 0.53                       | -                                  | -                    | N.D 1.1<br>(5 species)     |
|                                 | FY2016<br>Dec. | 1.72 , 7.1<br>(2 species)  | 15.1        | -                                       | -                          | 1.1 , 1.40<br>(2 species)          | -                    | -                          |
| Location L off<br>Soma City     | FY2016<br>Oct. | 1.12, 7.2<br>(2 species)   | ı           | ·                                       | N.D., 4.17<br>(2 species)  | 0.43 , 1.3<br>(2 species)          | -                    | 1.4 - 2.4<br>(4 species)   |
| (Matsukawaura Lagoon)           | FY2016<br>Aug. | 0.59 , 19.8<br>(2 species) | 1           | -                                       | 2.37 , 4.06<br>(2 species) | 1.2 , 3.53<br>(2 species)          | -                    | 1.7 - 3.62<br>(3 species)  |
|                                 | FY2016<br>Jun. | 1.91 , 6.6<br>(2 species)  | ı           | ٠                                       | 3.84                       | 1.72, 3.56<br>(2 species)          | -                    | 1.8 , 27.4<br>(2 species)  |
|                                 | FY2016<br>Dec. | 6.03                       | ı           | N.D., 1.2<br>(2 species)                | •                          | 0.74                               | -                    | 0.53 - 4.96<br>(6 species) |
| Location M off<br>Iwaki City    | FY2016<br>Oct. | 2.13                       | 1           | 0.31                                    | -                          | 0.42                               | -                    | 0.73 - 4.79<br>(9 species) |
| (Hisanohama)                    | FY2016<br>Sep. | 3.35                       | -           | 0.43                                    | 4.00                       | 0.88                               | N.D.                 | N.D 5.84<br>(10 species)   |
|                                 | FY2016<br>Jun. | 4.78                       | -           | 1.5                                     | -                          | 0.42                               | N.D.                 | N.D 14.3<br>(14 species)   |

<sup>\*</sup> ND means to be below the detection limit.

<sup>\*</sup> Organisms were collected in or around the targeted water areas.

<sup>\*</sup> Basically, measurement was conducted for all targeted samples, not limited to edible parts.