

OResults of Radioactive Material Monitoring of Aquatic Organisms (Location D along the Mano River)

<Location D along the Mano River: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
D-4 a	○	○	○	○	○	○

<Location D along the Mano River: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Transparency (cm)
D-4 a	37.7308°	140.9081°	2017/11/6	10:16	10:36	12.8	13.4	Gravel with sand	2.5Y4/3	None	0.67	>50

<Location D along the Mano River: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
D-4 a	37.7308°	140.9081°	2017/11/6	10:16	7.2	<0.5	3.1	11.3	7.7	0.05	1.5	2	1.5	0.0049	0.031	0.0011

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location D along the Mano River: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{SHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
D-4 a	37.7308°	140.9081°	2017/11/6	10:36	7.2	302	18.2	1.5	2.2	2.712	36.5	32.3	24.3	4.8	0.6	1.5	1.4	19	39	310	0.83

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location D along the Mano River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
D-3	The main stream of the Mano River	37.7051°	140.9623°	2017/10/20	Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	4	0.36	Immature fish	Japanese mitten Crab	Viscera removed	6.03	0.83	5.2	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Zacco platypus</i>	Pale chub	7	0.085	Immature fish,Mature fish	Algae	Viscera removed	33.7	3.7	30	-
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis</i>	Sweetfish	29	1.0	Immature fish,Mature fish	-	-	21.6	2.6	19	-
D-4b	The main stream of the Mano River	37.7312°	140.9096°	2017/10/20	Algae/plant	-	-	-	-	Riverbed Deposits (Include algae)	-	0.013	-	-	-	61.6	7.6	54	-
					Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp	230	0.044	Juvenile,Imago	-	-	32.4	4.4	28	-
					Mollusca	Bivalvia	Unionoida	Unionidae	<i>Inversium yokohamensis</i>	Inversium yokohamensis	20	0.084	Imago	-	Molluscos part	62.5	8.5	54	-
					Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	Semisulcospira libertina	30	0.023	Imago	-	Molluscos part	260	30	230	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	2	0.038	Immature fish	-	-	35.1	4.1	31	-
					Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	3	0.021	Immature fish,Mature fish	-	-	86.1	9.1	77	-
					Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.22	-	-	-	37.7	4.7	33	-
D-5	The main stream of the Mano River	37.7214°	140.8889°	2017/10/20	Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	6	0.47	Immature fish,Mature fish	Grasshopper, Kamimuria tibialis, Dragonfly(larva),Water strider, Baetidae,Moth,Antocha sp., Stenopsyche marmorata	Viscera removed	294	34	260	-
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	2	0.47	Immature fish	Obscure digesta	Viscera removed	202	22	180	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.