

Results of Radioactive Material Monitoring of Aquatic Organisms (Location H in Lake Akimoto)

<Location H in Lake Akimoto: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
H-1	○	○	○	○	○	○

<Location H in Lake Akimoto: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
H-1(Surface layer)	37.6575°	140.1264°	2017/10/18	08:35	09:10	14.5	12.5	Ooze	7.5Y 3/2	Plant pieces	13.8	4.5
H-1(Bottom layer)						13.8						

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
H-1(Surface layer)	37.6575°	140.1264°	2017/10/18	08:35	7.0	<0.5	3.2	8.9	5.1	0.03	1.3	2	1.4	N.D.(0.0019)	0.0095	-
H-1(Bottom layer)					6.9	1.0	3.3	8.4	5.2	0.03	1.7	2	2.3	0.0015	0.013	0.0010

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{N,H,E} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution							Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)	
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)				Maximum grain diameter (mm)
H-1	37.6575°	140.1264°	2017/10/18	09:10	6.7	203	69.1	9.6	21.5	2.564	0.0	0.0	0.1	0.7	57.5	41.7	0.0067	2.0	100	780	1.2

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)		
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
H-1 H-2 H-3	In the lake	37.6575° 37.6616° 37.6653°	140.1264° 140.1226° 140.1329°	2017/10/18	Arthropoda	Malacostraca	Decapoda	Astacidae	<i>Pacifastacus leniusculus trowbridgii</i>	Signal crayfish	11	0.64	Imago	-	-	-	21.5	3.5	18	7.5	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	12	2.2	Mature fish	-	-	-	49.2	5.2	44	0.71	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	<i>Carassius auratus langsdorffii</i>	Carassius auratus langsdorffii	6	0.80	Mature fish	-	-	-	42.1	5.1	37	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	<i>Hemibarbus barbus</i>	Hemibarbus barbus	3	2.0	Mature fish	Obscure digesta	Viscera removed	-	43.0	5.0	38	1.3
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Hypomesus nipponensis</i>	<i>Hypomesus nipponensis</i>	Japanese smelt	78	0.31	Immature fish, Mature fish	-	-	-	21.2	2.2	19	-
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	<i>Salvelinus leucomaenis</i>	Char	4	1.0	Mature fish	Japanese smelt	Viscera removed	-	33.3	3.3	30	0.30
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	<i>Oncorhynchus masou</i>	Seema	1	0.90	Mature fish	Empty stomach	Viscera removed	-	34.1	4.1	30	0.20
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus salmoides</i>	<i>Micropterus salmoides</i>	Largemouth bass	2	1.7	Immature fish, Mature fish	Japanese smelt	Viscera removed	-	47.4	5.4	42	1.5
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	<i>Micropterus dolomieu</i>	Small mouth bass	8	3.1	Immature fish, Mature fish	Signal crayfish, Japanese dace, Epophthalmia elegans	Viscera removed	-	91	10	81	0.48
Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Lepomis macrochirus</i>	<i>Lepomis macrochirus</i>	Bluegill	1	0.26	Mature fish	Elodea nuttallii	Viscera removed	-	22.9	2.9	20	-					
H-3	Inflowing rivers	37.6653°	140.1329°	2017/10/18	Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.23	-	-	-	37.9	3.9	34	-		
H-4	Within the lake and rivers in the vicinity	37.6551°	140.1181°	2017/10/18	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.026	-	-	-	N.D.	N.D.(1.5)	N.D.(1.3)	-		
					Algae/plant	Monocotyledoneae	Alismatales	Hydrocharitaceae	<i>Elodea nuttallii</i>	Western Waterweed	-	0.11	-	-	-	-	23.4	2.4	21	-	
					Arthropoda	Insecta	Plecoptera	Perlidae	<i>Oyamia sp.</i>	Oyamia	21	0.0040	Larva	-	-	-	N.D.	N.D.(10)	N.D.(8.7)	-	
					Arthropoda	Insecta	Odonata	Cordulegastriidae	<i>Anotogaster sieboldii</i>	<i>Anotogaster sieboldii</i>	-	-	-	-	-	-	-	-	-	-	-
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Nihonogomphus viridis</i>	<i>Nihonogomphus viridis</i>	26	0.022	Larva (Dragonfly larva)	-	-	-	12.9	1.9	11	-	
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Onychogomphus viridicostus</i>	<i>Melligomphus viridicostus</i>	-	-	-	-	-	-	-	-	-	-	
					Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	<i>Semisulcospira libertina</i>	30	0.029	Imago	-	Molluscos part	-	4.3	N.D.(1.6)	4.3	-	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Phoxinus lagowskii steindachneri</i>	<i>Phoxinus lagowskii steindachneri</i>	Amur Minnow	15	0.072	Immature fish, Mature fish	-	-	-	8.3	1.0	7.3	-
					Vertebrata	Amphibia	Anura	Ranidae	<i>Rana ornativentris</i>	<i>Rana ornativentris</i>	Montane brown frog	5	0.021	Imago	-	-	-	11	N.D.(2.0)	11	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.