

OResults of Radioactive Material Monitoring of Aquatic Organisms (Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J)

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Samples collected>

Items	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
Locations						
J-1	○	○	○	○	○	○

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Site measurement item>

Items	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
J-1(Surface layer)	37.4203°	140.1008°	2017/12/1	12:34	13:20	7.9	7.9	Sand gravel	7.5Y 5/3	Freshwater clam	3.5	>3.5
J-1(Bottom layer)												

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
	Latitude	Longitude	Date	Time (water)												
Locations						(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
J-1(Surface layer)	37.4203°	140.1008°	2017/12/1	12:34	7.1	<0.5	1.2	11.7	10.5	0.06	0.8	<1	0.3	N.D.(0.0014)	0.0070	-
J-1(Bottom layer)					6.9	<0.5	2.2	11.1	10.7	0.06	1.0	<1	0.5	N.D.(0.0014)	0.0072	0.00069

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: General survey items/Analysis of radioactive materials Sediment>

Items	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>NHLE</sub> (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
Locations																					
J-1	37.4203°	140.1008°	2017/12/1	13:20	6.1	341	28.5	1.5	2.3	2.740	2.2	5.1	47.7	37.4	4.8	2.8	0.27	9.5	7.6	55	N.D.(0.12)

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Lake Inawashiro (north lakeside) I / Lake Inawashiro (south lakeside) J: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
I-1 I-2 (north lakeside)	Within the lake and Nagase River	37.5047° 37.4995°	140.1143° 140.1409°	2017/12/1	Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.22	-	-	-	0.54	N.D.(0.48)	0.54	-
J-1 (south lakeside)	Within the lake and around the Oninuma	37.4203°	140.1008°	2017/12/1	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.021	-	-	-	N.D.	N.D.(1.6)	N.D.(1.6)	-
					Mollusca	Gastropoda	Architaenioglossa	Viviparidae	<i>Bellamyia japonica</i>	Japanese mysterysnail	5	0.017	Imago	-	Molluscous part	N.D.	N.D.(3.7)	N.D.(3.1)	-
					Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	84	0.038	Immature fish,Mature fish	Obscure digesta	Viscera removed	1.6	N.D.(1.7)	1.6	-
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Gymnogobius urotaenia</i>	Goby	8	0.012	Immature fish	-	-	9.3	N.D.(4.4)	9.3	-
					Vertebrata	Amphibia	Anura	-	-	Frog	132	0.055	Larva (Tadpole)	-	-	7.1	N.D.(1.6)	7.1	-

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.