OResults of Radioactive Material Monitoring of Aquatic Organisms (Location L off Soma City)

<Location L off Soma City: Samples collected>

Items	Genera	ıl items	Radioactive materials								
Locations	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)					
L-2	0	0	0	0	0	0					

<Location L off Soma City: Site measurement item>

Items	Latitude and le loca	ongitude of the		Survey date and time		Water		Sedi	ment		Ot	ther
Locations	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Property Color		Water depth (m)	Secchi disk depth (m)
L-2	37.8155°	140.9763°	2017/12/7	11:13	11:40	10.8	10.4	Sand	7.5Y4/3	None	1.1	>1.1

<Location L off Soma City: General survey items/Analysis of radioactive materials Water>

	Items	Latitude and longitude of the location		Survey dat	te and time	pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
	Locations	Latitude	Longitude	Date	Time (water)		(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
Ī	L-2	37.8155°	140.9763°	2017/12/7	11:13	8.1	<0.5	1.4	10.3	4830	32.24	1.0	<1	0.5	0.0012	0.0091	0.00081

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location L off Soma City: General survey items/Analysis of radioactive materials Sediment>

Latitude and longitude of the		ongitude of the	Survey do	ate and time										Grain siz	ze distribution						
itens	location		Sui vey da	ne and time	pH	Redox potential	Water content	IL	TOC	Soil particle	Gravel	Coarse sand	Medium sand	Fine sand	Silt	Clay	Median grain	Maximum	Cs-134	Cs-137	Sr-90
Locations	Latitude	Lanaituda	Data	Time (sediment)	1	$E_{N.H.E}$				density	(2-75mm)	(0.85-2mm)	(0.25-0.85mm)	(0.075-0.25mm)	(0.005-0.075mm)	(Less than 0.005mm)	diameter	grain diameter			
Locations	Latitude	Longitude	Date	Time (sediment)		(mV)	(%)	(%)	(mg/g-dry)	(g/cm ³)	(%)	(%)	(%)	(%)	(%)	(%)	(mm)	(mm)	(Bq/kg-dry)	(Bq/kg-dry)	(Bq/kg-dry)
L-2	37.8155°	140.9763°	2017/12/7	11:40	7.7	264	24.6	1.4	1.4	2.673	0.0	1.5	48.8	45.1	2.9	1.7	0.25	2.0	3.4	29	N.D.(0.14)

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location L off Soma City: Analysis items Aquatic organisms>

-Eocation E on Boi	na City. Anarysis item:																														
Locations	Sampling point	Latitude and longitude of the		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight	Note			Radioactive cesium (Bq/kg-wet)			Sr-90												
	bumping point	Latitude	Longitude	Sumpring date	Division	Class	Orac.	1	Selentine name	Linguist manie	ropulation	(kg-wet)	Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	(Bq/kg-wet)												
			140.06108	2017/12/3	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.013	-	ı	-	10	N.D.(2.8)	10	-												
				140.06100	140.06109	140.06100	140.06109	140.06100	140.06100	140.06109	140.06100	140.9610°	2017/12/7	Algae/plant	Monocotyledoneae	Najadales	Zosteraceae	Zostera marina	Eel grass	-	0.27	-	ı	-	0.67	N.D.(0.58)	0.67	-			
L-1		37.8210°												Arthropoda	Malacostraca	Mysida	Mysidae	Mysidae	Mysidae	-	0.29	Imago	-	-	0.72	N.D.(0.32)	0.72	-			
L-1 L-2	Matsukawaura	37.8210° 37.8155°	140.9763°	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	2017/12/3	Arthropoda	Malacostraca	Decapoda	Alpheidae	Alpheidae	Alpheidae	37	0.038	Juvenile,Imago	-	-	2.6	N.D.(1.4)	2.6	-
L-2 L-3	Lagoon	37.8217°		140.9765°						Arthropoda	Malacostraca	Decapoda	Palaemonidae	Palaemon macrodactylus	Palaemon macrodactylus	190	0.047	Juvenile,Imago	ı	-	1.9	N.D.(1.3)	1.9	-							
L.3	L-3		140.9703					Mollusca	Bivalvia	Ostreoida	Ostreidae	Crassostrea gigas	Oyster	13	0.31	Imago	1	Molluscous part	0.68	N.D.(0.39)	0.68	-									
				2017/12/4	Mollusca	Bivalvia	Veneroida	Veneridae	Ruditapes philippinarum	Japanese littleneck	50	0.26	Imago		Molluscous part	N.D.	N.D.(0.48)	N.D.(0.57)	-												
				2017/12/5	Vartabrata	Octoiobthyac	Saarmaanifarmas	Uavaarammidaa	Hoyagrammos otakii	Fat greenling	2	0.11	Immeture fich	Ragworm Shrimn	Viccore removed	1.2	N D (0.50)	1.2	_												

*1: Organisms were collected in or around the targeted water areas.

^{*2:} When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

 $^{{\}bf *3:} For a sample made of multiple types of a quatic organisms, the English name of the dominant one largest in number is underlined.$

^{*4:} Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

 $^{*5:} Plankton \ (suspended \ algae) \ is \ the \ residue \ remaining \ after \ the \ filtration \ of \ lake \ water \ or \ seawater \ with \ a \ plankton \ net \ (40\mu m-mesh).$

^{*6:} River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

^{*7:} N.D. means to be below the detection limit and figures in parentheses show the detection limit.

^{*8:} Activity concentrations include counting errors, but the details are omitted here.