

○ Results (water)

Locations			2018 May - June Survey												
	Latitude	Longitude	pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)	
Abukuma River System	A-1(Surface layer)	37.6210°	140.5218°	7.3	1.6	4.5	8.0	19.9	0.11	2.3	5	3.1	0.0017	0.015	0.0013
	A-1(Bottom layer)	37.6210°	140.5218°	7.3	1.7	4.7	8.3	19.9	0.11	2.4	8	3.6	N.D.(0.0016)	0.020	-
	A-2	37.5673°	140.3946°	7.3	0.8	3.6	9.8	13.1	0.07	1.5	4	2.5	0.0032	0.026	-
	B-2	37.8121°	140.5058°	7.4	1.0	3.4	9.5	9.9	0.06	1.3	8	3.3	0.0027	0.021	-
	B-3	37.8182°	140.4679°	7.4	0.9	3.0	10.2	5.6	0.03	1.1	8	2.9	N.D.(0.0015)	0.0096	-
Uda River	C-6	37.7764°	140.8877°	7.4	<0.5	2.9	9.2	11.0	0.06	1.3	3	2.8	0.0016	0.014	0.00061
Mano River	D-4 a	37.7308°	140.9081°	7.1	<0.5	2.6	9.5	10.8	0.06	1.1	2	1.8	0.0016	0.013	0.00075
Niida River	E-2 a	37.6640°	140.9447°	7.0	<0.5	2.4	9.7	8.1	0.05	1.0	3	2.4	0.0037	0.031	0.0015
Ota River	F-1	37.5975°	140.9252°	7.2	<0.5	2.7	9.7	5.2	0.03	1.1	1	0.8	0.015	0.14	0.0035
Lake Hayama (Mano Dam)	G-1(Surface layer)	37.7321°	140.8127°	7.2	0.5	3.6	9.0	7.0	0.04	1.6	5	4.1	0.0046	0.038	-
	G-1(Bottom layer)	37.7321°	140.8127°	6.9	<0.5	3.8	7.9	7.1	0.04	1.6	6	4.7	0.0042	0.039	0.0012
	G-4	37.7382°	140.8035°	7.5	<0.5	2.4	9.4	7.6	0.04	1.0	2	1.2	N.D.(0.0014)	0.013	-
Lake Akimoto	H-1(Surface layer)	37.6575°	140.1264°	7.3	0.7	2.4	9.7	3.8	0.03	1.2	<1	0.7	N.D.(0.0016)	0.0071	-
	H-1(Bottom layer)	37.6575°	140.1264°	7.1	0.8	3.1	9.9	3.6	0.03	1.5	1	1.1	N.D.(0.0014)	0.0043	0.0011
Lake Inawashiro	J-1(Surface layer)	37.4203°	140.1008°	6.5	<0.5	1.4	11.0	10.7	0.06	0.6	<1	0.7	N.D.(0.0013)	0.0066	-
	J-1(Bottom layer)	37.4203°	140.1008°	6.6	1.1	1.8	10.7	10.8	0.06	0.9	1	0.8	N.D.(0.0016)	0.0062	0.00086
Off the mouth of the Abukuma River (Sea Area in front of the mouth of the Abukuma River)	K-3(Surface layer)	38.0458°	140.9518°	8.2	2.2	4.5	9.8	4650	30.84	2.2	3	1.9	N.D.(0.0017)	0.0056	-
	K-3(Bottom layer)	38.0458°	140.9518°	7.9	0.7	1.8	3.3	5050	34.12	1.3	4	1.9	0.0014	0.014	0.00081
Off Soma City (Matsukawaura)	L-2	37.8155°	140.9763°	8.1	0.9	3.2	8.5	4640	31.69	1.9	6	3.5	0.0025	0.031	0.00081
Off Iwaki City (Hisanohama)	M-2(Surface layer)	37.1996°	141.0853°	8.1	<0.5	1.6	8.7	5070	34.12	1.0	1	0.6	N.D.(0.0014)	0.0038	-
	M-2(Bottom layer)	37.1996°	141.0853°	8.1	<0.5	1.2	6.0	5090	34.08	0.9	2	0.5	N.D.(0.0015)	0.0038	0.00098

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

O Results (sediments)

Locations			2018 May - June Survey																	
			pH	Redox potential E _{NHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution							Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)		
									Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)				
Abukuma River System	A-1	37.6210°	140.5218°	7.0	43	30.8	3.9	5.5	2.673	0.0	1.6	62.6	12.8	10.4	12.6	0.35	4.8	30	290	0.17
	A-2	37.5673°	140.3946°	6.9	317	18.8	2.2	2.1	2.715	25.8	46.4	19.6	5.5	2.7		1.3	9.5	15	140	-
	B-2	37.8121°	140.5058°	7.1	332	24.4	2.0	1.6	2.772	0.1	0.8	62.3	34.3	2.5		0.29	4.8	12	130	-
	B-3	37.8182°	140.4679°	7.3	312	18.2	1.9	2.2	2.673	18.7	26.7	34.8	15.7	1.4	2.7	0.73	4.8	9.0	91	-
Uda River	C-6	37.7764°	140.8877°	7.2	283	17.0	1.8	1.5	2.728	27.6	40.6	28.8	2.3	0.7		1.2	4.8	13	130	0.39
Mano River	D-4 a	37.7308°	140.9081°	7.2	263	17.3	2.4	1.4	2.714	31.3	43.9	20.9	2.7	1.2		1.4	19	30	310	0.82
Niida River	E-2 a	37.6640°	140.9447°	7.0	122	21.9	3.0	4.5	2.693	15.2	15.0	36.1	19.0	6.6	8.1	0.38	19	150	1500	0.36
Ota River	F-1	37.5975°	140.9252°	7.1	331	19.9	1.6	1.6	2.655	9.7	33.9	45.5	8.4	2.5		0.75	4.8	170	1600	0.69
Lake Hayama (Mano Dam)	G-1	37.7321°	140.8127°	6.9	91	60.8	11.7	43.8	2.552	0.0	0.1	1.7	24.5	40.0	33.7	0.020	2.0	320	3000	3.4
	G-4	37.7382°	140.8035°	7.2	210	25.7	3.2	3.6	2.692	9.9	24.1	43.7	12.6	3.8	5.9	0.59	19	62	630	-
Lake Akimoto	H-1	37.6575°	140.1264°	6.8	75	61.0	9.2	17.9	2.581	0.0	0.1	0.2	0.7	52.4	46.6	0.0058	2.0	29	250	1.5
Lake Inawashiro	J-1	37.4203°	140.1008°	7.2	314	24.0	1.9	2.5	2.719	0.3	1.7	56.9	33.9	2.6	4.6	0.28	4.8	7.0	80	N.D.(0.13)
Off the mouth of the Abukuma River (Sea Area in front of the mouth of the Abukuma River)	K-3	38.0458°	140.9518°	7.7	12	34.2	5.4	5.1	2.698	0.0	0.1	0.1	29.6	56.0	14.2	0.053	2.0	19	190	N.D.(0.13)
Off Soma City (Matsukawaura)	L-2	37.8155°	140.9763°	7.7	204	19.8	1.7	0.9	2.724	1.6	6.8	81.2	8.6	1.8		0.43	9.5	0.50	5.0	N.D.(0.14)
Off Iwaki City (Hisanohama)	M-2	37.1996°	141.0853°	7.6	188	21.7	2.9	1.4	2.789	0.0	0.6	2.4	91.1	2.6	3.3	0.16	2.0	4.3	50	N.D.(0.12)

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

O Results (aquatic organisms)

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
Abukuma River System	A-2	Harase River	2018/5/30	Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0072	-	-	-	79.5	5.5	74	-	
				Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	<i>Stenopsyche marmorata</i>	60	0.021	Larva	-	-	33.2	3.2	30	-
				Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	<i>Macromia amphigena</i>	94	0.11	Larva(Dragonfly larva)	-	-	18.1	2.1	16	-
				Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	<i>Anotogaster sieboldii</i>									
				Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	<i>Sieboldius albardae</i>									
				Arthropoda	Insecta	Odonata	Gomphidae	<i>Asiagomphus melaeonops</i>	<i>Asiagomphus melaeonops</i>									
				Arthropoda	Malacostraca	Decapoda	Cambaridae	<i>Procambarus clarkii</i>	Red swamp crawfish	6	0.029	Juvenile,Imago	-	-	14.8	1.8	13	-
				Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Neocaridina sp.</i>	<i>Neocaridina</i>	357	0.11	Juvenile,Imago	-	-	14.4	1.4	13	-
				Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	<i>Semisulcospira libertina</i>	30	0.031	Imago	-	Molluscous part	154	14	140	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Phoxinus lagowskii steindachneri</i>	<i>Amur Minnow</i>	226	1.1	Immature fish,Mature fish	-	-	7.45	0.95	6.5	0.23
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	<i>Japanese dace</i>	14	0.16	Immature fish	-	-	9.64	0.94	8.7	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	<i>Pale chub</i>	6	0.058	Immature fish	-	-	7.2	N.D.(0.85)	7.2	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Pseudogobio esocinus esocinus</i>	<i>Pseudogobio esocinus esocinus</i>	4	0.054	Immature fish,Mature fish	-	-	6.2	N.D.(1.1)	6.2	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Candidia temminckii</i>	<i>Dark chub</i>	18	0.067	Immature fish	-	-	7.3	1.2	6.1	-
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	<i>Oriental weatherfish</i>	99	0.38	Immature fish,Mature fish	-	-	9.8	1.1	8.7	-
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Nemacheilus toni</i>	<i>Stone loach</i>	71	0.91	Immature fish	-	-	6.96	0.76	6.2	0.16
				Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	<i>Yamame trout</i>	1	0.092	Mature fish	Diptera,Ant,Aquatic insect	Viscera removed	10	N.D.(0.83)	10	-
				Vertebrata	Amphibia	Anura	-	-	Frog	28	0.014	Larva(Tadpole)	-	-	134	14	120	-
				Vertebrata	Amphibia	Anura	Ranidae	<i>Rana japonica</i>	<i>Japanese Brown Frog</i>	7	0.040	Imago	-	-	19.7	1.7	18	-
				Vertebrata	Amphibia	Anura	Glandirana	<i>Glandirana rugosa</i>	<i>Wrinkled Frog</i>									
				Vertebrata	Amphibia	Caudata	Salamandridae	<i>Cynops pyrrhogaster</i>	<i>Cynops pyrrhogaster</i>	11	0.068	Imago	-	-	6.78	0.68	6.1	-
				Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.18	-	-	-	32.3	3.3	29	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
Abukuma River System	B-2 The main stream of the Abukuma River	2018/5/17	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	27	7.8	Mature fish	Obscure digesta	Viscera removed	13.2	1.2	12	0.17
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	3	4.7	Mature fish	Obscure digesta	Viscera removed	28.7	2.7	26	0.28
			Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	3	5.2	Mature fish	Gomphidae,Crab	Viscera removed	28.0	3.0	25	0.22
			Vertebrata	Osteichthyes	Siluriformes	Ictaluridae	<i>Ictalurus punctatus</i>	Channel catfish	1	4.2	Mature fish	Small mammals,Crab	Viscera removed	10.6	1.1	9.5	0.34
			Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	1	1.7	Mature fish	Empty stomach	Viscera removed	40.9	3.9	37	0.27
	B-3 Surikami River	2018/5/28	Algae/plant	-	-	-	-	Riverbed Deposits (Include algae)	-	0.012	-	-	-	40.1	4.1	36	-
			Arthropoda	Insecta	Ephemeroptera	Siphlonuridae	<i>Siphlonuridae</i>	<i>Siphlonuridae</i>	402	0.022	Larva	-	-	29.1	2.1	27	-
			Arthropoda	Insecta	Ephemeroptera	Ephemeridae	<i>Ephemerella strigata</i>	Mont mayfly									
			Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata	405	0.14	Larva	-	-	24.4	2.4	22	-
			Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	Macromia amphigena	46	0.032	Larva(Dragonfly larva)	-	-	7.7	N.D.(1.4)	7.7	-
			Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	<i>Anotogaster sieboldii</i>				-	-				
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Melligomphus viridicostus</i>	<i>Melligomphus viridicostus</i>									
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	<i>Sieboldius albardae</i>									
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Davidius sp.</i>	<i>Davidius</i>									
			Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	<i>Protohermes grandis</i>									
			Arthropoda	Malacostraca	Decapoda	Cambaridae	<i>Procambarus clarkii</i>	Red swamp crawfish	44	0.051	Larva	-	-	1.6	N.D.(1.2)	1.6	-
			Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	14	0.15	Juvenile,Imago	-	-	11.2			
			Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus pollux</i>	Japanese fluvial sculpin	4	1.3	Immature fish,Mature fish	Amur minnow,Cobitis biwae	Viscera removed	13.2	1.2	12	0.12
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Phoxinus lagowskii steindachneri</i>	Amur Minnow	43	0.29	Immature fish	-	-	4.26	0.56	3.7	-
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	194	2.4	Immature fish,Mature fish	-	-	4.38	0.48	3.9	0.36
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Pseudogobio esocinus esocinus</i>	Pseudogobio esocinus esocinus	31	0.24	Immature fish	-	-	5.12	0.52	4.6	-
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Candidia temminckii</i>	Dark chub	2	0.051	Mature fish	-	-	4.1	N.D.(0.95)	4.1	-
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp	15	0.17	Immature fish,Mature fish	-	-	5.27			
			Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	1	1.6	Mature fish	Obscure digesta	Viscera removed	8.84	0.94	7.9	0.37
			Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Nemacheilus toni</i>	Stone loach	49	0.20	Immature fish,Mature fish	-	-	9.6	1.0	8.6	-
			Vertebrata	Osteichthyes	Siluriformes	Bagridae	<i>Tachysurus tokiensis</i>	Cut-tailed bullhead	133	1.2	Immature fish	-	-	4.49	0.39	4.1	0.27
			Vertebrata	Amphibia	Anura		Frog	6	0.072	Immature fish,Mature fish	-	-	3.5	N.D.(0.90)	3.5	-	
			Vertebrata	Amphibia	Anura	Lithobates	<i>Lithobates catesbeianus</i>	American Bullfrog	8	0.011	Larva(Tadpole)	-	-	82.8			
			Vertebrata	Amphibia	Anura	Glandirana	<i>Glandirana rugosa</i>	Wrinkled Frog	1	0.55	Imago	-	-	7.24	0.74	6.5	-
			Vertebrata	Amphibia	Anura	Pelophylax	<i>Pelophylax porosus porosus</i>	Tokyo Daruma pond frog	8	0.030	Imago	-	-	20.9	1.9	19	-
			Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.21	-	-	-	13.1	1.1	12	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
Uda River	C-6	The main stream of the Uda River	2018/6/2	Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0088	-	-	-	49.0	5.0	44	-	
				Algae/plant	Zygnematophyceae	Zygnematales	Zygnemataceae	<i>Spirogyra sp.</i>	<i>Spirogyra</i>	-	0.17	-	-	-	2.5	N.D.(0.42)	2.5	-
				Arthropoda	Insecta	Ephemeroptera	Isonychiidae	<i>Isonychia valida</i>	<i>Isonychia valida</i>	330	0.014	Larva	-	-	40.9	3.9	37	-
				Arthropoda	Malacostraca	Decapoda	Cambaridae	<i>Procambarus clarkii</i>	Red swamp crawfish	35	0.054	Juvenile,Imago	-	-	5.8	N.D.(1.2)	5.8	-
				Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp	150	0.031	Juvenile	-	-	5.3	N.D.(1.8)	5.3	-
				Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	9	0.19	Juvenile	-	-	13.4	1.4	12	-
				Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	2	1.1	Mature fish	Empty stomach	Viscera removed	26.9	1.9	25	0.069
				Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	3	0.90	Immature fish,Mature fish	Japanese mitten Crab	Viscera removed	26.0	3.0	23	0.072
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	1	0.0097	Immature fish	Obscure digesta	Viscera removed	7.7	N.D.(5.8)	7.7	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Candidia temminckii</i>	Dark chub	17	0.069	Immature fish,Mature fish	Obscure digesta	Viscera removed	5.6	N.D.(0.96)	5.6	-
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Cobitis biwae</i>	Cobitis biwae	14	0.040	Mature fish	-	-	5.0	N.D.(1.3)	5.0	-
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	34	0.12	Immature fish,Mature fish	-	-	4.50	0.50	4.0	-
				Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	83	1.1	Immature fish,Mature fish	-	-	23.2	2.2	21	0.10
				Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	1	0.16	Immature fish	Common prawn,Plant pieces	Viscera removed	22.0	2.0	20	-
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Gymnogobius urotaenia</i>	Goby	4	0.040	Immature fish	Ephemeroptera,Plec optera,Algae,Plant pieces	Viscera removed	15.3	1.3	14	-
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	34	0.059	Immature fish,Mature fish	Ephemeroptera	Viscera removed	13.5	1.5	12	-
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius nagoyae			Dusky tripletooth goby	Obscure digesta	Viscera removed	10.5	1.4	9.1	-
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Tridentiger brevipinnis</i>	Cut-tailed bullhead	2	0.046	Mature fish	Obscure digesta	Viscera removed	6.7	N.D.(2.6)	6.7	-
				Vertebrata	Osteichthyes	Siluriformes	Bagridae	<i>Tachysurus tokiensis</i>	Tokyo Daruma pond frog	1	0.013	Imago	-	-	N.D.	N.D.(3.0)	N.D.(2.7)	-
				Coarse Particulate Organic Matter	-	-	-	<i>Pelophylax porosus porosus</i>	Bottom fallen leaves	-	0.22	-	-	-	87.8	9.8	78	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)					
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
Mano River	D-3 The main stream of the Mano River	2018/6/1	Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	33	0.60	Immature fish,Mature fish	Attached algae	Viscera removed	25.9	1.9	24	-		
				Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0076	-	-	-	124	14	110	-		
				Zygnematophyceae	Zygnematales	Zygnemataceae	<i>Spirogyra sp.</i>	Spirogyra	-	0.28	-	-	-	30.0	2.0	28	-		
				Monocotyledoneae	Poales	Poaceae	<i>Phragmites australis</i>	Common reed	-	0.20	-	-	-	23.7	2.7	21	-		
				Monocotyledoneae	Najadales	Potamogetonaceae	<i>Potamogeton berchtoldii</i>	Small pondweed	-	0.24	-	-	-	188	18	170	-		
				Arthropoda	Insecta	Ephemeroptera	<i>Potamanthus formosus</i>	Potamanthus formosus	504	0.023	Larva	-	-	31.2	3.2	28	-		
				Arthropoda	Insecta	Ephemeroptera	<i>Isonychia valida</i>	Isonychia valida											
				Arthropoda	Insecta	Ephemeroptera	<i>Siphlonuridae</i>	<i>Siphlonuridae</i>											
				Arthropoda	Insecta	Ephemeroptera	<i>Ephemerellidae</i>	<i>Drunella ishiyamana</i>											
				Arthropoda	Insecta	Ephemeroptera	<i>Ephemeridae</i>	<i>Ephemeridae</i>											
				Arthropoda	Insecta	Trichoptera	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata			119	0.010	Larva	-	-	34	N.D.(4.3)	34	-
				Arthropoda	Insecta	Odonata	<i>Macromia amphigena amphigena</i>	Macromia amphigena											
				Arthropoda	Insecta	Odonata	<i>Cordulegastridae</i>	<i>Anotogaster sieboldii</i>											
				Arthropoda	Insecta	Odonata	<i>Gomphidae</i>	<i>Stylgomphus suzukii</i>											
				Arthropoda	Insecta	Odonata	<i>Melligomphus viridicostus</i>	Melligomphus viridicostus											
				Arthropoda	Insecta	Odonata	<i>Gomphidae</i>	<i>Sieboldius albardae</i>											
				Arthropoda	Insecta	Odonata	<i>Davidius sp.</i>	<i>Davidius</i>											
				Arthropoda	Insecta	Odonata	<i>Aeshnidae</i>	<i>Boyeria macchlani</i>											
				Arthropoda	Insecta	Megaloptera	<i>Corydalidae</i>	<i>Protohermes grandis</i>											
				Arthropoda	Malacostraca	Decapoda	<i>Cambaridae</i>	<i>Procambarus clarkii</i>	Red swamp crawfish	2	0.056	Imago	-	-	59.0	5.0	54	-	
				Arthropoda	Malacostraca	Decapoda	<i>Palaemonidae</i>	<i>Palaemon paucidens</i>	Common prawn	24	0.040	Imago	-	-	13.4	1.4	12	-	
				Arthropoda	Malacostraca	Decapoda	<i>Atyidae</i>	<i>Paratya improvisa</i>	Freshwater shrimp	170	0.047	Juvenile,Imago	-	-	18.2	2.2	16	-	
				Arthropoda	Malacostraca	Decapoda	<i>Varunidae</i>	<i>Eriocheir japonica</i>	Japanese mitten crab	8	0.15	Juvenile	-	-	22.1	2.1	20	-	
	D-4 b The main stream of the Mano River	2018/6/6	Mollusca	Bivalvia	Unionoidea	Unionidae	<i>Inversuum jokohamensis</i>	<i>Inversuum jokohamensis</i>	4	0.025	Imago	-	-	65.7	6.7	59	-		
				Mollusca	Gastropoda	Discopoda	<i>Pleuroceridae</i>	<i>Semisulcospira libertina</i>	Semisulcospira libertina	30	0.031	Imago	-	-	65.6	5.6	60	-	
				Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	4	1.1	Immature fish,Mature fish	Japanese dace,Cobitis biwae,Sieboldius albardae,Ephemericlidiae	Viscera removed	110	10	100	0.13	
				Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus pollux</i>	Japanese fluvial sculpin	25	1.3	Immature fish,Mature fish	Protohermes grandis,Mataeopsephe nus japonicus,Freshwater shrimp,Stenopsyche marmorata,Rhinogobius s	Viscera removed	39.0	5.0	34	0.50	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Phoxinus lagowskii steindachneri</i>	Amur Minnow	17	0.071	Immature fish,Mature fish	-	-	12.2	1.2	11	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	5	0.026	Immature fish	-	-	22.1	2.1	20	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	Pale chub	17	0.14	Immature fish	Obscure digesta	Viscera removed	18.8	1.8	17	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Gnathopogon elongatus elongatus</i>	Gnathopogon elongatus elongatus	2	0.0076	Immature fish	-	-	14	N.D.(6.0)	14	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Cobitis biwae</i>	Cobitis biwae	79	0.29	Mature fish	Obscure digesta	Viscera removed	19.1	2.1	17	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	25	0.23	Immature fish,Mature fish	Obscure digesta	Viscera removed	14.2	1.2	13	-	
				Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	11	0.058	Immature fish	Attached algae	Viscera removed	49.7	5.7	44	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	67	0.21	Mature fish	Obscure digesta	Viscera removed	24.3	2.3	22	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius nagoyae			Mature fish	Obscure digesta	Viscera removed	20.6	2.6	18	-	
				Vertebrata	Cephalaspidomorphi	Petromyzontiformes	Petromyzontidae	<i>Tridentiger brevispinis</i>	Dusky tripletooth goby	7	0.14	Mature fish	-	-	11	N.D.(4.4)	11	-	
				Vertebrata	Amphibia	Anura	Rhacophoridae	<i>Buergeria buergeri</i>	Kajika frog	1	0.0032	Imago	-	-	175	15	160	-	
				Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.21	-	-	-	48.7	4.7	44	-	

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)					
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
Niida River E-2 b	The main stream of the Niida River 2018/6/2	2018/6/2	Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0094	-	-	-	-	226	26	200	-		
				Algae/plant	Monocotyledoneae	Poales	Poaceae	<i>Phragmites australis</i>	Common reed	-	0.17	-	-	-	22.3	2.3	20	-	
				Arthropoda	Insecta	Trichoptera	Stenoplectidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata	89	0.015	Larva	-	-	105	10	95	-	
				Arthropoda	Malacostraca	Decapoda	Cambaridae	<i>Procambarus clarkii</i>	Red swamp crawfish	9	0.081	Juvenile,Imago	-	-	56.4	5.4	51	-	
				Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Common prawn	44	0.079	Imago	-	-	25.6	2.6	23	-	
				Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp	353	0.078	Juvenile,Imago	-	-	39.4	3.4	36	-	
				Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	8	0.037	Juvenile	-	-	40.5	4.5	36	-	
				Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	Semisulcospira libertina	30	0.022	Imago	-	-	Molluscous part	75	13	62	-
				Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	1	0.42	Mature fish	Empty stomach	Viscera removed	92.7	8.7	84	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	20	0.23	Immature fish,Mature fish	-	-	30.9	2.9	28	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	Pale chub	10	0.11	Immature fish	-	-	39.8	3.8	36	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Pseudogobio esocinus esocinus</i>	Pseudogobio esocinus esocinus	17	0.44	Immature fish,Mature fish	-	-	26.4	1.4	25	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Sarcocheilichthys variegatus variegatus</i>	Sarcocheilichthys variegatus variegatus	5	0.047	Immature fish,Mature fish	-	-	21.5	2.5	19	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	1	0.060	Mature fish	Obscure digesta	Viscera removed	26.5	2.5	24	-	
				Vertebrata	Osteichthyes	Cypriniformes	Cobitidae	<i>Misgurnus anguillicaudatus</i>	Oriental weatherfish	3	0.037	Mature fish	-	-	31.7	2.7	29	-	
				Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	1	0.093	Immature fish	Orthoptera,Ecdyonurus yoshidae,Siphlonuridae,Apatania,Simuli um vittatum	Viscera removed	30.4	3.4	27	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	5	0.038	Mature fish	-	-	51.1	6.1	45	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius nagoyae										
				Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	1	1.4	Mature fish	Red swamp crawfish	Viscera removed	130	10	120	1.0	
				Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	4	2.3	Mature fish	Freshwater shrimp,Plant pieces	Viscera removed	83.3	7.3	76	0.71	
				Vertebrata	Amphibia	Anura	-	-	Frog	22	0.013	Larva(Tadpole)	-	-	287	27	260	-	
				Vertebrata	Amphibia	Anura	Lithobates	<i>Lithobates catesbeianus</i>	American Bullfrog	2	0.90	Imago	-	-	32.8	4.8	28	0.69	
				Vertebrata	Amphibia	Anura	Glandirana	<i>Glandirana rugosa</i>	Wrinkled Frog	5	0.028	Imago	-	-	33.0	3.0	30	-	
				Vertebrata	Amphibia	Anura	Pelophylax	<i>Pelophylax porosus porosus</i>	Tokyo Daruma pond frog										
				Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.25	-	-	-	86.9	7.9	79	-	
E-3	The main stream of the Niida River	2018/7/1	Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	143	3.1	Immature fish,Mature fish	-	-	-	47.9	4.9	43	0.25	

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)					
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
Ota River	The main stream of the Ota River	2018/6/5	Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0062	-	-	-	-	403	43	360	-		
			Algae/plant	Zygnematophyceae	Zygnematales	Zygnemataceae	<i>Spirogyra sp.</i>	Spirogyra	-	0.35	-	-	-	56.8	5.8	51	-		
		2018/6/7	Algae/plant	Monocotyledoneae	Poales	Poaceae	<i>Phragmites australis</i>	Common reed	-	0.10	-	-	-	231	21	210	-		
			Algae/plant	Monocotyledoneae	Najadales	Potamogetonaceae	<i>Potamogeton berchtoldii</i>	Small pondweed	-	0.17	-	-	-	46.5	3.5	43	-		
			Arthropoda	Insecta	Ephemeroptera	Isonychiidae	<i>Isonychia valida</i>	Isonychia valida	113	0.013	Larva	-	-	144	14	130	-		
			Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata	44	0.014	Larva	-	-	134	14	120	-		
			Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	Macromia amphigena	70	0.033	Larva(Dragonfly larva)	-	-	147	17	130	-		
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Nihonogomphus viridis</i>	Nihonogomphus viridis											
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Stylogomphus suzukii</i>	Stylogomphus suzukii											
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	Sieboldius albardae											
			Arthropoda	Insecta	Odonata	Gomphidae	<i>Asiagomphus melaenops</i>	Asiagomphus melaenops											
			Arthropoda	Insecta	Odonata	Libellulidae	<i>Orthetrum japonicum</i>	Japanese skimmer											
			Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	Protohermes grandis			44	0.0073	Larva	-	-	79.6	7.6	72	-
			Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Parachauliodes japonicus</i>	Parachauliodes japonicus											
			Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Common prawn	20	0.029	Imago	-	-	224	24	200	-		
			Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp	134	0.038	Juvenile,Imago	-	-	106.2	7.2	99	-		
			Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	5	0.085	Juvenile	-	-	306	26	280	-		
			Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	2	0.47	Immature fish,Mature fish	Obscure digesta	Viscera removed	641	61	580	-		
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	10	0.029	Immature fish	-	-	371	41	330	-		
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	Pale chub	7	0.037	Immature fish	-	-	240	20	220	-		
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius cuvieri</i>	Japanese crucian carp	2	1.9	Mature fish	Obscure digesta	Viscera removed	433	43	390	2.1		
			Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	19	0.067	Mature fish	Diptera(imago),Isonychia japonica,Ephemerella cryptomeria	Viscera removed	414	44	370	-		
			Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.11	-	-	-	709	69	640	-		
F-5	The main stream of the Ota River	2018/6/5	Vertebrata	Osteichthyes	Anguilliformes	Anguillidae	<i>Anguilla japonica</i>	Japanese eel	3	0.87	Immature fish,Mature fish	Obscure digesta	Viscera removed	254	24	230	-		
			Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	204	0.35	Immature fish	-	-	90.8	8.8	82	0.31		

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
Lake Hayuma	G-1 G-2 G-3	In the lake	2018/6/1	Algae/plant	-	-	-	Plankton (Planktonic algae)	-	0.015	-	-	-	2.3	N.D.(2.6)	2.3	-	
			2018/6/4	Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	4	2.6	Mature fish	Fish	Viscera removed	142	12	130	0.91
				Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Lepomis macrochirus</i>	Bluegill	41	0.20	Immature fish	Obscure digesta	Viscera removed	38.4	3.4	35	-
				Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0053	-	-	-	119.7	9.7	110	-	
				Arthropoda	Insecta	Ephemeroptera	Siphlonuridae	<i>Siphlonuridae</i>	Siphlonuridae	1214	0.020	Larva	-	-	39.0	4.0	35	-
				Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	Macromia amphigena	47	0.014	Larva(Dragonfly larva)	-	18	N.D.(3.1)	18	-	
				Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	Anotogaster sieboldii									
				Arthropoda	Insecta	Odonata	Gomphidae	<i>Stylogomphus suzukii</i>	Stylogomphus suzukii									
				Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	Sieboldius albardae									
				Arthropoda	Insecta	Odonata	Gomphidae	<i>Davidius sp.</i>	<i>Davidius</i>									
				Arthropoda	Insecta	Odonata	Aeshnidae	<i>Boyeria macclachlani</i>	Boyeria macclachlani									
				Arthropoda	Malacostraca	Decapoda	Atyidae	<i>Paratya improvisa</i>	Freshwater shrimp			Juvenile,Imago	-	-	25.5	2.5	23	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Phoxinus lagowskii steindachneri</i>	Amur Minnow	15	0.026	Immature fish	-	-	33.6	2.6	31	-
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	138	1.3	Immature fish,Mature fish	-	-	69.5	6.5	63	0.29
				Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Hypomesus nipponensis</i>	Japanese smelt	75	0.060	Immature fish,Mature fish	-	-	77.6	8.6	69	-
	G-4	Inflowing rivers	2018/6/1	Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	10	0.18	Immature fish	Scarabaeidae,Leaf beetle,Tiger beetle,Lepidoptera(larva),Lepidostoma,Goera japonica,Stink bug,Apatania,Ostrovus,Eriocnemis	Viscera removed	30.1	3.1	27	-
				Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	4	0.37	Mature fish	<i>Sieboldius albardae</i> ,Japanese dace,Rhinogobius flumineus	Viscera removed	217	17	200	1.1
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius flumineus</i>	Rhinogobius flumineus	51	0.059	Immature fish,Mature fish	Midge,Simulium vittatum	Viscera removed	29.4	3.4	26	-
				Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	3	2.1	Immature fish,Mature fish	Fish	Viscera removed	331	31	300	0.39
				Vertebrata	Amphibia	Anura	-	-	Frog	8	0.0022	Larva(Tadpole)	-	-	386	56	330	-
				Vertebrata	Amphibia	Anura	Ranidae	<i>Rana ornativentris</i>	Montane brown frog	2	0.024	Imago	-	-	29.9	2.9	27	-
				Vertebrata	Amphibia	Caudata	Salamandridae	<i>Cynops pyrrhogaster</i>	Cynops pyrrhogaster	1	0.0041	Imago	-	-	20	N.D.(15)	20	-
				Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.25	-	-	-	144	14	130	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
H-1 H-2 H-3	In the lake	2018/5/30		Arthropoda	Malacostraca	Decapoda	Astacidae	<i>Pacifastacus leniusculus trowbridgii</i>	Signal crayfish	10	0.88	Imago	-	-	15.9	1.9	14	7.5
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	12	3.7	Mature fish	Obscure digesta	Viscera removed	43.0	4.0	39	0.74
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	13	2.1	Mature fish	Obscure digesta	Viscera removed	42.6	2.6	40	1.1
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp	1	0.33	Mature fish	Obscure digesta	Viscera removed	31.1	3.1	28	1.1
				Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	3	2.2	Mature fish	Obscure digesta	Viscera removed	20.5	1.5	19	1.0
				Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Hypomesus nippensis</i>	Japanese smelt	47	0.27	Mature fish	-	-	19.7	1.7	18	-
				Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	6	1.8	Mature fish	Terrestrial insect,Japanese smelt	Viscera removed	52.1	5.1	47	0.27
				Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou masou</i>	Seema	3	0.62	Immature fish	Terrestrial insect,Japanese smelt,Mont mayfly	Viscera removed	42.5	4.5	38	-
				Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	4	0.65	Immature fish	Japanese smelt,Signal crayfish,Mont mayfly	Viscera removed	42.7	3.7	39	-
Lake Akimoto	Inflowing rivers	2018/5/30		Arthropoda	Insecta	Ephemeroptera	Heptageniidae	<i>Heptageniidae</i>	Heptageniidae	147	0.013	Larva	-	-	6.1	N.D.(3.0)	6.1	-
				Arthropoda	Insecta	Ephemeroptera	Siphlonuridae	<i>Siphlonuridae</i>	Siphlonuridae				-	-	-	-	-	-
				Arthropoda	Insecta	Ephemeroptera	Ephemerellidae	<i>Drunella basalis</i>	Drunella basalis				-	-	-	-	-	-
				Arthropoda	Insecta	Ephemeroptera	Ephemeridae	<i>Ephemerina japonica</i>	Ephemerina japonica				-	-	-	-	-	-
				Arthropoda	Insecta	Plecoptera	Perlidae	<i>Acroneuria sp.</i>	Acroneuria	86	0.016	Larva	-	-	N.D.	N.D.(2.6)	N.D.(2.5)	-
				Arthropoda	Insecta	Plecoptera	Perlidae	<i>Calineuria sp.</i>	Calineuria				-	-	-	-	-	-
				Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata	96	0.026	Larva	-	-	2.9	N.D.(1.6)	2.9	-
				Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus pollux</i>	Japanese fluvial sculpin	16	0.078	Immature fish	Ameletus montanus,Trichopter a,Drunella basalis,Acroneuria, Midge,Ephemeropte ra,Simuliidae	Viscera removed	4.1	N.D.(3.9)	4.1	-
				Vertebrata	Amphibia	Anura	Rhacophoridae	<i>Buergeria buergeri</i>	Kajika frog	19	0.14	Imago	-	-	69.2	6.2	63	-
				Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.17	-	-	-	18.5	1.5	17	-
H-4	Within the lake and rivers in the vicinity	2018/5/30		Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.015	-	-	-	N.D.	N.D.(2.0)	N.D.(2.2)	-
				Arthropoda	Insecta	Ephemeroptera	Ephemeridae	<i>Ephemerina japonica</i>	Ephemerina japonica	395	0.031	Larva	-	-	14.9	1.9	13	-
				Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	Anotogaster sieboldii	55	0.052	Larva(Dragonfly larva)	-	-	26.7	2.7	24	-
				Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Common prawn	55	0.017	Juvenile,Imago	-	-	13	N.D.(2.2)	13	-
				Mollusca	Gastropoda	Discopoda	Pleuroceridae	<i>Semisulcospira libertina</i>	Semisulcospira libertina	30	0.028	Imago	-	Molluscous part	11.3	1.3	10	-
				Vertebrata	Amphibia	Anura	Ranidae	<i>Rana ornativentris</i>	Montane brown frog	7	0.023	Imago	-	-	25.3	2.3	23	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
I-1 I-2 (north lakeside)	Within the lake and Nagase River	2018/5/30	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	12	2.3	Mature fish	Obscure digesta	Viscera removed	28.9	2.9	26	0.23
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	58	2.5	Immature fish,Mature fish	-	-	16.0	1.0	15	0.21
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	8	2.4	Mature fish	Obscure digesta	Viscera removed	34.8	2.8	32	0.39
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	2	2.7	Mature fish	Empty stomach	Viscera removed	38.3	3.3	35	0.35
			Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	4	1.3	Mature fish	Empty stomach	Viscera removed	39.4	3.4	36	0.11
			Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropodus dolomieu</i>	Small mouth bass	4	2.6	Immature fish,Mature fish	Common prawn	Viscera removed	27.8	2.8	25	0.23
			Vertebrata	Osteichthyes	Perciformes	Actinopterygii	<i>Channa argus</i>	Snakehead	2	1.6	Immature fish	Empty stomach	Viscera removed	19.4	1.4	18	0.30
			Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	1	0.62	Mature fish	Empty stomach	Viscera removed	22.8	1.8	21	-
	2018/5/29	Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.19	-	-	-	-	8.52	0.82	7.7	-
Lake Inawashiro	J-1 (south lakeside)	Within the lake and around the Onuma	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.014	-	-	-	N.D.	N.D.(2.3)	N.D.(2.3)	-
			Algae/plant	Zygnematophyceae	Zygnematales	Zygnemataceae	<i>Spirogyra sp.</i>	Spirogyra	-	0.31	-	-	-	8.20	0.70	7.5	-
			Algae/plant	Dicotyledoneae	Nymphaeales	Nymphaeaceae	<i>Nuphar japonicum</i>	Cow lily	-	0.33	-	-	-	0.87	N.D.(0.26)	0.87	-
			Algae/plant	Dicotyledoneae	Solanales	Menyanthaceae	<i>Nymphaoides peltata</i>	Fringed water-lily	-	0.35	-	-	-	0.29	N.D.(0.27)	0.29	-
			Arthropoda	Insecta	Odonata	Corduliidae	<i>Epophthalmia elegans</i>	Epophthalmia elegans	74	0.028	Larva(Dragonfly larva)	-	-	5.0	N.D.(1.7)	5.0	-
			Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	Anotogaster sieboldii				-	-	-	-	-	-
			Arthropoda	Insecta	Odonata	Libellulidae	<i>Deielia phaon</i>	Frosted skimmer				-	-	-	-	-	-
			Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Common prawn	795	0.49	Imago	-	-	9.56	0.96	8.6	0.66
			Mollusca	Gastropoda	Architaenioglossa	Viviparidae	<i>Cipangopaludina japonica</i>	Japanese mysterysnail	8	0.055	Imago	-	Molluscous part	7.9	N.D.(1.3)	7.9	-
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	11	1.3	Mature fish	Obscure digesta	Viscera removed	11.0	1.0	10	0.26
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	3	1.4	Mature fish	Obscure digesta	Viscera removed	25.1	2.1	23	0.39
			Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	1	1.2	Mature fish	Obscure digesta	Viscera removed	18.7	1.7	17	0.39
			Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	1	0.12	Mature fish	Japanese smelt	Viscera removed	16.5	1.5	15	-
			Vertebrata	Amphibia	Anura	Glandirana	<i>Glandirana rugosa</i>	Wrinkled Frog	8	0.071	Imago	-	-	N.D.	N.D.(1.3)	N.D.(1.1)	-
			Vertebrata	Amphibia	Caudata	Salamandridae	<i>Cynops pyrrhogaster</i>	Cynops pyrrhogaster	22	0.11	Imago	-	-	1.3	N.D.(0.71)	1.3	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)					
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
Off the mouth of the Abukuma River	Surrounding water area off the mouth of the Abukuma River	2018/6/4	Sea area in front of the Abukuma River Estuary	Arthropoda	Malacostraca	Decapoda	Portunidae	<i>Portunus trituberculatus</i>	Japanese blue crab	3	0.57	Imago	-	-	N.D.	N.D.(0.30)	N.D.(0.31)	-	
				Mollusca	Cephalopoda	Octopoda	Octopodidae	<i>Octopus vulgaris</i>	Common octopus	1	0.27	Imago	-	-	N.D.	N.D.(0.26)	N.D.(0.27)	-	
				Vertebrata	Osteichthyes	Lophiiformes	Lophiidae	<i>Lophius budegassa</i>	Monkfish	2	1.3	Immature fish	Fish	Viscera removed	N.D.	N.D.(0.30)	N.D.(0.35)	-	
				Vertebrata	Osteichthyes	Scorpaeniformes	Hexagrammidae	<i>Hexagrammos otakii</i>	Fat greenling	3	0.36	Mature fish	Shrimp	Viscera removed	0.39	N.D.(0.37)	0.39	-	
				Vertebrata	Osteichthyes	Scorpaeniformes	Triglidae	<i>Chelidonichthys spinosus</i>	Gurnard	8	2.3	Mature fish	Crab	Viscera removed	N.D.	N.D.(0.25)	N.D.(0.29)	N.D.(0.016)	
				Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Kareius bicoloratus</i>	Stone flounder	1	0.66	Immature fish	Obscure digesta	Viscera removed	N.D.	N.D.(0.38)	N.D.(0.33)	-	
				Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Pleuronectes herzensteini</i>	Yellow striped flounder	2	0.63	Mature fish	Obscure digesta	Viscera removed	0.40	N.D.(0.37)	0.40	-	
				Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Eopsetta grigorjewi</i>	Shotched halibut	6	0.63	Immature fish	Obscure digesta	Viscera removed	N.D.	N.D.(0.35)	N.D.(0.28)	-	
				Vertebrata	Osteichthyes	Perciformes	Polyprionidae	<i>Stereolepis doederleini</i>	Striped jewfish	3	0.45	Immature fish	Fish	Viscera removed	N.D.	N.D.(0.39)	N.D.(0.39)	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Acanthogobius flavimanus</i>	Yellowfin Goby	6	0.074	Immature fish,Mature fish	-	-	1.7	N.D.(0.75)	1.7	-	
Off Soma City	Matsukawaura Lagoon	2018/6/4		Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.012	-	-	-	22.9	2.9	20	-	
				Algae/plant	Monocotyledoneae	Najadales	Zosteraceae	<i>Zostera marina</i>	Eel grass	-	0.23	-	-	-	N.D.	N.D.(0.33)	N.D.(0.30)	-	
		2018/6/2		Algae/plant	Chlorophyceae	Ulvales	Ulvaceae	<i>Ulva pertusa</i>	Ulva pertusa	-	0.29	-	-	-	3.0	N.D.(0.26)	3.0	-	
				Annelida	Polychaeta	Eunicida	Eunicidae	<i>Polychaeta</i>	Polychaeta	281	0.076	Imago	-	-	5.91	0.41	5.5	-	
				Annelida	Polychaeta	Phyllodocida	Nereididae	<i>Hediste sp.</i>	Hediste										
				Arthropoda	Malacostraca	Decapoda	Alpheidae	<i>Alpheidae</i>	Alpheidae	27	0.042	Juvenile,Imago	-	-	1.6	N.D.(1.0)	1.6	-	
				Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon sp.</i>	Palaemon	61	0.033	Juvenile,Imago	-	-	2.1	N.D.(1.5)	2.1	-	
				Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Hemigrapsus sp.</i>	Hemigrapsus	134	0.39	Juvenile,Imago	-	-	3.94	0.64	3.3	-	
				Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	1	0.12	Imago	-	-	4.1	N.D.(0.66)	4.1	-	
		2018/6/4		Mollusca	Bivalvia	Ostreoida	Ostreidae	<i>Crassostrea gigas</i>	Oyster	6	0.15	Imago	-	-	Molluscous part	0.99	N.D.(0.43)	0.99	-
				Mollusca	Bivalvia	Venerida	Veneridae	<i>Ruditapes philippinarum</i>	Japanese littleneck	30	0.16	Imago	-	-	Molluscous part	1.2	N.D.(0.40)	1.2	-
		2018/6/3		Vertebrata	Osteichthyes	Scorpaeniformes	Hexagrammidae	<i>Hexagrammos otakii</i>	Fat greenling	3	0.087	Immature fish	Ragworm	Viscera removed	N.D.	N.D.(0.92)	N.D.(0.80)	-	
				Vertebrata	Osteichthyes	Scorpaeniformes	Hexagrammidae	<i>Hexagrammos agrammus</i>	Hexagrammos agrammus	3	0.052	Immature fish	Gammarus,Skeleton shrimp	Viscera removed	N.D.	N.D.(0.96)	N.D.(0.88)	-	
				Vertebrata	Osteichthyes	Scorpaeniformes	Scorpaenidae	<i>Sebastes cheni</i>	Rockfish	5	0.14	Immature fish	Mysidae,Fish	Viscera removed	0.69	N.D.(0.67)	0.69	-	
		2018/6/2		Vertebrata	Osteichthyes	Perciformes	Pholididae	<i>Pholis crassispina</i>	Pholis crassispina	30	0.057	Immature fish,Mature fish	-	-	N.D.	N.D.(1.1)	N.D.(0.95)	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Tridentiger trigonocephalus</i>	Chameleon goby						2.7	N.D.(0.63)	2.7	-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Acentrogobius virgatus</i>	Acentrogobius virgatus	56	0.10	Immature fish	-	-	2.7			-	
				Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Favonigobius gymnauchen</i>	Favonigobius gymnauchen										
		2018/6/3		Vertebrata	Osteichthyes	Perciformes	Sebastidae	<i>Sebastes schlegelii</i>	Black rockfish	2	0.086	Immature fish	Crab	Viscera removed	0.92	N.D.(0.88)	0.92	-	
		2018/6/2		Vertebrata	Osteichthyes	Tetraodontiformes	Tetraodontidae	<i>Takifugu niphobles</i>	Takifugu niphobles	1	0.093	Mature fish	Crab	Viscera removed	0.97	N.D.(0.84)	0.97	-	

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae, etc.) that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.

Location	Sampling point	Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
											Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
Off Iwaki City	M-1 M-2 M-3	2018/6/5	Offshore of Hisanohama	Mollusca	Cephalopoda	Decapodiformes	<i>Loiginiidae</i>	<i>Loioius japonica</i>	Japanese squid	7	0.18	Imago	-	-	N.D.	N.D.(0.43)	N.D.(0.42)	-
				Echinodermata	Astroidea	Forcipulatida	<i>Asteriidae</i>	<i>Asterias amurensis</i>	Northern Pacific seastar	3	0.75	Imago	-	-	N.D.	N.D.(0.28)	N.D.(0.29)	-
				Vertebrata	Osteichthyes	Lophiiformes	<i>Lophiidae</i>	<i>Lophiomus setigerus</i>	Monkfish	4	3.6	Immature fish	Searobin	Viscera removed	N.D.	N.D.(0.30)	N.D.(0.38)	-
				Vertebrata	Osteichthyes	Scorpaeniformes	<i>Triglidae</i>	<i>Lepidotrigla microptera</i>	Searobin	13	2.3	Immature fish,Mature fish	Shrimp	Viscera removed	0.62	N.D.(0.32)	0.62	-
				Vertebrata	Osteichthyes	Pleuronectiformes	<i>Pleuronectidae</i>	<i>Kareius bicoloratus</i>	Stone flounder	4	3.6	Immature fish,Mature fish	Crab	Viscera removed	0.34	N.D.(0.30)	0.34	-
				Vertebrata	Osteichthyes	Pleuronectiformes	<i>Pleuronectidae</i>	<i>Pleuronectes herzensteini</i>	Yellow striped flounder	7	2.2	Immature fish,Mature fish	Ragworm	Viscera removed	1.8	N.D.(0.43)	1.8	-
				Vertebrata	Osteichthyes	Pleuronectiformes	<i>Pleuronectidae</i>	<i>Pleuronectes yokohamae</i>	Marbled sole	4	2.0	Mature fish	Ragworm	Viscera removed	1.7	N.D.(0.36)	1.7	-
				Vertebrata	Osteichthyes	Pleuronectiformes	<i>Pleuronectidae</i>	<i>Eopsetta grigorjewi</i>	Shotched halibut	2	0.61	Immature fish	Bivalvia	Viscera removed	1.2	N.D.(0.47)	1.2	-
				Vertebrata	Osteichthyes	Pleuronectiformes	<i>Paralichthyidae</i>	<i>Paralichthys olivaceus</i>	Bastard halibut	2	4.3	Mature fish	Fish	Viscera removed	0.77	N.D.(0.34)	0.77	0.014
				Vertebrata	Osteichthyes	Perciformes	<i>Sparidae</i>	<i>Evynnis japonica</i>	Crimson sea-bream	3	0.39	Immature fish	Mysid shrimp	Viscera removed	N.D.	N.D.(0.33)	N.D.(0.33)	-
				Vertebrata	Osteichthyes	Tetraodontiformes	<i>Tetraodontidae</i>	<i>Takifugu snyderi</i>	Obscure Puffer	5	0.97	Immature fish	Bivalvia,Ragworm,Crab	Viscera removed	0.63	N.D.(0.35)	0.63	-
				Vertebrata	Osteichthyes	Zeiformes	<i>Zeidae</i>	<i>Zenopsis nebulosa</i>	Dory	12	0.39	Immature fish	Empty stomach	Viscera removed	0.41	N.D.(0.33)	0.41	-
				Vertebrata	Osteichthyes	Zeiformes	<i>Zeidae</i>	<i>Zeus faber</i>	John Dory	2	1.6	Mature fish	Fish	Viscera removed	0.47	N.D.(0.36)	0.47	-
				Vertebrata	Chondrichthyes	Rajiformes	<i>Rajidae</i>	<i>Okamejei kenojei</i>	Common Skete	3	2.5	Immature fish	Shrimp,Crab	Viscera removed	2.03	0.33	1.7	-
	M-4	Hisanohama Coastal areas	2018/6/5	Algae/plant	Phaeophyceae	Laminariales	<i>Laminariaceae</i>	<i>Eisenia bicyclis</i>	<i>Eisenia bicyclis</i>	-	0.28	-	-	-	0.34	N.D.(0.35)	0.34	-
				Mollusca	Gastropoda	Archaeogastropoda	<i>Haliotidae</i>	<i>Haliothis sp.</i>	Abalone	3	0.27	Imago	-	Molluscous part	N.D.	N.D.(0.33)	N.D.(0.29)	-
				Echinodermata	Echinoidea	Echinida	<i>Strongylocentrotidae</i>	<i>Strongylocentrotus nudus</i>	Northern sea urchin	8	0.82	Imago	-	-	0.54	N.D.(0.31)	0.54	-
				Vertebrata	Osteichthyes	Scorpaeniformes	<i>Hexagrammidae</i>	<i>Hexagrammos agrammus</i>	Hexagrammos agrammus	1	0.10	Immature fish	Bivalvia	Viscera removed	1.1	N.D.(0.76)	1.1	-
				Vertebrata	Osteichthyes	Scorpaeniformes	<i>Scorpaenidae</i>	<i>Sebastes cheni</i>	Rockfish	13	0.34	Immature fish	Amphipod	Viscera removed	0.94	N.D.(0.32)	0.94	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.