

Results of Radioactive Material Monitoring of Aquatic Organisms (Location G in Lake Hayama)

<Location G in Lake Hayama: Samples collected>

Items Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
G-1	○	○	○	○	○	○
G-4	○	○	○	-	○	-

<Location G in Lake Hayama: Site measurement item>

Items Locations	Latitude and longitude of the location		Survey date and time			Water temperature (degrees C)	Sediment			Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)		Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
G-1(Surface layer)	37.7321°	140.8127°	2018/12/3	11:23	11:50	11.8	9.3	Sand sediment	7.5Y 2/2	Plant pieces	3.8
						11.5					>3.8
G-4	37.7382°	140.8035°	2018/12/4	15:00	13:10	9.5	9.3	Sand gravel	7.5Y 5/3	Plant pieces	0.2
G-4	37.7382°	140.8035°	2018/12/4	15:00	7.3	0.6	2.4	11.0	7.7	0.05	1.3
G-4	37.7382°	140.8035°	2018/12/4	15:00	7.3	0.6	2.4	11.0	7.7	0.05	<1
G-4	37.7382°	140.8035°	2018/12/4	15:00	7.3	0.6	2.4	11.0	7.7	0.05	1.3

<Location G in Lake Hayama: General survey items/Analysis of radioactive materials Water>

Items Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity (%)	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
G-1(Surface layer)	37.7321°	140.8127°	2018/12/3	11:23	7.1	<0.5	3.3	9.6	6.7	0.04	1.8	1	1.0	0.0050	0.047	-
					7.1	<0.5	3.3	10.2	6.9	0.04	1.7	1	1.2	N.D.(0.0017)	0.023	0.0011
G-4	37.7382°	140.8035°	2018/12/4	15:00	7.3	0.6	2.4	11.0	7.7	0.05	1.3	<1	0.3	N.D.(0.0014)	0.0065	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location G in Lake Hayama: General survey items/Analysis of radioactive materials Sediment>

Items Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution						Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)		
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
G-1	37.7321°	140.8127°	2018/12/3	11:50	7.2	274	43.3	8.8	30.1	2.613	12.9	6.9	21.7	24.5	19.6	14.4	0.18	19	200	2200	2.9
G-4	37.7382°	140.8035°		13:10	7.5	298	20.0	1.7	2.3	2.695	31.2	44.7	22.1	1.0	1.0	1.4	19	25	280	-	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location G in Lake Hayama: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)		
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137			
G-1	In the lake	37.7321°	140.8127°	2018/12/3		Algae/plant	-	-	-	Plankton (Planktonic algae)	-	0.015	-	-	-	7.2	N.D.(1.7)	7.2	-		
G-2		37.7267°	140.8223°			Algae/plant	Monocotyledoneae	Alismatales	Hydrocharitaceae	<i>Elodea nuttallii</i>	Western Waterweed	-	0.20	-	-	-	59.5	4.5	55	-	
G-3	Inflowing rivers	37.7302°	140.8307°	2018/12/3		Algae/plant	-	-	-	Riverbed Deposits (Include algae)	-	0.0080	-	-	-	317	27	290	-		
G-4		37.7382°	140.8035°			Anthropoda	Insecta	Ephemeroptera	<i>Ephemeridae</i>	<i>Ephemerida japonica</i>	Ephemerida japonica	548	0.032	Larva	-	-	104.5	8.5	96	-	
						Anthropoda	Insecta	Ephemeroptera	<i>Ephemeridae</i>	<i>Ephemerida strigata</i>	Mont mayfly										
						Anthropoda	Insecta	Plecoptera	Perlidae	<i>Kamimura uenoii</i>	Kamimura uenoii Kohno										
						Anthropoda	Insecta	Plecoptera	Perlidae	<i>Kamimura tibialis</i>	Kamimura tibialis										
						Anthropoda	Insecta	Plecoptera	Perlidae	<i>Paragnetina suzukii</i>	Paragnetina suzukii Okamoto	276	0.023	Larva	-	-	8.1	N.D.(7.9)	8.1	-	
						Anthropoda	Insecta	Trichoptera	Stenophryctidae	<i>Stenophrycte marmorata</i>	Stenophrycte marmorata	142	0.021	Larva	-	-	47	N.D.(9.4)	47	-	
						Anthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	Protohermes grandis		11	0.0075	Larva	-	-	16	N.D.(15)	16	-
						Anthropoda	Insecta	Megaloptera	Corydalidae	<i>Parachauliodes japonicus</i>	Parachauliodes japonicus										
						Course Particulate Organic Matter	-	-	-	Bottom fallen leaves	-	0.20	-	-	-	30.3	2.3	28	-		

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowel) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

*6: River bottom materials (in algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.