

OResults of Radioactive Material Monitoring of Aquatic Organisms (Location E along the Niida River)

<Location E along the Niida River: Samples collected>

Items	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
E-2 a	○	○	○	○	○	○

<Location E along the Niida River: Site measurement item>

Items	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Transparency (cm)
E-2 a	37.6640°	140.9447°	2019/8/28	12:00	12:20	21.3	21.6	Silt	5Y3/1	Plant pieces	0.40	>50

<Location E along the Niida River: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
	Latitude	Longitude	Date	Time (water)		(mg/L)	(mg/L)	(mg/L)	(mS/m)		(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)
E-2 a	37.6640°	140.9447°	2019/8/28	12:00	7.0	0.6	3.7	8.9	8.7	0.05	1.6	4	3.5	0.0020	0.039	0.0017

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location E along the Niida River: General survey items/Analysis of radioactive materials Sediment>

Items	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution							Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)	
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)				Maximum grain diameter (mm)
Locations																					
E-2 a	37.6640°	140.9447°	2019/8/28	12:20	7.1	63	64.3	13.3	47.1	2.505	1.3	3.4	4.1	17.5	41.0	32.7	0.018	4.8	480	6400	1.3

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location E along the Niida River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
E-2 b	The main stream of the Niida River	37.6635°	140.9452°	2019/8/29	Algae/plant	-	-	-	-	Riverbed Deposits (Include algae)	-	0.012	-	-	-	45.8	3.8	42	-
					Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata	185	0.024	Larva	-	-	151	11	140	-
					Arthropoda	Insecta	Odonata	Corduliidae	<i>Macromia amphigena amphigena</i>	<u>Macromia amphigena</u>	30	0.013	Larva(Dragonfly larva)	-	-	20	N.D.(3.0)	20	-
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Meligomphus viridicostus</i>	Meligomphus viridicostus									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Sieboldius albardae</i>	Sieboldius albardae									
					Arthropoda	Insecta	Odonata	Gomphidae	<i>Davidius sp.</i>	Davidius									
					Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	Protohermes grandis	12	0.010	Larva	-	-	17	N.D.(3.1)	17	-
					Arthropoda	Malacostraca	Decapoda	Palaemonidae	<i>Palaemon paucidens</i>	Common prawn	26	0.036	Imago	-	-	29.5	1.5	28	-
					Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	13	0.11	Juvenile	-	-	43.8	2.8	41	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	60	0.79	Immature fish	-	-	32.2	2.2	30	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Opsariichthys platypus</i>	Pale chub	50	0.28	Immature fish	-	-	20.3	1.3	19	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Pseudogobio esocinus esocinus</i>	Pseudogobio esocinus esocinus	1	0.023	Mature fish	-	-	18	N.D.(3.2)	18	-
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	10	0.24	Immature fish,Mature fish	-	-	94.3	6.3	88	-
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	21	0.073	Mature fish	-	-	39.3	2.3	37	-
					Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius nagoyae									
					Vertebrata	Amphibia	Anura	Ranidae	<i>Rana japonica</i>	Japanese brown frog	4	0.012	Imago	-	-	12	N.D.(3.0)	12	-
					Vertebrata	Amphibia	Anura	Hylidae	<i>Hyla japonica</i>	<u>Japanese tree flog</u>									
					Vertebrata	Amphibia	Anura	Pelophylax	<i>Pelophylax porosus porosus</i>	Tokyo daruma pond frog									
					Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.24	-	-	-	106.6	6.6	100	-

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.