

## Results of Radioactive Material Monitoring of Aquatic Organisms (Location D along the Mano River)

<Location D along the Mano River: Samples collected>

Items	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
D-4 a	○	○	○	○	○	○

<Location D along the Mano River: Site measurement item>

Items	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other		
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Transparency (cm)	
D-4 a	37.7308°	140.9081°	2019/11/5	13:05	13:20	14.9	14.9	Sand	2.5Y 4/3	None	0.78	19	

<Location D along the Mano River: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity (mg/L)	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
D-4 a	37.7308°	140.9081°	2019/11/5	13:05	7.4	<0.5	3.3	10.9	7.9	0.04	1.6	13	26.5	0.011	0.14	0.0013

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location D along the Mano River: General survey items/Analysis of radioactive materials Sediment>

Items	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>NHE</sub> (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution						Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)	
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (Less than 0.005mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)		
D-4 a	37.7308°	140.9081°	2019/11/5	13:20	7.2	530	16.4	1.7	2.2	2.725	48.8	24.7	17.7	7.4	1.4	2.0	9.5	12	160	0.94

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location D along the Mano River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)	
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
D-3	The main stream of the Mano River	37.7051°	140.9623°	2019/11/3	Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus reinii</i>	Sculpin	1	0.030	Immature fish	-	-	9.6	N.D.(4.9)	9.6	-	
				2019/11/5	Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Plecoglossus altivelis altivelis</i>	Sweetfish	109	2.7	Immature fish,Mature fish	-	-	24.8	1.8	23	0.18	
D-4 b	The main stream of the Mano River	37.7312°	140.9096°	2019/11/2		Algae/plant	Monocotyledoneae	Najadales	Potamogetonaceae	<i>Potamogeton berchtoldii</i>	Small pondweed	-	0.11	-	-	-	16.2	1.2	15	-
						Arthropoda	Insecta	Megaloptera	Corydalidae	<i>Protohermes grandis</i>	Protohermes grandis	29	0.013	Larva	-	-	13	N.D.(6.9)	13	-
						Arthropoda	Malacostraca	Decapoda	Cambaridae	<i>Procambarus clarkii</i>	Red swamp crawfish	4	0.11	Imago	-	-	43.5	2.5	41	-
						Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus keta</i>	Salmon	1	2.0	Mature fish	-	-	N.D.	N.D.(0.82)	N.D.(0.90)	N.D.(0.018)
						Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius fluviatilis</i>	Rhinogobius fluviatilis	13	0.036	Mature fish	-	-	23	N.D.(2.5)	23	-
						Vertebrata	Osteichthyes	Perciformes	Gobiidae	<i>Rhinogobius nagoyae</i>	Rhinogobius nagoyae			-	-	6.1	N.D.(3.9)	6.1	-	
						Vertebrata	Amphibia	Anura	Ranidae	<i>Rana japonica</i>	Japanese brown frog	3	0.036	Imago	-	-	25.0	2.0	23	-
						Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.23	-	-	-	Viscera removed	394	24	370
D-5	The main stream of the Mano River	37.7214°	140.8889°	2019/11/3	Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	2	0.12	Immature fish,Mature fish	Empty stomach	Viscera removed	394	24	370	-	

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.