

**Results of Radioactive Material Monitoring of Aquatic Organisms (Location F along the Ota River)**

<Location F along the Ota River: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
F-1	○	○	○	○	○	○

<Location F along the Ota River: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Transparency (cm)
F-1	37.5975°	140.9252°	2019/11/5	08:58	09:18	13.5	13.1	Sand	2.5Y 4/3	None	0.55	18

<Location F along the Ota River: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
F-1	37.5975°	140.9252°	2019/11/5	08:58	7.5	<0.5	3.7	11.6	3.9	0.03	1.3	17	24.0	0.034	0.48	0.0041

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location F along the Ota River: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>NHE</sub> (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
F-1	37.5975°	140.9252°	2019/11/5	09:18	6.9	463	21.2	1.4	3.4	2.670	7.8	22.7	43.5	19.6	2.8	3.6	0.48	9.5	15	230	0.71

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location F along the Ota River: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137	
F-1	The main stream of the Ota River	37.5975°	140.9252°	2019/11/3	Arthropoda	Malacostraca	Decapoda	Varunidae	<i>Eriocheir japonica</i>	Japanese mitten crab	2	0.0079	Juvenile	-	-	385	25	360	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	4	0.074	Immature fish	-	-	392	22	370	-
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou</i>	Yamame trout	1	0.020	Immature fish	-	-	423	33	390	-
					Vertebrata	Amphibia	Anura	Ranidae	<i>Rana japonica</i>	Japanese brown frog	2	0.034	Imago	-	-	30.0	2.0	28	-
					Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.23	-	-	-	169.1	9.1	160	-
F-3	The main stream of the Ota River	37.6045°	140.9636°	2019/11/3	Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	1	0.066	Mature fish	Obscure digesta	Viscera removed	163	13	150	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp	1	0.066	Immature fish	Obscure digesta	Viscera removed	148.6	8.6	140	-
F-5	The main stream of the Ota River	37.6022°	140.9868°	2019/11/3	Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus keta</i>	Salmon	1	3.2	Mature fish	Empty stomach	Viscera removed	N.D.	N.D.(0.33)	N.D.(0.28)	0.045

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.