

## ○Results of Radioactive Material Monitoring of Aquatic Organisms (Location H in Lake Akimoto)

<Location H in Lake Akimoto: Samples collected>

Items	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
H-1	○	○	○	○	○	○
H-2	○	○	○	-	○	-

<Location H in Lake Akimoto: Site measurement item>

Items	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)		Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)
H-1(Surface layer)	37.6575°	140.1264°	2020/7/3	08:42	08:55	20.0	10.6	Ooze	7.5Y 3/2	Plant pieces	10.8	3.5
H-1(Bottom layer)						10.2						
H-2(Surface layer)	37.6616°	140.1226°		09:10	09:22	19.4	13.2	Ooze	7.5Y 5/3	Plant pieces	6.0	2.5
H-2(Bottom layer)						13.6						

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Water>

Items	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
H-1(Surface layer)	37.6575°	140.1264°	2020/7/3	08:42	7.3	0.5	3.5	8.5	4.6	0.05	2.0	1	1.3	N.D.(0.0015)	0.0069	-
H-1(Bottom layer)					6.9	<0.5	2.9	8.4	4.8	0.05	1.8	3	2.1	N.D.(0.0015)	0.0043	0.00093
H-2(Surface layer)	37.6616°	140.1226°		09:10	7.1	0.7	3.4	8.5	5.0	0.03	1.4	2	2.1	N.D.(0.0015)	0.0069	-
H-2(Bottom layer)					6.5	0.7	3.2	1.6	5.7	0.03	1.2	12	10.4	0.0022	0.059	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Sediment>

Items	Latitude and longitude of the location		Survey date and time		pH	Redox potential E <sub>NHE</sub> (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm <sup>3</sup> )	Grain size distribution						Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)		
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
H-1	37.6575°	140.1264°	2020/7/3	08:55	7.0	78	67.5	10.7	35.0	2.543	0.0	0.1	0.3	0.1	53.9	45.6	0.0057	2.0	150	2600	1.4
H-2	37.6616°	140.1226°		09:22	6.9	50	75.1	14.2	46.4	2.443	0.0	0.1	0.2	0.3	18.8	80.6	-	2.0	120	2200	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

## &lt;Location H in Lake Akimoto: Analysis items Aquatic organisms&gt;

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)				
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
H-1 H-2 H-3	In the lake	37.6575° 37.6616° 37.6653°	140.1264° 140.1226° 140.1329°	2020/7/3	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.014	-	-	-	2.2	N.D.(2.0)	2.2	-	
				2020/6/17	Arthropoda	Malacostraca	Decapoda	Astacidae	<i>Pacifastacus leniusculus trowbridgii</i>	Signal crayfish	18	0.42	Imago	-	-	-	20.0	1.0	19	5.9
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	17	3.5	Mature fish	-	-	-	32.5	1.5	31	0.60
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsfordii	5	2.7	Mature fish	Obscure digesta	Viscera removed	33.4	1.4	32	1.2	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Cyprinus carpio</i>	Common carp	1	4.1	Mature fish	Obscure digesta	Viscera removed	11.74	0.74	11	0.55	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	3	1.2	Immature fish,Mature fish	Obscure digesta	Viscera removed	32.0	2.0	30	0.99	
				2020/7/3	Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenoides</i>	Char	3	0.43	Mature fish	Japanese smelt,Plant piece,Diptera(imago)	Viscera removed	18.96	0.96	18	-	
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou masou</i>	Seema	7	0.80	Immature fish	Fish,Diptera(imago),Terrestrial insect,Ephemera,Plant piece	Viscera removed	13.69	0.69	13	-	
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	7	2.1	Immature fish,Mature fish	Plant piece,Common prawn,Ephemeroptera(imago)	Viscera removed	45.3	2.3	43	1.0	
				2020/7/3	Algae/plant	Monocotyledoneae	Alismatales	Hydrocharitaceae	<i>Elodea nuttallii</i>	Western waterweed	-	0.28	-	-	-	6.24	0.44	5.8	-	
H-3	Inflowing rivers	37.6653°	140.1329°	2020/7/3	Arthropoda	Insecta	Ephemeroptera	Ameletidae	<i>Ameletus sp.</i>	Ameletus	220	0.0096	Larva	-	-	N.D.	N.D.(3.8)	N.D.(3.2)	-	
					Arthropoda	Insecta	Ephemeroptera	Heptageniidae	<i>Epeorus latifolium</i>	<i>Epeorus latifolium</i>				-	-				-	
					Arthropoda	Insecta	Ephemeroptera	Ephemerellidae	<i>Drunella trispina</i>	Drunella trispina				-	-				-	
					Arthropoda	Insecta	Plecoptera	Perlidae	<i>Acronerura sp.</i>	Acronerura	102	0.025	Larva	-	-	N.D.	N.D.(1.9)	N.D.(1.9)	-	
					Arthropoda	Insecta	Plecoptera	Perlidae	<i>Kamimuria quadrata</i>	Kamimuria quadrata				-	-				-	
				2020/7/1	Arthropoda	Insecta	Plecoptera	Perlidae	<i>Calineuria sp.</i>	Calineuria	12	0.0043	Larva	-	-	N.D.	N.D.(7.7)	N.D.(6.9)	-	
					Arthropoda	Insecta	Trichoptera	Stenopsychidae	<i>Stenopsyche marmorata</i>	Stenopsyche marmorata				-	-				-	
					Arthropoda	Insecta	Odonata	Corduliidae	<i>Somatochlora uchidai</i>	Somatochlora uchidai				-	-				-	
					Arthropoda	Insecta	Odonata	Cordulegastridae	<i>Anotogaster sieboldii</i>	Anotogaster sieboldii				-	-	4.9	N.D.(4.8)	4.9	-	
					Arthropoda	Insecta	Odonata	Libellulidae	<i>Orthetrum albistylum speciosum</i>	Common skimmer				-	-				-	
				2020/7/3	Arthropoda	Insecta	Odonata	Aeshnidae	<i>Aeshna juncea</i>	Common hawk	27	0.010	Juvenile,Imago	-	-	5.4	N.D.(4.5)	5.4	-	
					Vertebrata	Osteichthyes	Scorpaeniformes	Cottidae	<i>Cottus pollux</i>	Japanese fluvial sculpin				-	-		8.1	N.D.(0.39)	8.1	-
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenoides</i>	Char				-	-				-	
				2020/7/1	Vertebrata	Amphibia	Anura	-	-	Frog	240	0.13	Larva(Tadpole)	-	-	137.3	7.3	130	-	
				2020/7/3	Coarse Particulate Organic Matter	-	-	-	-	Bottom fallen leaves	-	0.22	-	-	-	8.0	N.D.(0.36)	8.0	-	

\*1: Organisms were collected in or around the targeted water areas.

\*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

\*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

\*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

\*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40μm-mesh).

\*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

\*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

\*8: Activity concentrations include counting errors, but the details are omitted here.