

OResults of Radioactive Material Monitoring of Aquatic Organisms (Location M off Iwaki City)

<Location M off Iwaki City: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
M-2	○	○	○	○	○	○

<Location M off Iwaki City: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
M-2(Surface layer)	37.1996°	141.0853°	2020/7/14	08:36	08:50	19.6	16.2	Sand	5Y3/1	Shell fragments	43.5	5.2
M-2(Bottom layer)				08:10		13.9						

<Location M off Iwaki City: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD	COD	DO	Electric conductivity	Salinity	TOC	SS	Turbidity	Cs-134	Cs-137	Sr-90
	Latitude	Longitude	Date	Time (water)	(mg/L)	(mg/L)	(mg/L)	(mS/m)	(mg/L)	(mg/L)	(FNU)	(Bq/L)	(Bq/L)	(Bq/L)		
M-2(Surface layer)	37.1996°	141.0853°	2020/7/14	08:36	8.1	0.7	2.6	8.0	4780	31.42	1.1	3	1.5	N.D.(0.0015)	0.0066	-
M-2(Bottom layer)				08:10	8.0	<0.5	2.3	8.5	5090	33.70	1.0	2	0.8	N.D.(0.0014)	0.0039	0.00077

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location M off Iwaki City: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHLE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution						Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)		
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)				Median grain diameter (mm)	Maximum grain diameter (mm)
											0.0	0.9	3.2	93.0	2.9					0.17	2.0
M-2	37.1996°	141.0853°	2020/7/14	08:50	7.7	171	22.3	2.0	3.6	2.763	0.0	0.9	3.2	93.0	2.9		0.17	2.0	1.5	32	N.D.(0.16)

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location M off Iwaki City: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)	
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
M-1 M-2 M-3	Offshore of Hisanohama	37.1736° 37.1996° 37.2324°	141.0788° 141.0853° 141.0935°	2020/7/14	Echinodermata	Asteroidea	Forcipulatida	Asteriidae	<i>Asterias amurensis</i>	Northern pacific seastar	2	0.98	Imago	-	-	-	N.D.	N.D.(0.42)	N.D.(0.38)	-
					Vertebrata	Osteichthyes	Scorpaeniformes	Triglidae	<i>Lepidotrigla microptera</i>	Searobin	6	1.1	Immature fish,Mature fish	Amphipod,Shrimp	Viscera removed	0.61	N.D.(0.32)	0.61	-	
					Vertebrata	Osteichthyes	Scorpaeniformes	Triglidae	<i>Chelidonichthys spinosus</i>	Gurnard	1	0.19	Mature fish	Amphipod,Fish	Viscera removed	0.64	N.D.(0.88)	0.64	-	
					Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Pleuronichthys japonicus</i>	Finespotted flounder	9	1.1	Immature fish,Mature fish	Ragworm	Viscera removed	0.63	N.D.(0.45)	0.63	-	
					Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Pleuronectes herzensteini</i>	Yellow striped flounder	1	0.43	Mature fish	Shrimp	Viscera removed	0.86	N.D.(0.35)	0.86	-	
					Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Pleuronectes yokohamae</i>	Marbled sole	1	0.51	Mature fish	Empty stomach	Viscera removed	0.87	N.D.(0.41)	0.87	-	
					Vertebrata	Osteichthyes	Pleuronectiformes	Pleuronectidae	<i>Eopsetta grigorjewi</i>	Shotted halibut	4	1.1	Immature fish	Shrimp,Crab	Viscera removed	1.3	N.D.(0.50)	1.3	N.D.(0.019)	
					Vertebrata	Osteichthyes	Pleuronectiformes	Paralichthyidae	<i>Paralichthys olivaceus</i>	Bastard halibut	1	2.6	Mature fish	Fish	Viscera removed	1.1	N.D.(0.32)	1.1	-	
					Vertebrata	Osteichthyes	Perciformes	Sparidae	<i>Pagrus major</i>	Red seabream	1	1.2	Mature fish	Shellfish	Viscera removed	0.58	N.D.(0.57)	0.58	N.D.(0.020)	
					Vertebrata	Osteichthyes	Tetraodontiformes	Tetraodontidae	<i>Takifugu snyderi</i>	Obscure puffer	2	0.71	Immature fish	Shellfish,Sea urchin	Viscera removed	0.72	N.D.(0.42)	0.72	-	
					Vertebrata	Osteichthyes	Zeiformes	Zeidae	<i>Zeus faber</i>	John dory	3	1.1	Immature fish,Mature fish	Fish,Squid	Viscera removed	N.D.	N.D.(0.47)	N.D.(0.41)	N.D.(0.025)	
					Vertebrata	Chondrichthyes	Rajiformes	Rajidae	<i>Okamejei kenojei</i>	Common skate	2	1.8	Immature fish	Crab,Shrimp,Squid	Viscera removed	1.3	N.D.(0.31)	1.3	0.045	

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.