

Results of Radioactive Material Monitoring of Aquatic Organisms (Location H in Lake Akimoto)

<Location H in Lake Akimoto: Samples collected>

Locations	General items		Radioactive materials			
	Water	Sediment	Water (Cs)	Water (Sr)	Sediment (Cs)	Sediment (Sr)
H-1	○	○	○	○	○	○
H-2	○	○	○	-	○	-

<Location H in Lake Akimoto: Site measurement item>

Locations	Latitude and longitude of the location		Survey date and time			Water	Sediment				Other	
	Latitude	Longitude	Date	Time (water)	Time (sediment)	Water temperature (degrees C)	Sediment temperature (degrees C)	Property	Color	Contaminants	Water depth (m)	Secchi disk depth (m)
H-1(Surface layer)	37.6575°	140.1264°	2020/10/23	08:43	08:59	14.4	13.3	Ooze	7.5Y 4/2	Plant pieces	13.0	4.5
H-1(Bottom layer)						13.6						
H-2(Surface layer)	37.6616°	140.1226°		09:25	09:36	14.2	14.0	Ooze	7.5Y 4/1	Plant pieces	5.0	2.5
H-2(Bottom layer)												

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Water>

Locations	Latitude and longitude of the location		Survey date and time		pH	BOD (mg/L)	COD (mg/L)	DO (mg/L)	Electric conductivity (mS/m)	Salinity	TOC (mg/L)	SS (mg/L)	Turbidity (FNU)	Cs-134 (Bq/L)	Cs-137 (Bq/L)	Sr-90 (Bq/L)
	Latitude	Longitude	Date	Time (water)												
H-1(Surface layer)	37.6575°	140.1264°	2020/10/23	08:43	7.2	0.7	3.9	9.0	5.5	0.03	1.5	2	1.0	N.D.(0.0011)	0.0078	-
H-1(Bottom layer)					7.0	<0.5	3.3	8.2	5.4	0.03	1.4	3	2.0	N.D.(0.0011)	0.0052	0.00092
H-2(Surface layer)	37.6616°	140.1226°		09:25	7.0	0.8	3.7	9.2	5.9	0.03	1.9	3	2.4	N.D.(0.0011)	0.0087	-
H-2(Bottom layer)					7.0	<0.5	3.7	8.6	5.5	0.03	1.6	4	3.1	N.D.(0.0013)	0.018	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: General survey items/Analysis of radioactive materials Sediment>

Locations	Latitude and longitude of the location		Survey date and time		pH	Redox potential E _{NHLE} (mV)	Water content (%)	IL (%)	TOC (mg/g-dry)	Soil particle density (g/cm ³)	Grain size distribution								Cs-134 (Bq/kg-dry)	Cs-137 (Bq/kg-dry)	Sr-90 (Bq/kg-dry)
	Latitude	Longitude	Date	Time (sediment)							Gravel (2-75mm) (%)	Coarse sand (0.85-2mm) (%)	Medium sand (0.25-0.85mm) (%)	Fine sand (0.075-0.25mm) (%)	Silt (0.005-0.075mm) (%)	Clay (Less than 0.005mm) (%)	Median grain diameter (mm)	Maximum grain diameter (mm)			
H-1	37.6575°	140.1264°	2020/10/23	08:59	6.7	41	64.8	9.1	31.7	2.564	-	0.0	0.1	0.2	54.6	45.1	0.0062	2.0	30	700	1.2
H-2	37.6616°	140.1226°		09:36	6.7	10	74.0	12.0	43.5	2.486	0.0	0.1	0.1	0.5	37.4	61.9	0.0024	4.8	36	660	-

Note) N.D. means to be below the detection limit and figures in parentheses show the detection limit.

<Location H in Lake Akimoto: Analysis items Aquatic organisms>

Locations	Sampling point	Latitude and longitude of the location		Sampling date	Division	Class	Order	Family	Scientific name	English name	Population	Sample weight (kg-wet)	Note			Radioactive cesium (Bq/kg-wet)			Sr-90 (Bq/kg-wet)	
		Latitude	Longitude										Growth stage	Stomach contents	Measurement site	Total	Cs-134	Cs-137		
H-1 H-2 H-3	In the lake	37.6575° 37.6616° 37.6653°	140.1264° 140.1226° 140.1329°	2020/10/23	Algae/plant	-	-	-	-	Plankton (Planktonic algae)	-	0.019	-	-	-	N.D.	N.D.(2.2)	N.D.(2.1)	-	
				2020/10/30	Arthropoda	Malacostraca	Decapoda	Astacidae	<i>Pacifastacus leniusculus trowbridgii</i>	Signal crayfish	10	0.45	Imago	-	-	-	16	N.D.(1.7)	16	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Tribolodon hakonensis</i>	Japanese dace	10	1.6	Mature fish	Obscure digesta	Viscera removed	18.2	1.2	17	0.58	
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Carassius auratus</i>	Carassius auratus langsdorffii	Mature fish	5	2.1	Mature fish	Obscure digesta	Viscera removed	16	N.D.(1.2)	16	-
					Vertebrata	Osteichthyes	Cypriniformes	Cyprinidae	<i>Hemibarbus barbus</i>	Hemibarbus barbus	Mature fish	4	2.7	Mature fish	Obscure digesta	Viscera removed	64.8	2.8	62	-
					Vertebrata	Osteichthyes	Salmoniformes	Osmeridae	<i>Hypomesus nipponensis</i>	Japanese smelt	Immature fish, Mature fish	30	0.15	Immature fish, Mature fish	-	-	14	N.D.(0.90)	14	-
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Salvelinus leucomaenis</i>	Char	Mature fish	2	0.32	Mature fish	Empty stomach	Viscera removed	9.5	N.D.(1.1)	9.5	-
					Vertebrata	Osteichthyes	Salmoniformes	Salmonidae	<i>Oncorhynchus masou masou</i>	Seema	Immature fish	2	0.79	Immature fish	Terrestrial insect	Viscera removed	23.3	1.3	22	-
					Vertebrata	Osteichthyes	Perciformes	Centrarchidae	<i>Micropterus dolomieu</i>	Small mouth bass	Immature fish	8	1.7	Immature fish	Caterpillar	Viscera removed	35.6	1.6	34	-
					Vertebrata	Osteichthyes	Siluriformes	Siluridae	<i>Silurus asotus</i>	Amur catfish	Mature fish	2	1.1	Mature fish	Empty stomach	Viscera removed	39.9	1.9	38	0.74

*1: Organisms were collected in or around the targeted water areas.

*2: When multiple types of aquatic organisms were collected, a sample was prepared by mixing them.

*3: For a sample made of multiple types of aquatic organisms, the English name of the dominant one largest in number is underlined.

*4: Basically, measurement was conducted for all organism samples. Viscera (stomach and bowels) were removed for the measurement when possible so that undigested food and sediments, etc. in the digestive system would be excluded.

*5: Plankton (suspended algae) is the residue remaining after the filtration of lake water or seawater with a plankton net (40µm-mesh).

*6: River bottom materials (incl. algae) are algae, etc. that were scratched off stones with a brush, etc. and may include very fine particles such as inorganic silt and clay.

*7: N.D. means to be below the detection limit and figures in parentheses show the detection limit.

*8: Activity concentrations include counting errors, but the details are omitted here.