


何も手を打たなければ、海に流入するプラスチックの量は 20年後には10倍に増加する



Jamebeck et al. (2015), Science

マイクロプラスチックのリスクについて不確かなことあるが、国際的には**予防原則の立場**から、対策は進められています。

←何も手を打たなければ、
海洋プラスチック汚染は深刻化

←海洋のマイクロプラスチックは除去できない

国際的には予防原則的に動いている

2015年12月 アメリカで**マイクロビーズ**配合禁止の
連邦法成立

2014年8月：米カリフォルニア州で**レジ袋**禁止の法案成立

2014年11月：EUが加盟国へレジ袋削減案策定を義務づけ

2025年までにレジ袋の消費を
1人1年40枚まで削減がEUの目標

日本では年間300億枚以上のレジ袋が使われている。
1人あたりでは年間300枚

2016年： イギリスで**レジ袋**に課税

世界20カ国以上で
規制が行われている


国際的には予防原則的に動いている

2014年3月：米サンフランシスコ市で**ペットボトル**での飲料水の販売を禁止

2016年9月：フランスで「**プラスチック製使い捨て容器や食器を禁止**する法律」成立（2020年より）

プラスチックの原料の石油は数百万年以上前の生物遺骸が地殻中で変成したもの
 →石油由来のプラスチックは循環型ではない。

プラスチック削減は温暖化対策にもつながる



地球規模での炭素の循環 (単位: ギガトン; 矢印は年間移動量; 枠内は現存量)

国連環境計画の年鑑


UNEP YEAR BOOK

EMERGING ISSUES
IN OUR GLOBAL ENVIRONMENT

2011



United Nations Environment Programme



Plastic Debris in the Ocean

Every year large amounts of plastic debris enter the ocean, where it slowly fragments and accumulates in convergence zones. Scientists are concerned about the possible impacts of small plastic fragments—microplastics—in the environment. The role of plastics as a vector for transporting chemicals and species in the ocean is as yet poorly understood, but it is a potential threat to ecosystems and human health. Improved waste management is the key to preventing plastic and other types of litter from entering the ocean.

Box 3: Plastic pellets


Plastic resin pellets are small granules, generally in the shape of a cylinder or disc, with a diameter of a few millimetres. These particles are an industrial raw material that is remelted and moulded into final products. They enter the ocean as a result of spills or accidental releases. Like other plastic particles, they have been shown to accumulate PBTs. In the case of thin plastic films, for example those 50 micrometres or less, it may take only a few days for this process of accumulation or release to occur (Adams et al. 2007). In the case of pellets, equilibrium between the concentration of a given compound in a pellet and in the surrounding water or sediment may take many weeks or months. Older pellets consequently tend to have higher concentrations of contaminants and have been used to map the distribution of pollution in coastal waters around the world (Ogata et al. 2009, International Pellet Watch 2011) (Figure 5). Their consistent size makes them a useful monitoring tool.

Transport by plastic particles does not represent a significant additional flux of PBTs on a global scale compared with atmospheric or water transport (Zarfl and Matthies 2010). However, the concentration of contaminants by microplastic particles presents the possibility of increasing exposure to organisms through ingestion and entrance into the food chain—with the prospect of biomagnification in top-end predators in the food chain such as swordfish and seals. Ingestion of small particles by a wide variety of organisms has been well reported. However, the basic information needed on the biochemical and physiological response of organisms to ingested plastics contaminated with PBTs in order to quantify the scale of the problem is currently unavailable (Arthur et al. 2009, GESAMP 2010). It is conceivable that PBTs in plastic particles will be less bioavailable than those from the surrounding water or food sources (Gouin et al. 2011).



Collected from beaches around the world, plastic pellets like these have been found to accumulate persistent, bio-accumulating and toxic substances. The pellets are used in the manufacture of plastic products and have been introduced into the ocean through accidental releases. They may also be released as a result of poor handling or waste management. While there is evidence that quantities entering the marine environment have been reduced as a result of improved industrial practices, pellets already released will persist for many years. Credit: International Pellet Watch

Figure 5: Concentration of PCBs in beached plastic resin pellets, in nanograms per gram of pellet. Samples of polyethylene pellets have been collected at 56 beaches in 29 countries and analyzed for concentrations of organochlorine compounds. PCB concentrations were highest in pellets collected in the United States, Western Europe and Japan. They were lowest in those collected in tropical Asia and Africa. This spatial pattern reflects regional differences in the use of PCBs. Source: Ogata et al. (2009) with additional data provided by International Pellet Watch in 2010



国連でも議論され、海のプラスチック汚染が国際的に共通した懸念に



第17回「海洋及び海洋法に関する
国連総会非公式協議プロセス
(ICP)」

2016年6月13日～17日



ニューヨーク 国連本部

マイクロプラスチック国際条約についても議論が始まる



Marine Litter Advisory Group Meeting

9th - 11th May 2017, Windsor Golf Hotel & Country Club Nairobi-Kenya





THE
OCEAN
CONFERENCE
UNITED NATIONS, NEW YORK, 5-9 JUNE 2017



Side event

3 R as the Basis for Moving Towards **Zero Plastic Waste** in Coastal and Marine Environment


“Issue of microplastics in the coastal and marine environment and 3R solutions” Hideshige Takada (Japan)



持続可能な開発目標



14 LIFE BELOW WATER



14.1

By 2025, prevent and significantly reduce marine pollution of all kinds, in particular from land-based activities, including **marine debris** and nutrient pollution

14.1.1

Index of coastal eutrophication and floating **plastic debris** density



**United Nations Conference to Support the
Implementation of Sustainable Development Goal 14:
Conserve and sustainably use the oceans, seas and
marine resources for sustainable development**

New York, 5-9 June 2017

Item 11 of the provisional agenda*

Outcome of the Conference

Draft call for action

4. We are particularly alarmed by the adverse impacts of climate change on the ocean, including the rise in ocean temperatures, ocean and coastal acidification, deoxygenation, sea-level rise, the decrease in polar ice coverage, coastal erosion and extreme weather events. We acknowledge the need to address the adverse impacts that impair the crucial ability of the ocean to act as climate regulator, source of marine biodiversity and as key provider of food and nutrition, tourism and ecosystem services and as an engine for sustainable economic development and growth. We recognize, in this regard, the particular **importance of the Paris Agreement**, adopted under the United Nations Framework Convention on Climate Change.

(g) Accelerate actions to prevent and significantly reduce marine pollution of all kinds, particularly from land-based activities, including marine debris, plastics and microplastics, nutrient pollution, untreated wastewater, solid waste discharges, hazardous substances, pollution from ships and abandoned, lost or otherwise discarded fishing gear, as well as to address, as appropriate, the adverse impacts of other human-related activities on the ocean and on marine life, such as ship strikes, underwater noise and invasive alien species;

(h) Promote waste prevention and minimization; develop sustainable consumption and production patterns; adopt the 3Rs — reduce, reuse and recycle — including through incentivizing market-based solutions to reduce waste and its generation, improving mechanisms for environmentally sound waste management, disposal and recycling and developing alternatives such as reusable or recyclable products or products that are biodegradable under natural conditions;

(i) Implement long-term and robust strategies to reduce the use of plastics and microplastics, in particular plastic bags and single-use plastics, including by partnering with stakeholders at relevant levels to address their production, marketing and use;

海洋プラスチック汚染低減のための具体的対策

- ・ 使い捨てプラスチック(特に、レジ袋)の使用規制
- ・ 再使用・リサイクルが容易になるような商品や包装(簡易包装も含む)を生産者や流通業者がとり組むように指導、規制
- ・ 紙や木などのバイオマスの高度利用の促進
- ・ バイオマスベースのプラスチックの利用促進
- ・ リサイクルを促進する社会的なシステムの開発と実装
- ・ 生分解プラスチックの改良と陸上での処理装置での分解促進
- ・ 食品包装へのバイオマスベース生分解性プラスチックの適用とコンポスト化の促進
- ・ 海岸清掃(行政、ボランティア)
- ・ 市民の意識の3R(削減ファースト)意識の啓発

パリ協定(温暖化ガスの実質的排出ゼロ)や他の環境問題対策との整合性

ポストプラスチック社会の将来像


プラスチックの削減が最優先。紙や木の利用も促進して、リサイクルできないプラスチックは極力減らし、それでも残る食品包装等に必要なプラスチックについては石油ベースのプラスチックからバイオマスベースかつ生分解性のプラスチックに置き換え、それらを食品残渣と共にコンポスト化して、農地還元する。

(g) Accelerate actions to prevent and significantly reduce marine pollution of all kinds, particularly from land-based activities, including marine debris, plastics and microplastics, nutrient pollution, untreated wastewater, solid waste discharges, hazardous substances, pollution from ships and abandoned, lost or otherwise discarded fishing gear, as well as to address, as appropriate, the adverse impacts of other human-related activities on the ocean and on marine life, such as ship strikes, underwater noise and invasive alien species;

(h) Promote waste prevention and minimization; develop sustainable consumption and production patterns; adopt the 3Rs — reduce, reuse and recycle — including through incentivizing market-based solutions to reduce waste and its generation, improving mechanisms for environmentally sound waste management, disposal and recycling and developing alternatives such as reusable or recyclable products or products that are biodegradable under natural conditions;


(i) Implement long-term and robust strategies to reduce the use of plastics and microplastics, in particular plastic bags and single-use plastics, including by partnering with stakeholders at relevant levels to address their production, marketing and use;

東京湾流域における窒素の動態



東京湾

1930年



東京湾

赤潮
貧酸素水塊
青潮

現在

トン/年


使い捨てプラスチック削減を最優先にして、 多様なアイデアを取り入れた行政のイニシアティブを



以降補足スライド


5つの主要なプラスチックの生産量割合と密度

ポリエチレン (PE) 29%




密度: 0.90~0.97 g/cm³

ポリスチレン (PS) 7%



1.04 g/cm³


ポリプロピレン (PP)



19%

0.90 g/cm³


ポリエチレンテレフタレート (PET)



7%

1.34~
1.37
g/cm³

ポリ塩化ビニル (PVC)



13%

1.4 g/cm³



マイクロ
プラスチック

5 mm以下の
プラスチック

日本列島から1000km離れた太平洋上で気象庁が採取したマイクロプラスチック。

SCIENTIFIC REPORTS

Microplastic fragments and microbeads in digestive tracts of planktivorous fish from urban coastal waters

Kosuke Tanaka & Hideshige Takada

Laboratory of Organic Geochemistry, Tokyo University of Agriculture and Technology, Fuchu, Tokyo 183-8509, Japan. Correspondence and requests for materials should be addressed to H.T. (email: shige@cc.tuat.ac.jp)

Received: 15 March 2016

Accepted: 13 September 2016


Published: 30 September 2016

マイクロプラスチックはいろいろな起源から供給される

5 mm以下のプラスチック

- プラスチック製品の破片
- 化学繊維
- レジンペレット
- マイクロビーズ(スクラブ)
- メラミンフォームスポンジ

ノニルフェノール：環境ホルモン



プラスチック添加剤

酸化防止剤
剥離剤
静電防止剤

- ・子宮内膜症、乳癌の増加
- ・メスとオスが一緒になった魚（雌雄同体）

堆積物コアを利用して汚染のトレンドを解析：東京におけるコア採取




アジア・アフリカにおけるマイクロプラスチック汚染の歴史変遷の解析




マイクロプラスチック汚染の進行は世界的現象


Thailand (Gulf of Thailand)



Malaysia



South Africa



Vertical profiles of microplastics in sediment cores from Asian and African waters

Regulation of plastic shopping bag : priority option

Polyethylene : high affinity with POPs

Light floating; long travel; thin

Easy to be microplastics

Table 1. Regulation on plastic shopping bags in the world

	Ban		Charge		Tax
	France		Sweden		Denmark
	Italy		Finland		Belgium
	Eritrea		The Netherlands		Luxembourg
	Rwanda		Germany		Iceland
	Bhutan		Australia		Ireland
	Bangladesh		Spain		Kenya
	Cameroon		Botswana		
			South Africa		
			Korea		
			China		
			UK		

わが国における食料供給と窒素循環

川島 博之*

摘 要

わが国における食料供給システムを窒素収支概念より考察した。わが国の農耕地での空中窒素固定量は 1.5×10^5 [ton year⁻¹] 程度と推定される。これに対し 1992 年において、化学肥料として 6.1×10^5 [ton year⁻¹]、漁獲物として 2.6×10^5 [ton year⁻¹]、輸入食料・飼料として 8.3×10^5 [ton year⁻¹] の窒素が食料供給に伴いわが国に流入した。1 億 2 千万人余りの人々が多量の動物性蛋白質を摂取する豊かな食生活は、自然界の窒素循環を大きく改変することにより成立しており、内湾や湖沼の富栄養化の遠因となっている。人間活動が自然界の物質循環に大きな影響を及ぼすことのない社会を構築するためには、農耕地への化学肥料投入を人糞尿の農地還元置き換えること、輸入飼料に依存した畜産に代わり食肉を直接輸入するなどを検討する必要がある。