

1,1-ビス[tert-ブチルペー オキシ]-3,3,5-トリメチルシクロヘキサンの
ラットを用いた 52 週間反復経口投与毒性試験

Table
(図及び群別表)

目 次

	頁
Figure 1 Body weight changes (Male)	1
Figure 2 Body weight changes (Female)	2
Figure 3 Food consumption (Male)	3
Figure 4 Food consumption (Female)	4
Table 1 Clinical signs - Male, Female	5
Table 2 Body weights - Male, Female	39
Table 3 Food consumption - Male, Female	41
Table 4 Urinary findings - Male, Female, 13w	43
Table 5 Urinary findings - Male, Female, 52w	47
Table 6 Hematological findings - Male, Female, 13w	51
Table 7 Hematological findings - Male, Female, 52w	54
Table 8 Biochemical findings - Male, Female, 13w	57
Table 9 Biochemical findings - Male, Female, 52w	60
Table 10 Necropsy findings - Male, Female, 13w	63
Table 11 Necropsy findings - Male, Female, 52w	64
Table 12 Absolute and relative organ weights - Male, Female ,13w	66
Table 13 Absolute and relative organ weights - Male, Female, 52w	70
Table 14 Histopathological findings - Male, Female, 13w	74
Table 15 Histopathological findings - Male, Female, 52w	83

Study No.P030098




Figure 1 Body weight changes (Male)

Study No.P030098




Figure 2 Body weight changes (Female)




Figure 3 Food consumption (Male)

Study No.P030098




Figure 4 Food consumption (Female)