

## Comparison between ICRP Recommendations and Domestic Laws and Regulations

		Occupational exposure		Public exposure	
		ICRP	Laws and regulations concerning the prevention of radiation hazards (Japan)	ICRP	Laws and regulations concerning the prevention of radiation hazards (Japan)
mSv: millisieverts					
<b>Effective dose limits</b>		The average annual dose for the prescribed five years should not exceed 20 mSv. The annual dose for any single year should not exceed 50 mSv. (*1)	The average annual dose for the prescribed five years should not exceed 20 mSv. The annual dose for any single year should not exceed 50 mSv. (*3)	1 mSv/year (In special circumstances, a higher value of effective dose could be allowed in a single year, provided that the average over 5 years does not exceed 1 mSv per year.) (*1)	No dose limit is specified, but doses at the boundaries of the site, including those due to exhaust gas and discharged water, are regulated not to exceed the dose limit of 1 mSv/year. (*3)
<b>Equivalent dose limits</b>	<b>The Lens of the Eye</b>	The average annual dose for the five years should not exceed 20 mSv/year and the annual dose for any single year should not exceed 50 mSv. (*2)	The average annual dose for the five years should not exceed 20 mSv/year and the annual dose for any single year should not exceed 50 mSv. (*3)	15 mSv/year (*1)	—
	<b>Skin</b>	500 mSv/year (*1)	500 mSv/year (*3)	50 mSv/year (*1)	—
	<b>Fingers and toes</b>	500 mSv/year (*1)	—	—	—
<b>Dose limits for female radiation workers</b>		The effective dose of an embryo/a fetus during gestation after reporting pregnancy should not exceed 1 mSv. (*1)	5 mSv/3 months Equivalent dose limit for the abdominal surface after coming to know of pregnancy until delivery: 2 mSv Internal exposure: 1 mSv (*3)	—	—

Source: Prepared based on the following:  
 \*1 2007 Recommendations of the ICRP;  
 \*2 ICRP Publication 118 "ICRP Statement on Tissue Reactions and Early and Late Effects of Radiation in Normal Tissues and Organs - Threshold Doses for Tissue Reactions in a Radiation Protection Context"; and  
 \*3 Japanese laws and regulations concerning the prevention of radiation hazards (as of December 2021)

Present laws and regulations in Japan have not yet completed the incorporation of the 2007 Recommendations of the ICRP, but dose limits specified in the 2007 Recommendations are mostly the same as those in the 1990 Recommendations. Therefore, dose limits in Japan also mostly coincide with those specified in the 2007 Recommendations. Japan has uniquely specified dose limits for female radiation workers (5 mSv per three months).

The ICRP Statement on Tissue Reactions provides recommendations concerning the equivalent dose limit for the lens of the eye under planned occupational exposure. Following the Statement, in Japan, the Radiation Council compiled and provided some insights on “Approach to Radiation Protection of the Lens of the Eye” in 2018, for relevant administrative bodies, and all related laws and regulations (the Regulation on Prevention of Ionizing Radiation Hazards, etc.) were amended in 2021.

Included in this reference material on March 31, 2013

Updated on March 31, 2023

Dose Limits ICRP Recommendations and Responses of the Japanese Government			
	2007 Recommendations of the ICRP		Responses at the time of the accident at Tokyo Electric Power Company (TEPCO)'s Fukushima Daiichi NPS
Occupational exposure	Rescue activities (Volunteers who have obtained the relevant information)	When benefits for other people outweigh the rescuers' risks, dose limits are not applied.	Special Provisions of the Ordinance on Prevention of Ionizing Radiation Hazards (Ministry of Health, Labour and Welfare) The emergency exposure dose limit was temporarily raised to <b>250 mSv</b> from the conventional level of <b>100 mSv</b> (from March 14 to December 16, 2011).
	Other emergency activities	1,000 mSv or 500 mSv	The Ordinance on Prevention of Ionizing Radiation Hazards was partially amended to raise the exceptional emergency dose limit to <b>250 mSv</b> (enforced on April 1, 2016).
Public exposure	Emergency exposure situations	The limit is to be set within the range of <b>20 to 100 mSv/year</b> .	e.g. Standards for evacuation in Deliberate Evacuation Areas: <b>20 mSv/year</b>
	Reconstruction period (Existing exposure situations)	The limit is to be set within the range of <b>1 to 20 mSv/year</b> .	e.g. Additional exposure dose to be achieved in the long term: <b>1 mSv/year</b>

mSv: millisieverts

Source: Prepared based on the 2007 Recommendations of the ICRP and the Special Provisions of the Ordinance on Prevention of Ionizing Radiation Hazards (Ministry of Health, Labour and Welfare: MHLW)

The accident at TEPCO's Fukushima Daiichi NPS occurred while deliberations were continuing over the incorporation of the 2007 Recommendations of the ICRP into domestic laws and regulations.

The accident changed exposure situations, and the idea of reference levels, which had been unfamiliar to Japanese laws and regulations, was adopted for public exposure. In exposure dose control using reference levels, an initial reference level is first set based on the standards for respective exposure situations specified in the 2007 Recommendations of the ICRP so as to ensure that no one receives an unduly high dose. Secondly, if the situation has improved and there is almost no one who receives a high dose exceeding the reference level, a new lower reference level is set as necessary to efficiently achieve exposure dose reduction.

In the meantime, regarding occupational exposure, the emergency dose limit was temporarily raised from 100 mSv to 250 mSv as an exception for an unavoidable case for the purpose of preventing the expansion of the disaster at the NPS. Later, as the work to achieve stable cold shut-down conditions of the reactors was completed, this exceptional measure was abandoned.

Considering the need to develop regulations on the prevention of radiation hazards during emergency work in preparation for any possible nuclear emergencies at nuclear facilities in the future, the Ordinance on Prevention of Ionizing Radiation Hazards was partially amended to raise the exceptional emergency dose limit to 250 mSv. The amended Ordinance was put into force on April 1, 2016.

(Related to p.168 of Vol. 1, "Reduction of Exposure Doses Using Reference Levels")

Included in this reference material on March 31, 2013

Updated on March 31, 2019

Radionuclide	Japan	Codex Alimentarius Commission	EU	US
Radioactive cesium (Bq/kg)	Milk 50 Infant foods 50 General foods 100	Infant foods 1,000 General foods 1,000	Milk 1000 Infant foods 400 General foods 1,250	All foods 1,200
Upper limits for additional doses	1mSv	1mSv	1mSv	5mSv
Assumed percentages of foods containing radioactive materials	50%	10%	10%	30%

- \* The Codex Alimentarius Commission is an intergovernmental body created in 1963 by the Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) for the purpose of protecting consumers' health and ensuring fair-trade practices in the food trade, etc.; The Commission establishes international standards for foods.
- \* Standard limits incorporate effects of the amount of food intake and assumed percentages of foods containing radioactive materials. Therefore, the values are not suitable for inter-comparison.
- \* Indicated standard limits for drinking water are the WHO guidance levels of radioactive materials, which are referred to in respective countries, and standard limits for radioactive materials vary by country due to differences in adopted preconditions. Therefore, the values are not suitable for inter-comparison.

Source: Modified "Food and Radiation Q&A" published by Consumer Affairs Agency

In Japan, the new standard limits for radionuclides in foods were established and were put into force on April 1, 2012. Under the new standard limits, foods are classified into four categories, and the standard limit for drinking water, which people take most frequently, was set at 10 Bq/kg.

The standard limit for general foods was set at 100 Bq/kg. However, for "infant foods" consumed by infants under one year old and for "milk" whose intake by children is extremely high, the standard limit was set at 50 Bq/kg, respectively.


All foods other than infant foods were categorized as general foods based on the idea to minimize gaps in additional doses caused by differences in individuals' eating habits. The value was set with sufficient room to ensure safety no matter what foods people eat as long as radioactive Cs concentrations therein are within the standard limit.

The standard limits vary by country due to differences in annual exposure dose limits based on which the respective countries set their standard limits and in contamination rates in foods, etc. (In Japan, regulation values were set on the safe side based on the annual exposure dose limit of 1 mSv and on the assumption that 50% of general foods and 100% of milk and infant foods are contaminated. On the other hand, the Codex Alimentarius Commission specifies the annual exposure dose limit as 1 mSv and assumes that 10% of foods are contaminated.)

(Related to p.55 of Vol. 2, "Standard Limits Applied from April 2012," p.61 of Vol. 2, "Approach for the Calculation of the Standard Limits (1/2)," and p.62 of Vol. 2, "Approach for the Calculation of the Standard Limits (2/2)")

Included in this reference material on March 31, 2013

Updated on March 31, 2023



There is scientific evidence for the fact that radiation doses of 100 to 200 mSv or over in a relatively short time increase deterministic effects (tissue reactions) and cancer risks. Therefore, in an emergency due to a radiation accident, the initial reference level is set to avoid annual exposure doses of 100 mSv or over in order to prevent serious physical disorders. When the situation improves as the accident is brought under control and there is almost no one who receives a high dose exceeding the initial reference level, a new lower reference level (such as 1 to 20 mSv per year) is set to curb increases in risks of any possible cancer in the future, thereby further promoting exposure dose reduction (p.162 of Vol. 1, “Exposure Situations and Protection Measures”).

As the standard limit in normal times, 1 mSv/year is adopted. As a result, some misunderstand that radiation exposure exceeding 1 mSv per year is dangerous or that they may be exposed to radiation up to that level. However, dose limits do not represent the threshold dividing the safety and the danger.

It is not that radiation exposure up to 1 mSv per year is permissible. Principally, radiation exposure should be reduced as low as practically achievable in light of various circumstances.

(Related to p.117 of Vol. 1, “Relationship between Solid Cancer Deaths and Doses”)

Included in this reference material on March 31, 2013

Updated on March 31, 2021