


- **Distribution of age at the time of radiation exposure of childhood thyroid cancer patients observed in Chernobyl and Fukushima**  
(Among the total number of incidence in respective regions)



Source: Williams D. Eur Thyroid J 2015; 4: 164–173

This figure shows the incidence rates of childhood thyroid cancer by age at the time of radiation exposure (aged 18 or younger), in comparison with those after the Chernobyl accident and those in three years after the accident at TEPCO's Fukushima Daiichi NPS (the percentage in the figure shows the ratio by age, i.e., what percentage the incidence for each age accounts for against the total number of incidence of thyroid cancer in respective regions; the sum of all percentages comes to 100%). The figure shows clear difference in age distribution although an accurate comparison is difficult as thyroid cancer screening in Chernobyl has not been conducted in a uniform manner as in Fukushima and such information as the number of examinees and observation period is not clearly indicated.

Generally speaking, risks of radiation-induced thyroid cancer are higher at younger ages (especially 5 years old or younger). In Chernobyl, it is observed that people exposed to radiation at younger ages have been more likely to develop thyroid cancer. On the other hand, in Fukushima, incidence rates of thyroid cancer among young children have not increased three years after the accident and incidence rates have only increased in tandem with examinees' ages. This tendency is the same as increases observed in incidence rates of ordinary thyroid cancer.

The document by Williams suggests that thyroid cancer detected three years after the accident at Fukushima Daiichi NPS is not attributable to the effects of the radiation exposure due to the accident in light of the facts that daily iodine intake from foods is larger in Japan than in areas around Chernobyl and that the maximum estimated thyroid exposure doses among children is much smaller in Japan (66 mGy in Fukushima and 5,000 mGy in Chernobyl).

(Related to p.131 of Vol. 1, "Comparison between the Chernobyl Accident and the Accident at Tokyo Electric Power Company (TEPCO)'s Fukushima Daiichi NPS (Thyroid Doses)")

Included in this reference material on March 31, 2017