

Exposure Doses from Foods (Example of Calculation)

(e.g.) An adult consumed 0.5 kg of foods containing 100 Bq/kg of Cesium-137

$$\begin{array}{ccccccc}
 100 & \times & 0.5 & \times & 0.013 & = & 0.65 \text{ } \mu\text{Sv} \\
 (\text{Bq/kg}) & & (\text{kg}) & & (\mu\text{Sv/Bq}) & & \\
 & & & & & & = 0.00065 \text{ mSv}
 \end{array}$$

Committed effective dose coefficients ($\mu\text{Sv/Bq}$)



| | Iodine-131 | Cesium-137 |
|------------------|------------|------------|
| Three months old | 0.18 | 0.021 |
| One year old | 0.18 | 0.012 |
| Five years old | 0.10 | 0.0096 |
| Adult | 0.022 | 0.013 |

Bq: becquerels; μSv : microsieverts; mSv: millisieverts

Source: Prepared based on ICRP Publication 119, Compendium of Dose Coefficients based on ICRP Publication 60, 2012, International Commission on Radiological Protection (ICRP)

For example, the dose that an adult who consumed foods containing Cesium-137 will receive is calculated here.

Suppose the person has consumed 0.5 kg of foods containing 100 Bq of Cesium-137 per 1 kg.

The amount of Cesium-137 actually consumed is 50 Bq. This value is multiplied by an effective dose coefficient to calculate committed effective dose (p.56 of Vol. 1, "Committed Effective Doses").

Committed effective dose coefficients are defined in detail for each type of radioactive material, each intake route (inhalation or ingestion), and each age group (p.57 of Vol. 1, "Conversion Factors to Effective Doses").

Included in this reference material on March 31, 2013

Updated on March 31, 2015